



Full wwPDB X-ray Structure Validation Report ⓘ

Jun 22, 2024 – 10:33 AM EDT

PDB ID : 5I5Y
Title : X-RAY CRYSTAL STRUCTURE AT 1.81Å RESOLUTION OF HUMAN MITOCHONDRIAL BRANCHED CHAIN AMINOTRANSFERASE (BCATM) COMPLEXED WITH AN ARYL ACETATE COMPOUND AND AN INTERNAL ALDIMINE LINKED PLP COFACTOR.
Authors : Somers, D.O.
Deposited on : 2016-02-15
Resolution : 1.81 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.37.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.37.1

2 Entry composition

There are 7 unique types of molecules in this entry. The entry contains 7065 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Branched-chain-amino-acid aminotransferase, mitochondrial.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	363	2950	1903	510	517	20	0	7	0
1	B	359	2943	1901	509	513	20	0	9	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP O15382
A	-2	SER	-	expression tag	UNP O15382
A	-1	HIS	-	expression tag	UNP O15382
A	0	MET	-	expression tag	UNP O15382
B	-3	GLY	-	expression tag	UNP O15382
B	-2	SER	-	expression tag	UNP O15382
B	-1	HIS	-	expression tag	UNP O15382
B	0	MET	-	expression tag	UNP O15382

- Molecule 2 is PYRIDOXAL-5'-PHOSPHATE (three-letter code: PLP) (formula: C₈H₁₀NO₆P).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	P		
2	A	1	15	8	1	5	1	0	0
2	B	1	15	8	1	5	1	0	0

- Molecule 3 is {2-[(5-methyl-4-oxo-3,4-dihydrothieno[2,3-d]pyrimidine-6-carbonyl)amino]phenyl}acetic acid (three-letter code: 68D) (formula: C₁₆H₁₃N₃O₄S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
3	A	1	24	16	3	4	1	0	0

Continued on next page...


Continued from previous page...

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
3	B	1	Total	C	N	O	S	0	0
			24	16	3	4	1		

- Molecule 4 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	Cl	0	0
			1	1		
4	B	3	Total	Cl	0	0
			3	3		

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	A	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		
5	B	1	Total	C	O	0	0
			4	2	2		

- Molecule 6 is GLYCEROL (three-letter code: GOL) (formula: $C_3H_8O_3$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	B	1	Total	C	O	0	0
			6	3	3		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	499	Total	O	0	0
			499	499		
7	B	549	Total	O	0	0
			549	549		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Branched-chain-amino-acid aminotransferase, mitochondrial



- Molecule 1: Branched-chain-amino-acid aminotransferase, mitochondrial



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.92Å 109.92Å 59.92Å 90.00° 96.98° 90.00°	Depositor
Resolution (Å)	20.00 – 1.81 29.74 – 1.81	Depositor EDS
% Data completeness (in resolution range)	98.5 (20.00-1.81) 98.4 (29.74-1.81)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.49 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.7.0032	Depositor
R, R_{free}	0.150 , 0.190 0.164 , 0.202	Depositor DCC
R_{free} test set	2733 reflections (4.03%)	wwPDB-VP
Wilson B-factor (Å ²)	19.6	Xtrriage
Anisotropy	0.181	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 54.3	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.017 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7065	wwPDB-VP
Average B, all atoms (Å ²)	24.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.54% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 68D, GOL, CL, EDO, PLP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.41	0/3029	0.66	1/4113 (0.0%)
1	B	0.40	0/3020	0.62	0/4099
All	All	0.41	0/6049	0.64	1/8212 (0.0%)

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	220	ARG	NE-CZ-NH1	5.60	123.10	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2950	0	2967	23	0
1	B	2943	0	2967	10	0
2	A	15	0	6	3	0
2	B	15	0	6	1	0
3	A	24	0	0	0	0
3	B	24	0	0	0	0
4	A	1	0	0	1	0
4	B	3	0	0	2	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	A	12	0	18	1	0
5	B	24	0	36	0	0
6	B	6	0	8	0	0
7	A	499	0	0	9	1
7	B	549	0	0	7	1
All	All	7065	0	6008	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:403:CL:CL	7:A:783:HOH:O	1.94	1.22
4:B:404:CL:CL	7:B:851:HOH:O	1.94	1.17
1:A:179:VAL:CG1	1:A:346:GLN:HE22	1.96	0.78
4:B:405:CL:CL	7:B:620:HOH:O	2.44	0.71
1:A:179:VAL:HG12	1:A:346:GLN:HE22	1.56	0.69
1:B:123:ARG:NH1	7:B:501:HOH:O	2.26	0.68
1:A:220:ARG:NE	7:A:501:HOH:O	2.31	0.62
1:B:105:MET:HE3	7:B:618:HOH:O	2.02	0.59
1:A:315:CYS:SG	1:A:318:CYS:HB2	2.44	0.58
1:A:220:ARG:HG2	1:A:220:ARG:HH11	1.70	0.57
1:A:40[A]:MET:HG3	1:A:54:GLN:NE2	2.23	0.53
1:A:42:GLU:HG3	1:A:162[B]:LEU:CD2	2.39	0.53
1:B:105:MET:HG2	7:B:618:HOH:O	2.08	0.53
1:A:105:MET:CE	7:A:637:HOH:O	2.56	0.53
1:B:40[B]:MET:HG3	1:B:54:GLN:NE2	2.25	0.52
1:B:237:GLU:OE1	2:B:401:PLP:N1	2.43	0.51
1:A:105:MET:HE3	7:A:637:HOH:O	2.10	0.51
1:A:101:LEU:HG	7:A:637:HOH:O	2.11	0.50
1:A:202:LYS:NZ	2:A:401:PLP:O3	2.45	0.49
1:A:173:TYR:HA	5:A:404:EDO:O1	2.14	0.47
1:A:34:PHE:CE2	1:A:168[B]:CYS:SG	3.07	0.47
1:A:179:VAL:CG1	1:A:346:GLN:NE2	2.72	0.47
1:A:19:HIS:CE1	7:A:573:HOH:O	2.68	0.46
1:A:237:GLU:OE1	2:A:401:PLP:N1	2.49	0.46
1:A:105:MET:HG2	7:A:637:HOH:O	2.16	0.46
1:B:101:LEU:CD1	7:B:618:HOH:O	2.63	0.46
1:B:309:PHE:HB3	1:B:342:ILE:HD11	1.97	0.45
1:A:309:PHE:HB3	1:A:342:ILE:HD11	1.99	0.45

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:101:LEU:CD1	7:A:637:HOH:O	2.66	0.43
1:A:31:GLY:HA2	1:A:168[A]:CYS:SG	2.58	0.43
1:B:34:PHE:CE1	1:B:168[B]:CYS:SG	3.12	0.43
1:A:220:ARG:NH2	7:A:506:HOH:O	2.51	0.42
1:A:170[B]:VAL:HG12	1:A:171:GLY:O	2.19	0.42
1:B:242:ASN:O	1:B:310:GLY:HA2	2.20	0.41
1:B:101:LEU:HG	7:B:618:HOH:O	2.20	0.41
1:A:207:TYR:OH	2:A:401:PLP:O3	2.25	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
7:A:593:HOH:O	7:B:591:HOH:O[1_554]	2.10	0.10

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	368/369 (100%)	357 (97%)	11 (3%)	0	100	100
1	B	364/369 (99%)	352 (97%)	12 (3%)	0	100	100
All	All	732/738 (99%)	709 (97%)	23 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	322/319 (101%)	321 (100%)	1 (0%)	92	91
1	B	322/319 (101%)	321 (100%)	1 (0%)	92	91
All	All	644/638 (101%)	642 (100%)	2 (0%)	92	91

All (2) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	247	TRP
1	B	247	TRP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	346	GLN
1	B	234	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 18 ligands modelled in this entry, 4 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	A	404	-	3,3,3	0.49	0	2,2,2	0.39	0
2	PLP	B	401	1	15,15,16	2.78	3 (20%)	20,22,23	1.70	6 (30%)
5	EDO	B	409	-	3,3,3	0.39	0	2,2,2	0.47	0
5	EDO	B	406	-	3,3,3	0.43	0	2,2,2	0.35	0
5	EDO	B	408	-	3,3,3	0.39	0	2,2,2	0.28	0
3	68D	A	402	-	22,26,26	0.97	1 (4%)	23,37,37	1.15	2 (8%)
5	EDO	B	411	-	3,3,3	0.48	0	2,2,2	0.30	0
5	EDO	A	406	-	3,3,3	0.45	0	2,2,2	0.13	0
2	PLP	A	401	1	15,15,16	2.98	3 (20%)	20,22,23	1.70	6 (30%)
5	EDO	A	405	-	3,3,3	0.36	0	2,2,2	0.40	0
3	68D	B	402	-	22,26,26	1.01	2 (9%)	23,37,37	1.53	6 (26%)
6	GOL	B	412	-	5,5,5	0.28	0	5,5,5	0.19	0
5	EDO	B	410	-	3,3,3	0.42	0	2,2,2	0.32	0
5	EDO	B	407	-	3,3,3	0.40	0	2,2,2	0.30	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	A	404	-	-	0/1/1/1	-
2	PLP	B	401	1	-	0/6/6/8	0/1/1/1
5	EDO	B	409	-	-	0/1/1/1	-
5	EDO	B	406	-	-	0/1/1/1	-
5	EDO	B	408	-	-	1/1/1/1	-
3	68D	A	402	-	-	2/8/12/12	0/3/3/3
5	EDO	B	411	-	-	0/1/1/1	-
5	EDO	A	406	-	-	0/1/1/1	-
2	PLP	A	401	1	-	0/6/6/8	0/1/1/1
5	EDO	A	405	-	-	1/1/1/1	-
3	68D	B	402	-	-	2/8/12/12	0/3/3/3
6	GOL	B	412	-	-	2/4/4/4	-
5	EDO	B	410	-	-	1/1/1/1	-
5	EDO	B	407	-	-	1/1/1/1	-

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	401	PLP	C5-C4	7.64	1.48	1.40
2	A	401	PLP	C3-C2	7.44	1.48	1.40
2	B	401	PLP	C5-C4	7.17	1.48	1.40
2	B	401	PLP	C3-C2	6.39	1.47	1.40
2	B	401	PLP	C3-C4	4.06	1.48	1.40
2	A	401	PLP	C3-C4	3.53	1.47	1.40
3	B	402	68D	O23-C22	-2.49	1.22	1.30
3	B	402	68D	C11-C2	2.27	1.43	1.39
3	A	402	68D	C11-C2	2.24	1.43	1.39

All (20) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	402	68D	C20-C21-C22	3.62	120.97	112.99
2	A	401	PLP	C4A-C4-C5	3.53	124.57	120.94
3	B	402	68D	C11-C9-N8	-3.38	112.48	115.49
2	B	401	PLP	C4A-C4-C5	3.38	124.42	120.94
2	A	401	PLP	O4P-C5A-C5	3.15	115.34	109.35
2	B	401	PLP	C3-C4-C5	-2.84	115.67	118.74
3	A	402	68D	C11-C9-N8	-2.76	113.04	115.49
2	A	401	PLP	O2P-P-O4P	-2.72	99.51	106.73
3	B	402	68D	C21-C20-C15	2.59	125.10	121.41
2	B	401	PLP	O2P-P-O4P	-2.56	99.91	106.73
3	A	402	68D	C20-C21-C22	2.44	118.38	112.99
2	B	401	PLP	C6-C5-C4	2.40	120.05	118.16
2	A	401	PLP	C6-N1-C2	2.38	123.58	119.17
2	B	401	PLP	C6-N1-C2	2.29	123.40	119.17
3	B	402	68D	C3-C12-N14	2.28	116.18	113.74
2	A	401	PLP	O3-C3-C2	2.26	122.43	117.49
2	A	401	PLP	C3-C4-C5	-2.16	116.40	118.74
3	B	402	68D	O10-C9-N8	2.10	122.88	120.32
2	B	401	PLP	O3-C3-C2	2.03	121.92	117.49
3	B	402	68D	C21-C20-C19	-2.02	116.59	120.06

There are no chirality outliers.

All (10) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	B	412	GOL	O1-C1-C2-C3
6	B	412	GOL	O1-C1-C2-O2
5	B	410	EDO	O1-C1-C2-O2
3	A	402	68D	C15-C20-C21-C22

Continued on next page...

Continued from previous page...



Mol	Chain	Res	Type	Atoms
3	B	402	68D	C15-C20-C21-C22
5	A	405	EDO	O1-C1-C2-O2
5	B	407	EDO	O1-C1-C2-O2
3	A	402	68D	C19-C20-C21-C22
3	B	402	68D	C19-C20-C21-C22
5	B	408	EDO	O1-C1-C2-O2

There are no ring outliers.

3 monomers are involved in 5 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	404	EDO	1	0
2	B	401	PLP	1	0
2	A	401	PLP	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	363/369 (98%)	0.21	24 (6%) 18 14	11, 21, 51, 83	0
1	B	359/369 (97%)	-0.04	9 (2%) 57 52	11, 19, 35, 78	0
All	All	722/738 (97%)	0.08	33 (4%) 32 27	11, 20, 42, 83	0

All (33) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	27	PRO	7.7
1	A	28	LEU	6.7
1	B	27	PRO	6.3
1	A	176	GLY	6.2
1	A	175	PRO	5.8
1	A	177	GLY	4.9
1	A	173	TYR	4.9
1	B	28	LEU	4.8
1	A	84	LYS	4.5
1	A	25	GLY	4.1
1	A	356	ILE	3.9
1	A	20	LYS	3.5
1	A	174	PHE	3.2
1	A	172	ALA	3.2
1	A	26	GLU	3.2
1	A	179	VAL	3.2
1	B	19	HIS	3.1
1	A	3	SER	3.0
1	A	357	ARG	2.9
1	A	18	PRO	2.8
1	A	29	VAL	2.8
1	A	19	HIS	2.7
1	B	3	SER	2.7
1	B	327	ASP	2.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	16	GLN	2.4
1	A	23	GLY	2.3
1	A	178	SER	2.3
1	A	17	LYS	2.3
1	B	46	LYS	2.2
1	A	180	THR	2.2
1	B	328	ARG	2.1
1	B	45	ASP	2.1
1	B	20	LYS	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	EDO	A	404	4/4	0.73	0.16	35,37,39,41	0
5	EDO	B	406	4/4	0.80	0.15	40,40,42,43	0
6	GOL	B	412	6/6	0.85	0.32	54,55,57,61	0
5	EDO	B	410	4/4	0.87	0.15	36,36,37,39	0
5	EDO	A	406	4/4	0.87	0.16	28,31,34,39	0
5	EDO	B	408	4/4	0.90	0.14	24,28,30,34	0
5	EDO	A	405	4/4	0.91	0.12	33,33,34,34	0
5	EDO	B	409	4/4	0.92	0.14	41,42,43,47	0
5	EDO	B	407	4/4	0.94	0.27	42,45,45,45	0
3	68D	A	402	24/24	0.96	0.09	17,20,22,22	0
3	68D	B	402	24/24	0.97	0.08	14,15,16,17	0
2	PLP	A	401	15/16	0.98	0.09	14,17,19,21	0
2	PLP	B	401	15/16	0.98	0.09	11,13,17,17	0
5	EDO	B	411	4/4	0.98	0.07	21,22,23,26	0


Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	CL	B	405	1/1	0.98	0.05	32,32,32,32	0
4	CL	B	404	1/1	0.99	0.05	24,24,24,24	0
4	CL	A	403	1/1	0.99	0.04	25,25,25,25	0
4	CL	B	403	1/1	0.99	0.13	14,14,14,14	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





6.5 Other polymers [i](#)

There are no such residues in this entry.