



Full wwPDB X-ray Structure Validation Report ⓘ

May 23, 2020 – 03:11 am BST

PDB ID : 1I7R
Title : CRYSTAL STRUCTURE OF CLASS I MHC A2 IN COMPLEX WITH PEPTIDE P1058
Authors : Busslep, J.; Zhao, R.; Loftus, D.; Appella, E.; Collins, E.J.
Deposited on : 2001-03-10
Resolution : 2.20 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtrriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION







The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4898 (2.20-2.20)
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)
RSRZ outliers	127900	4800 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	275	 4% 83% 13% •
1	D	275	 6% 82% 14% •
2	B	100	 % 82% 16% ••
2	E	100	 3% 76% 20% •
3	C	9	 11% 78% 22%
3	F	9	 22% 78% 22%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 6427 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			
1	D	275	Total	C	N	O	S	0	0	0
			2247	1403	409	426	9			

- Molecule 2 is a protein called BETA-2-MICROGLOBULIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			
2	E	100	Total	C	N	O	S	0	0	0
			837	533	141	159	4			

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	0	MET	-	CLONING ARTIFACT	UNP P01884
E	0	MET	-	CLONING ARTIFACT	UNP P01884

- Molecule 3 is a protein called 9 RESIDUE PEPTIDE.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	9	Total	C	N	O	0	0	0
			77	57	9	11			
3	F	9	Total	C	N	O	0	0	0
			76	57	9	10			


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	34	Total 34	O 34	0	0
4	B	22	Total 22	O 22	0	0
4	C	1	Total 1	O 1	0	0
4	D	32	Total 32	O 32	0	0
4	E	17	Total 17	O 17	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN




- Molecule 1: HLA CLASS I HISTOCOMPATIBILITY ANTIGEN, A-2 ALPHA CHAIN





- Molecule 2: BETA-2-MICROGLOBULIN





- Molecule 2: BETA-2-MICROGLOBULIN



- Molecule 3: 9 RESIDUE PEPTIDE



● Molecule 3: 9 RESIDUE PEPTIDE



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	50.86Å 63.82Å 75.87Å 82.09° 76.20° 78.13°	Depositor
Resolution (Å)	30.00 – 2.20 31.09 – 2.20	Depositor EDS
% Data completeness (in resolution range)	96.7 (30.00-2.20) 96.8 (31.09-2.20)	Depositor EDS
R_{merge}	0.08	Depositor
R_{sym}	0.08	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.90 (at 2.20Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, R_{free}	0.264 , 0.307 0.242 , 0.282	Depositor DCC
R_{free} test set	2252 reflections (5.08%)	wwPDB-VP
Wilson B-factor (Å ²)	18.9	Xtrriage
Anisotropy	0.006	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 48.2	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.90	EDS
Total number of atoms	6427	wwPDB-VP
Average B, all atoms (Å ²)	19.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.23% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.47	0/2312	1.29	20/3137 (0.6%)
1	D	0.47	0/2312	1.25	20/3137 (0.6%)
2	B	0.48	0/860	1.19	2/1162 (0.2%)
2	E	0.49	0/860	1.13	1/1162 (0.1%)
3	C	0.60	0/82	1.03	0/110
3	F	0.59	0/81	1.12	0/110
All	All	0.48	0/6507	1.24	43/8818 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
2	B	0	1
2	E	0	1
All	All	0	2

There are no bond length outliers.

All (43) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	44	ARG	NE-CZ-NH1	-14.71	112.94	120.30
1	A	273	ARG	NE-CZ-NH1	-13.66	113.47	120.30
1	D	44	ARG	NE-CZ-NH2	11.40	126.00	120.30
2	B	97	ARG	NE-CZ-NH2	11.13	125.87	120.30
1	D	44	ARG	NE-CZ-NH1	-10.47	115.06	120.30
1	A	234	ARG	NE-CZ-NH1	-9.06	115.77	120.30
1	A	273	ARG	NE-CZ-NH2	8.73	124.67	120.30
1	A	157	ARG	NE-CZ-NH2	-8.41	116.09	120.30
1	A	44	ARG	NE-CZ-NH2	7.64	124.12	120.30
1	A	97	ARG	CD-NE-CZ	7.64	134.30	123.60

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	234	ARG	NE-CZ-NH2	7.24	123.92	120.30
1	D	97	ARG	CD-NE-CZ	7.01	133.41	123.60
1	A	219	ARG	NE-CZ-NH2	-6.94	116.83	120.30
1	D	219	ARG	NE-CZ-NH1	6.74	123.67	120.30
2	B	97	ARG	NE-CZ-NH1	-6.74	116.93	120.30
1	A	35	ARG	CD-NE-CZ	6.51	132.71	123.60
1	D	116	TYR	CA-CB-CG	6.46	125.68	113.40
1	D	35	ARG	CD-NE-CZ	6.26	132.36	123.60
2	E	97	ARG	NE-CZ-NH1	-6.20	117.20	120.30
1	D	219	ARG	CD-NE-CZ	6.16	132.23	123.60
1	D	157	ARG	NE-CZ-NH1	6.00	123.30	120.30
1	A	122	ASP	CB-CG-OD2	5.95	123.66	118.30
1	D	256	ARG	CD-NE-CZ	5.94	131.91	123.60
1	A	116	TYR	CA-CB-CG	5.85	124.51	113.40
1	A	219	ARG	CD-NE-CZ	5.83	131.76	123.60
1	D	223	ASP	CB-CG-OD1	5.80	123.52	118.30
1	A	17	ARG	NE-CZ-NH1	5.76	123.18	120.30
1	A	84	TYR	CA-CB-CG	5.54	123.93	113.40
1	D	84	TYR	CA-CB-CG	5.52	123.89	113.40
1	D	230	LEU	CA-CB-CG	5.50	127.95	115.30
1	A	201	LEU	CA-CB-CG	5.42	127.78	115.30
1	D	44	ARG	CD-NE-CZ	-5.40	116.04	123.60
1	D	61	ASP	CB-CG-OD1	5.39	123.15	118.30
1	D	99	TYR	CB-CG-CD1	-5.36	117.78	121.00
1	A	170	ARG	NE-CZ-NH2	-5.36	117.62	120.30
1	A	244	TRP	CA-CB-CG	5.27	123.72	113.70
1	D	244	TRP	CA-CB-CG	5.27	123.72	113.70
1	A	223	ASP	CB-CG-OD1	5.25	123.02	118.30
1	D	77	ASP	CB-CG-OD1	5.21	122.99	118.30
1	D	122	ASP	CB-CG-OD2	5.20	122.98	118.30
1	D	170	ARG	NE-CZ-NH2	-5.10	117.75	120.30
1	A	37	ASP	CB-CG-OD1	5.10	122.89	118.30
1	D	223	ASP	CA-CB-CG	5.00	124.41	113.40

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
2	B	0	MET	Mainchain
2	E	0	MET	Mainchain

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2247	0	2096	22	0
1	D	2247	0	2096	22	0
2	B	837	0	803	10	0
2	E	837	0	803	12	0
3	C	77	0	71	1	0
3	F	76	0	71	1	0
4	A	34	0	0	1	0
4	B	22	0	0	1	0
4	C	1	0	0	0	0
4	D	32	0	0	1	0
4	E	17	0	0	1	0
All	All	6427	0	5940	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:0:MET:O	2:B:0:MET:HG2	1.67	0.92
2:B:73:THR:HG22	2:B:76:ASP:H	1.48	0.79
2:E:73:THR:HG22	2:E:76:ASP:H	1.49	0.77
1:D:19:GLU:HB3	1:D:20:PRO:HD2	1.71	0.73
1:A:19:GLU:HB3	1:A:20:PRO:HD2	1.70	0.73
2:B:85:VAL:HG13	4:B:100:HOH:O	1.97	0.65
2:B:12:ARG:HB3	2:B:21:ASN:HD21	1.62	0.65
1:D:99:TYR:OH	3:F:3:PRO:HD2	1.98	0.62
1:A:99:TYR:OH	3:C:3:PRO:HD2	2.02	0.59
1:A:253:GLN:NE2	1:A:256:ARG:HH11	2.00	0.59
1:A:263:HIS:HD2	1:A:265:GLY:H	1.51	0.58
2:B:13:HIS:HD2	2:B:21:ASN:ND2	2.02	0.58
1:D:263:HIS:HD2	1:D:265:GLY:H	1.52	0.57
2:E:12:ARG:HB3	2:E:21:ASN:HD21	1.70	0.56
1:A:44:ARG:NH1	4:A:278:HOH:O	2.39	0.56
1:D:263:HIS:CD2	1:D:265:GLY:H	2.24	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:0:MET:O	2:B:0:MET:CG	2.48	0.54
1:D:274:TRP:O	1:D:275:GLU:HB3	2.07	0.54
2:E:13:HIS:HD2	2:E:21:ASN:ND2	2.07	0.53
1:A:263:HIS:CD2	1:A:265:GLY:H	2.26	0.53
2:B:22:PHE:CZ	2:B:69:GLU:HG2	2.44	0.52
1:D:9:PHE:HE2	1:D:99:TYR:CE2	2.27	0.52
1:D:234:ARG:HD2	1:D:242:GLN:HB2	1.92	0.51
2:B:21:ASN:CG	2:B:22:PHE:H	2.14	0.51
1:D:253:GLN:NE2	1:D:256:ARG:HH11	2.09	0.50
1:D:28:VAL:HG11	1:D:179:LEU:HD13	1.92	0.50
1:D:111:ARG:HD3	1:D:113:TYR:CZ	2.47	0.50
2:E:21:ASN:CG	2:E:22:PHE:H	2.16	0.50
1:A:111:ARG:HD3	1:A:113:TYR:CZ	2.49	0.48
1:A:28:VAL:HG11	1:A:179:LEU:HD13	1.96	0.48
1:A:9:PHE:HE2	1:A:99:TYR:CE2	2.31	0.48
2:E:85:VAL:HG13	4:E:101:HOH:O	2.14	0.47
1:D:117:ALA:HB2	2:E:60:TRP:CE2	2.49	0.47
1:A:98:MET:HB3	1:A:98:MET:HE2	1.68	0.47
1:D:98:MET:HE2	1:D:98:MET:HB3	1.68	0.47
2:E:21:ASN:ND2	2:E:22:PHE:H	2.14	0.46
1:D:44:ARG:HD3	1:D:44:ARG:HH11	1.44	0.46
1:D:127:LYS:HD2	1:D:132:SER:HB2	1.98	0.45
1:A:182:THR:CG2	1:A:265:GLY:HA2	2.47	0.45
2:B:21:ASN:ND2	2:B:22:PHE:H	2.15	0.45
1:A:273:ARG:HD2	1:A:273:ARG:HH11	1.48	0.45
1:A:44:ARG:HD3	1:A:44:ARG:HH11	1.47	0.44
1:D:15:PRO:HG2	1:D:91:GLY:O	2.18	0.44
2:E:83:ASN:OD1	2:E:90:PRO:HG3	2.17	0.44
1:D:16:GLY:O	1:D:17:ARG:NE	2.50	0.44
2:E:22:PHE:CZ	2:E:69:GLU:HG2	2.53	0.44
1:A:234:ARG:HD2	1:A:242:GLN:HB2	2.00	0.44
1:A:111:ARG:HD3	1:A:113:TYR:OH	2.18	0.43
1:D:253:GLN:HE21	1:D:256:ARG:HD3	1.83	0.43
2:E:51:HIS:HA	2:E:65:LEU:O	2.18	0.43
1:D:44:ARG:NH1	4:D:279:HOH:O	2.52	0.42
1:A:97:ARG:HD2	1:A:116:TYR:CZ	2.54	0.42
1:A:127:LYS:HD2	1:A:132:SER:HB2	2.02	0.42
1:A:197:HIS:ND1	1:A:197:HIS:N	2.68	0.42
1:D:55:GLU:OE2	1:D:170:ARG:NH2	2.52	0.42
1:A:117:ALA:HB2	2:B:60:TRP:CE2	2.54	0.42
1:D:184:ALA:HB2	1:D:265:GLY:O	2.21	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:E:73:THR:CG2	2:E:75:LYS:HG2	2.50	0.41
1:D:111:ARG:HD3	1:D:113:TYR:OH	2.20	0.41
1:D:97:ARG:HD2	1:D:116:TYR:CZ	2.56	0.41
2:E:25:CYS:HB2	2:E:39:LEU:HD21	2.03	0.41
1:A:182:THR:HG23	1:A:265:GLY:HA2	2.02	0.40
1:A:257:TYR:O	1:A:273:ARG:HG3	2.22	0.40
1:A:15:PRO:HG2	1:A:91:GLY:O	2.21	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	273/275 (99%)	266 (97%)	7 (3%)	0	100	100
1	D	273/275 (99%)	267 (98%)	6 (2%)	0	100	100
2	B	98/100 (98%)	95 (97%)	3 (3%)	0	100	100
2	E	98/100 (98%)	96 (98%)	2 (2%)	0	100	100
3	C	7/9 (78%)	7 (100%)	0	0	100	100
3	F	7/9 (78%)	7 (100%)	0	0	100	100
All	All	756/768 (98%)	738 (98%)	18 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	231/231 (100%)	217 (94%)	14 (6%)	18	21
1	D	231/231 (100%)	215 (93%)	16 (7%)	15	16
2	B	95/95 (100%)	86 (90%)	9 (10%)	8	8
2	E	95/95 (100%)	85 (90%)	10 (10%)	7	6
3	C	7/7 (100%)	6 (86%)	1 (14%)	3	2
3	F	7/7 (100%)	6 (86%)	1 (14%)	3	2
All	All	666/666 (100%)	615 (92%)	51 (8%)	13	13

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	11	SER
1	A	14	ARG
1	A	98	MET
1	A	105	SER
1	A	132	SER
1	A	182	THR
1	A	196	ASP
1	A	201	LEU
1	A	207	SER
1	A	223	ASP
1	A	225	THR
1	A	230	LEU
1	A	234	ARG
1	A	268	LYS
2	B	0	MET
2	B	34	ASP
2	B	48	LYS
2	B	58	LYS
2	B	70	PHE
2	B	73	THR
2	B	75	LYS
2	B	83	ASN
2	B	88	SER
3	C	6	PHE
1	D	11	SER
1	D	14	ARG
1	D	35	ARG
1	D	98	MET

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	105	SER
1	D	108	ARG
1	D	132	SER
1	D	182	THR
1	D	186	LYS
1	D	196	ASP
1	D	201	LEU
1	D	207	SER
1	D	223	ASP
1	D	225	THR
1	D	230	LEU
1	D	234	ARG
2	E	0	MET
2	E	34	ASP
2	E	48	LYS
2	E	58	LYS
2	E	70	PHE
2	E	73	THR
2	E	75	LYS
2	E	83	ASN
2	E	88	SER
2	E	91	LYS
3	F	6	PHE

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	174	ASN
1	A	253	GLN
1	A	263	HIS
1	D	174	ASN
1	D	253	GLN
1	D	263	HIS
2	E	2	GLN
2	E	13	HIS
2	E	21	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	275/275 (100%)	0.14	11 (4%) 38 36	8, 18, 31, 40	0
1	D	275/275 (100%)	0.27	16 (5%) 23 22	9, 20, 33, 41	0
2	B	100/100 (100%)	-0.05	1 (1%) 82 81	8, 15, 29, 33	0
2	E	100/100 (100%)	0.11	3 (3%) 50 48	8, 16, 31, 35	0
3	C	9/9 (100%)	0.95	1 (11%) 5 4	15, 22, 29, 29	1 (11%)
3	F	9/9 (100%)	1.11	2 (22%) 0 0	15, 22, 28, 31	1 (11%)
All	All	768/768 (100%)	0.18	34 (4%) 34 32	8, 18, 32, 41	2 (0%)

All (34) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	99	TYR	4.2
1	D	86	ASN	4.0
1	A	86	ASN	3.9
1	A	18	GLY	3.9
1	D	196	ASP	3.7
1	D	223	ASP	3.7
3	F	5	PHE	3.6
1	D	195	SER	3.5
1	D	16	GLY	3.4
1	A	17	ARG	3.4
1	A	267	PRO	3.3
1	D	17	ARG	3.3
3	C	5	PHE	3.3
1	D	194	VAL	3.3
1	A	99	TYR	3.1
1	A	16	GLY	3.1
1	A	268	LYS	3.1
2	B	0	MET	2.9
1	D	193	ALA	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
2	E	0	MET	2.8
1	A	54	GLN	2.7
1	A	196	ASP	2.7
1	D	88	SER	2.5
1	D	192	HIS	2.5
1	D	18	GLY	2.5
1	D	197	HIS	2.5
1	D	226	GLN	2.4
1	D	131	ARG	2.4
1	A	1	GLY	2.4
2	E	47	GLU	2.3
3	F	9	LEU	2.3
1	A	195	SER	2.3
2	E	48	LYS	2.1
1	D	105	SER	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.