



Full wwPDB X-ray Structure Validation Report ⓘ

Feb 6, 2023 – 04:14 PM JST

PDB ID : 8I7L
Title : Crystal structure of indoleamine 2,3-dioxygenase 1 (IDO1) complexed with a novel inhibitor
Authors : Li, K.; Liu, W.; Dong, X.
Deposited on : 2023-02-01
Resolution : 2.80 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.32.1
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.32.1


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3140 (2.80-2.80)
Clashscore	141614	3569 (2.80-2.80)
Ramachandran outliers	138981	3498 (2.80-2.80)
Sidechain outliers	138945	3500 (2.80-2.80)
RSRZ outliers	127900	3078 (2.80-2.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	403	
1	B	403	

2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 5825 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Indoleamine 2,3-dioxygenase 1.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	357	2840	1833	478	512	17	0	0	0
1	B	359	2876	1857	490	512	17	0	0	0

- Molecule 2 is THIOSULFATE (three-letter code: THJ) (formula: O₃S₂).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	3	2	0	0

- Molecule 3 is 1-[3-[(4-chloranyl-2-fluoranyl-phenyl)carbamoylamino]-4-[cyclohexyl(2-methylpropyl)amino]phenyl]pyrrole-2-carboxylic acid (three-letter code: OIH) (formula: C₂₈H₃₂ClFN₄O₃) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf		
			Total	C	Cl	F	N			O	
3	A	1	Total	37	28	1	1	4	3	0	0
3	B	1	Total	37	28	1	1	4	3	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
4	A	18	Total	18	0	0
4	B	12	Total	12	0	0

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	87.11Å 97.11Å 128.45Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	48.55 – 2.80 48.55 – 2.80	Depositor EDS
% Data completeness (in resolution range)	87.6 (48.55-2.80) 87.7 (48.55-2.80)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.02 (at 2.81Å)	Xtrriage
Refinement program	PHENIX 1.16-3549	Depositor
R, R_{free}	0.270 , 0.293 0.271 , 0.292	Depositor DCC
R_{free} test set	1206 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	87.8	Xtrriage
Anisotropy	0.252	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.35 , 64.9	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5825	wwPDB-VP
Average B, all atoms (Å ²)	107.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: THJ, OIH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.29	0/2906	0.54	0/3932
1	B	0.31	0/2942	0.58	2/3975 (0.1%)
All	All	0.30	0/5848	0.56	2/7907 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed($^{\circ}$)	Ideal($^{\circ}$)
1	B	297	ARG	NE-CZ-NH1	-6.83	116.88	120.30
1	B	270	PHE	N-CA-CB	-6.07	99.67	110.60

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	12	SER	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2840	0	2834	19	0
1	B	2876	0	2901	21	0
2	A	5	0	0	0	0
3	A	37	0	0	0	0
3	B	37	0	0	1	0
4	A	18	0	0	0	0
4	B	12	0	0	0	0
All	All	5825	0	5735	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:290:GLN:HA	1:B:297:ARG:HH22	1.02	1.18
1:A:290:GLN:HA	1:B:297:ARG:NH2	1.80	0.95
1:A:290:GLN:CA	1:B:297:ARG:HH22	1.90	0.84
1:A:50:ILE:HD11	1:A:89:ALA:HB1	1.67	0.77
1:A:317:ARG:NH2	1:A:332:TYR:OH	2.25	0.70
1:B:317:ARG:NH2	1:B:398:SER:O	2.26	0.67
1:B:271:GLN:NE2	1:B:292:LEU:O	2.31	0.59
1:A:383:ASP:N	1:A:383:ASP:OD1	2.35	0.59
1:B:383:ASP:N	1:B:383:ASP:OD1	2.35	0.58
1:A:193:ARG:NE	1:A:325:ASP:OD2	2.35	0.57
1:B:321:LEU:HD21	1:B:400:LEU:HD22	1.90	0.54
1:B:279:ILE:HD13	1:B:395:THR:HG23	1.91	0.53
1:A:343:ARG:HE	1:A:391:VAL:HG12	1.73	0.53
1:A:321:LEU:HD21	1:A:400:LEU:HD22	1.91	0.52
1:B:193:ARG:NE	1:B:325:ASP:OD2	2.40	0.50
1:A:317:ARG:HH12	1:A:398:SER:C	2.17	0.48
1:B:384:LEU:H	1:B:384:LEU:HD23	1.78	0.47
1:A:279:ILE:HD13	1:A:395:THR:HG23	1.95	0.47
1:B:35:PHE:CE2	1:B:65:LEU:HD22	2.49	0.47
1:A:232:ILE:HD11	1:A:233:TYR:CZ	2.51	0.46
1:A:84:GLY:O	1:A:88:MET:HG2	2.17	0.45
1:A:317:ARG:NH1	1:A:398:SER:O	2.44	0.45
1:A:361:GLN:HG3	1:A:362:PRO:HD2	1.99	0.45
1:B:281:GLN:HB3	1:B:292:LEU:HD12	1.99	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:14:GLU:H	1:B:14:GLU:HG2	1.50	0.43
1:B:277:LEU:HD11	1:B:335:CYS:HB3	2.00	0.43
1:B:181:ILE:N	1:B:182:PRO:HD2	2.34	0.43
1:A:355:LEU:HD23	1:A:355:LEU:HA	1.89	0.42
1:B:136:LYS:HD3	1:B:139:PRO:HA	2.00	0.42
1:B:12:SER:O	1:B:14:GLU:N	2.52	0.42
1:B:84:GLY:O	1:B:88:MET:HG2	2.19	0.42
1:A:46:LEU:O	1:A:50:ILE:HG12	2.20	0.41
1:A:80:ARG:HH21	1:A:125:VAL:HG22	1.85	0.41
1:B:115:SER:HB3	1:B:120:LEU:O	2.21	0.41
1:B:355:LEU:HD23	1:B:355:LEU:HA	1.86	0.41
1:A:123:ILE:O	1:A:125:VAL:HG13	2.21	0.41
1:B:262:GLY:HA3	3:B:501:OIH:CL1	2.58	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	347/403 (86%)	337 (97%)	10 (3%)	0	100	100
1	B	349/403 (87%)	339 (97%)	9 (3%)	1 (0%)	41	72
All	All	696/806 (86%)	676 (97%)	19 (3%)	1 (0%)	51	81

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	13	LYS

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	311/348 (89%)	304 (98%)	7 (2%)	50	82
1	B	316/348 (91%)	312 (99%)	4 (1%)	69	91
All	All	627/696 (90%)	616 (98%)	11 (2%)	59	86

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	14	GLU
1	A	29	GLN
1	A	232	ILE
1	A	247	LEU
1	A	270	PHE
1	A	287	HIS
1	A	383	ASP
1	B	14	GLU
1	B	287	HIS
1	B	290	GLN
1	B	383	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	THJ	A	501	-	2,4,4	2.01	1 (50%)	2,6,6	1.05	0
3	OIH	B	501	-	39,40,40	2.62	12 (30%)	49,56,56	1.85	10 (20%)
3	OIH	A	502	-	39,40,40	2.59	9 (23%)	49,56,56	1.80	11 (22%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OIH	B	501	-	-	7/27/36/36	0/4/4/4
3	OIH	A	502	-	-	11/27/36/36	0/4/4/4

All (22) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	OIH	C26-CL1	-8.95	1.55	1.74
3	B	501	OIH	C26-CL1	-8.93	1.55	1.74
3	A	502	OIH	C7-N2	-6.95	1.32	1.38
3	B	501	OIH	C7-N2	-6.79	1.32	1.38
3	A	502	OIH	C22-N3	5.68	1.49	1.37
3	B	501	OIH	C22-N3	5.53	1.49	1.37
3	B	501	OIH	C22-N4	5.11	1.48	1.37
3	B	501	OIH	C10-C11	4.78	1.56	1.51
3	A	502	OIH	C22-N4	4.75	1.47	1.37

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	502	OIH	C10-C11	4.43	1.55	1.51
3	A	502	OIH	C3-N1	3.75	1.53	1.41
3	B	501	OIH	C3-N1	3.66	1.52	1.41
3	B	501	OIH	C23-N4	3.53	1.48	1.41
3	A	502	OIH	C23-N4	3.40	1.48	1.41
3	A	502	OIH	C13-N1	2.56	1.51	1.47
3	A	502	OIH	C6-N2	2.54	1.48	1.44
3	B	501	OIH	C13-N1	2.48	1.51	1.47
3	B	501	OIH	C6-N2	2.36	1.47	1.44
3	B	501	OIH	C4-N3	2.18	1.45	1.41
3	B	501	OIH	C13-C14	2.06	1.58	1.52
2	A	501	THJ	O2-S1	2.03	1.51	1.45
3	B	501	OIH	O3-C22	-2.02	1.19	1.23

All (21) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	502	OIH	C17-C12-N1	-6.32	104.32	112.80
3	B	501	OIH	C25-C24-C23	-5.33	118.42	123.50
3	B	501	OIH	C26-C25-C24	4.97	121.14	117.48
3	A	502	OIH	C25-C24-C23	-4.01	119.68	123.50
3	B	501	OIH	C23-N4-C22	-3.75	117.06	125.39
3	A	502	OIH	C18-C17-C12	3.72	118.23	110.82
3	B	501	OIH	C21-C12-N1	-3.69	107.85	112.80
3	A	502	OIH	C26-C25-C24	3.59	120.12	117.48
3	B	501	OIH	N4-C22-N3	3.50	118.61	112.49
3	B	501	OIH	C28-C23-C24	3.40	120.60	117.17
3	A	502	OIH	C28-C23-C24	3.20	120.40	117.17
3	A	502	OIH	C21-C12-N1	-2.85	108.98	112.80
3	A	502	OIH	N4-C22-N3	2.58	117.00	112.49
3	B	501	OIH	O3-C22-N4	-2.57	119.28	123.62
3	A	502	OIH	O3-C22-N4	-2.52	119.37	123.62
3	B	501	OIH	C20-C21-C12	2.48	115.75	110.82
3	A	502	OIH	C23-N4-C22	-2.37	120.13	125.39
3	A	502	OIH	C21-C12-C17	2.29	116.46	111.19
3	B	501	OIH	F1-C24-C23	2.25	120.32	117.50
3	A	502	OIH	C20-C21-C12	2.05	114.89	110.82
3	B	501	OIH	O2-C11-O1	-2.02	118.87	123.35

There are no chirality outliers.

All (18) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	A	502	OIH	N2-C10-C11-O1
3	A	502	OIH	N2-C10-C11-O2
3	A	502	OIH	C21-C12-N1-C13
3	B	501	OIH	N2-C10-C11-O1
3	B	501	OIH	N2-C10-C11-O2
3	B	501	OIH	C21-C12-N1-C13
3	A	502	OIH	C24-C23-N4-C22
3	B	501	OIH	C24-C23-N4-C22
3	A	502	OIH	N1-C13-C14-C15
3	B	501	OIH	C4-C3-N1-C13
3	A	502	OIH	N1-C13-C14-C16
3	A	502	OIH	C21-C12-N1-C3
3	B	501	OIH	C17-C12-N1-C13
3	B	501	OIH	C2-C3-N1-C13
3	A	502	OIH	C2-C3-N1-C13
3	A	502	OIH	C9-C10-C11-O1
3	A	502	OIH	C4-C3-N1-C13
3	A	502	OIH	C28-C23-N4-C22


There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	501	OIH	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	357/403 (88%)	0.85	39 (10%) 5 3	66, 108, 148, 192	0
1	B	359/403 (89%)	0.67	29 (8%) 12 6	65, 103, 143, 183	0
All	All	716/806 (88%)	0.76	68 (9%) 8 4	65, 105, 144, 192	0

All (68) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	69	HIS	6.6
1	A	287	HIS	6.3
1	A	289	ALA	5.6
1	B	362	PRO	5.5
1	A	101	LYS	4.8
1	B	281	GLN	4.5
1	B	287	HIS	4.4
1	A	294	ASP	4.0
1	A	290	GLN	3.9
1	B	360	GLN	3.6
1	A	292	LEU	3.6
1	A	280	GLN	3.6
1	A	362	PRO	3.5
1	A	293	GLN	3.5
1	A	361	GLN	3.4
1	A	251	GLY	3.4
1	B	239	GLY	3.4
1	A	291	PHE	3.4
1	A	197	LEU	3.2
1	A	295	MET	3.1
1	B	114	LEU	3.0
1	A	288	ALA	3.0
1	A	207	LEU	3.0
1	B	65	LEU	3.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	99	VAL	3.0
1	A	137	LYS	3.0
1	A	383	ASP	2.9
1	B	308	CYS	2.8
1	B	140	ASN	2.8
1	B	297	ARG	2.7
1	B	361	GLN	2.7
1	B	354	ILE	2.7
1	A	297	ARG	2.7
1	B	39	TRP	2.7
1	B	386	ASN	2.6
1	A	32	LEU	2.6
1	A	270	PHE	2.6
1	B	241	PRO	2.5
1	A	27	ASN	2.5
1	A	139	PRO	2.4
1	A	92	TRP	2.4
1	A	70	LEU	2.4
1	B	355	LEU	2.4
1	A	66	SER	2.4
1	A	272	CYS	2.4
1	B	356	ILE	2.4
1	A	96	HIS	2.3
1	A	271	GLN	2.3
1	A	138	ASP	2.3
1	A	151	LEU	2.3
1	B	35	PHE	2.2
1	A	50	ILE	2.2
1	B	293	GLN	2.2
1	B	288	ALA	2.1
1	B	289	ALA	2.1
1	A	319	PHE	2.1
1	A	143	LEU	2.1
1	A	59	VAL	2.1
1	B	215	HIS	2.1
1	B	67	ILE	2.1
1	B	330	GLU	2.1
1	A	107	ILE	2.0
1	B	69	HIS	2.0
1	A	57	GLU	2.0
1	B	59	VAL	2.0
1	B	107	ILE	2.0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	247	LEU	2.0
1	B	200	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)




In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
3	OIH	B	501	37/37	0.81	0.33	90,97,117,119	0
3	OIH	A	502	37/37	0.88	0.33	72,86,95,100	0
2	THJ	A	501	5/5	0.90	0.20	145,146,147,150	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.




Electron density around OIH B 501:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



Electron density around OIH A 502:

$2mF_o-DF_c$ (at 0.7 rmsd) in gray
 mF_o-DF_c (at 3 rmsd) in purple (negative)
and green (positive)



6.5 Other polymers [i](#)

There are no such residues in this entry.