



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 28, 2024 – 07:39 AM EDT

PDB ID : 3I8N  
Title : A domain of a conserved functionally known protein from *Vibrio parahaemolyticus* RIMD 2210633.  
Authors : Tan, K.; Tesar, C.; Bearden, J.; Joachimiak, A.; Midwest Center for Structural Genomics (MCSG)  
Deposited on : 2009-07-09  
Resolution : 2.15 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 2022.3.0, CSD as543be (2022)  
Xtriage (Phenix) : 1.20.1  
EDS : 3.0  
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)  
CCP4 : 9.0.003 (Gargrove)  
Density-Fitness : 1.0.11  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.39


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.15 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	164625	3336 (2.16-2.12)
Clashscore	180529	3585 (2.16-2.12)
Ramachandran outliers	177936	3554 (2.16-2.12)
Sidechain outliers	177891	3553 (2.16-2.12)
RSRZ outliers	164620	3337 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	130	
1	B	130	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 2120 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called uncharacterized protein VP2912.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	124	998	642	169	181	6	0	2	0
1	B	126	1009	648	172	183	6	0	1	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	186	SER	-	expression tag	UNP Q87KR4
A	187	ASN	-	expression tag	UNP Q87KR4
A	188	ALA	-	expression tag	UNP Q87KR4
B	186	SER	-	expression tag	UNP Q87KR4
B	187	ASN	-	expression tag	UNP Q87KR4
B	188	ALA	-	expression tag	UNP Q87KR4

- Molecule 2 is FORMIC ACID (three-letter code: FMT) (formula: CH<sub>2</sub>O<sub>2</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			3	1	2		
2	B	1	Total	C	O	0	0
			3	1	2		


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	50	Total	O	0	0
			50	50		
3	B	57	Total	O	0	0
			57	57		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: uncharacterized protein VP2912



- Molecule 1: uncharacterized protein VP2912



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 65 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.30Å 69.30Å 255.14Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	49.04 – 2.15 49.04 – 2.15	Depositor EDS
% Data completeness (in resolution range)	98.3 (49.04-2.15) 98.1 (49.04-2.15)	Depositor EDS
$R_{merge}$	0.15	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.05 (at 2.14Å)	Xtrriage
Refinement program	PHENIX (phenix.refine)	Depositor
R, $R_{free}$	0.219 , 0.265 0.223 , 0.263	Depositor DCC
$R_{free}$ test set	1056 reflections (5.14%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	46.4	Xtrriage
Anisotropy	0.590	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 60.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	2120	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	67.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.62% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: FMT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.33	0/1017	0.54	0/1367
1	B	0.31	0/1025	0.49	0/1381
All	All	0.32	0/2042	0.51	0/2748

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	998	0	1029	41	0
1	B	1009	0	1036	33	0
2	A	3	0	1	0	0
2	B	3	0	1	0	0
3	A	50	0	0	2	0
3	B	57	0	0	1	0
All	All	2120	0	2067	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:200:ARG:HH22	1:A:226:ARG:HD3	1.12	1.12
1:B:194:THR:HA	1:B:197:MSE:HE3	1.02	0.99
1:B:194:THR:CA	1:B:197:MSE:HE3	1.92	0.98
1:B:194:THR:HA	1:B:197:MSE:CE	1.97	0.94
1:B:210:MSE:HG3	1:B:214:GLU:HG3	1.59	0.84
1:B:197:MSE:HE1	1:B:292:VAL:HG11	1.62	0.79
1:A:200:ARG:NH2	1:A:226:ARG:HD3	1.94	0.79
1:B:211:THR:OG1	1:B:214:GLU:HG2	1.84	0.78
1:B:206[A]:VAL:HG13	1:B:210:MSE:HE3	1.73	0.70
1:B:197:MSE:CE	1:B:292:VAL:HG11	2.22	0.69
1:B:257:LYS:HD2	1:B:258:GLN:H	1.59	0.67
1:B:295:GLU:HG3	3:B:23:HOH:O	1.94	0.67
1:B:206[A]:VAL:CG1	1:B:210:MSE:HE3	2.28	0.64
1:A:242:HIS:HE1	1:A:288:GLN:HG3	1.63	0.64
1:B:289:LEU:HG	1:B:302:LEU:HD11	1.77	0.64
1:A:243:ARG:HD2	3:A:40:HOH:O	1.98	0.63
1:A:242:HIS:CE1	1:A:288:GLN:HG3	2.34	0.62
1:A:194:THR:HA	1:A:197:MSE:O	2.00	0.62
1:A:225:SER:OG	1:A:226:ARG:HD2	1.99	0.61
1:B:197:MSE:HE1	1:B:292:VAL:CG1	2.31	0.60
1:A:211:THR:HG22	1:A:214:GLU:H	1.64	0.60
1:A:270:LEU:HD13	1:B:265:PRO:HB3	1.84	0.59
1:A:200:ARG:HH22	1:A:226:ARG:CD	2.02	0.59
1:A:211:THR:CG2	1:A:214:GLU:H	2.15	0.59
1:B:197:MSE:HE2	1:B:292:VAL:HG21	1.84	0.58
1:A:211:THR:HG22	1:A:214:GLU:HG3	1.87	0.57
1:B:257:LYS:CD	1:B:258:GLN:H	2.17	0.57
1:A:243:ARG:H	1:A:243:ARG:CZ	2.18	0.56
1:B:249:MSE:HE2	1:B:262:VAL:HG13	1.87	0.56
1:A:275:LEU:HD23	1:A:311:HIS:HD2	1.71	0.56
1:B:257:LYS:HG3	1:B:261:ALA:HB2	1.87	0.56
1:B:249:MSE:HE2	1:B:262:VAL:CG1	2.36	0.56
1:B:208:ALA:HB2	1:B:229:VAL:CG1	2.36	0.56
1:B:271:ASN:HD22	1:B:271:ASN:N	2.05	0.55
1:A:285:HIS:NE2	1:B:286:ARG:HD3	2.25	0.52
1:B:257:LYS:HD2	1:B:258:GLN:N	2.25	0.51
1:A:211:THR:HG22	1:A:214:GLU:CG	2.40	0.51
1:B:271:ASN:HD22	1:B:271:ASN:H	1.57	0.51
1:A:275:LEU:HD23	1:A:311:HIS:CD2	2.46	0.50
1:A:266:ILE:HG12	1:A:291:LEU:HD22	1.93	0.50
1:A:211:THR:HG23	1:A:213:ASN:N	2.26	0.50

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:225:SER:O	1:A:227:PRO:HD3	2.13	0.49
1:A:302:LEU:C	1:A:302:LEU:HD12	2.33	0.48
1:A:304:THR:O	1:A:308:ILE:HG12	2.13	0.48
1:A:245:GLU:O	1:A:249:MSE:HG3	2.15	0.47
1:A:196:VAL:HG22	1:A:311:HIS:CE1	2.50	0.47
1:A:285:HIS:CD2	1:B:286:ARG:HD2	2.49	0.47
1:A:225:SER:C	1:A:227:PRO:HD3	2.34	0.47
1:B:200:ARG:NH2	1:B:235:ASP:HB2	2.31	0.46
1:A:192:PRO:HA	1:A:195:GLN:NE2	2.30	0.46
1:A:265:PRO:HG3	1:B:281:GLN:OE1	2.16	0.46
1:A:308:ILE:HG22	1:A:312:LEU:HD22	1.97	0.46
1:A:288:GLN:HG2	3:A:21:HOH:O	2.16	0.46
1:A:289:LEU:C	1:A:289:LEU:HD12	2.37	0.45
1:A:226:ARG:CZ	1:A:304:THR:HG21	2.47	0.44
1:B:259:LEU:HD12	1:B:259:LEU:HA	1.78	0.44
1:B:222:THR:HB	1:B:223:PRO:HD2	1.99	0.44
1:A:211:THR:HG23	1:A:213:ASN:H	1.83	0.43
1:B:283:MSE:HE2	1:B:283:MSE:HB3	1.88	0.43
1:A:249:MSE:HE1	1:A:262:VAL:CG1	2.50	0.42
1:A:239:GLY:HA3	1:A:263:MSE:SE	2.69	0.42
1:A:204:PHE:CE2	1:A:227:PRO:HB3	2.55	0.42
1:B:206[A]:VAL:HG13	1:B:210:MSE:CE	2.48	0.42
1:A:312:LEU:HD12	1:A:312:LEU:HA	1.85	0.41
1:B:238:ILE:HG13	1:B:263:MSE:SE	2.71	0.41
1:A:243:ARG:H	1:A:243:ARG:NH2	2.19	0.41
1:A:228:LEU:HD13	1:A:237:ILE:HG21	2.04	0.40
1:A:232:GLU:HG3	1:B:296:TYR:CE1	2.56	0.40
1:A:210:MSE:HB3	1:A:259:LEU:HD22	2.04	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [\(i\)](#)

### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	124/130 (95%)	117 (94%)	7 (6%)	0	100	100
1	B	125/130 (96%)	121 (97%)	4 (3%)	0	100	100
All	All	249/260 (96%)	238 (96%)	11 (4%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	115/112 (103%)	103 (90%)	12 (10%)	5	2
1	B	116/112 (104%)	104 (90%)	12 (10%)	6	2
All	All	231/224 (103%)	207 (90%)	24 (10%)	5	2

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	211	THR
1	A	217	ASP
1	A	226	ARG
1	A	259	LEU
1	A	270	LEU
1	A	275	LEU
1	A	289	LEU
1	A	291	LEU
1	A	295	GLU
1	A	300	LEU
1	A	312	LEU
1	A	313	VAL
1	B	195	GLN
1	B	200	ARG
1	B	205	ARG
1	B	216	LEU
1	B	233	GLN
1	B	246	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	256	GLN
1	B	259	LEU
1	B	271	ASN
1	B	295	GLU
1	B	310	GLU
1	B	312	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (16) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	195	GLN
1	A	213	ASN
1	A	233	GLN
1	A	236	ASN
1	A	242	HIS
1	A	250	GLN
1	A	267	GLN
1	A	281	GLN
1	A	311	HIS
1	B	236	ASN
1	B	250	GLN
1	B	251	GLN
1	B	256	GLN
1	B	258	GLN
1	B	271	ASN
1	B	272	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	FMT	B	2	-	2,2,2	0.74	0	1,1,1	0.21	0
2	FMT	A	1	-	2,2,2	0.68	0	1,1,1	0.15	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	118/130 (90%)	0.54	6 (5%) 34 40	31, 60, 93, 143	2 (1%)
1	B	120/130 (92%)	0.72	11 (9%) 16 20	39, 66, 103, 184	1 (0%)
All	All	238/260 (91%)	0.63	17 (7%) 23 28	31, 62, 102, 184	3 (1%)

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	194	THR	4.7
1	A	231	SER	4.0
1	B	309	PHE	3.5
1	A	193	VAL	3.5
1	B	188	ALA	3.3
1	B	313	VAL	3.2
1	A	192	PRO	2.9
1	A	226	ARG	2.9
1	B	236	ASN	2.4
1	B	272	ASN	2.3
1	A	314	GLY	2.3
1	B	190	ASP	2.3
1	B	195	GLN	2.3
1	B	233	GLN	2.2
1	B	261	ALA	2.1
1	B	307	ASP	2.1
1	B	191	VAL	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	FMT	A	1	3/3	0.75	0.25	58,58,76,77	0
2	FMT	B	2	3/3	0.85	0.14	81,81,87,88	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.