



# Full wwPDB X-ray Structure Validation Report ⓘ

Jun 13, 2024 – 06:17 AM EDT

PDB ID : 4I82  
Title : Crystal Structure of Hypothetical Thioesterase Protein SP\_1851 from *Streptococcus pneumoniae* TIGR4  
Authors : Khandokar, Y.B.; Forwood, J.K.  
Deposited on : 2012-12-02  
Resolution : 2.50 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.20.1  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4661 (2.50-2.50)
Clashscore	141614	5346 (2.50-2.50)
Ramachandran outliers	138981	5231 (2.50-2.50)
Sidechain outliers	138945	5233 (2.50-2.50)
RSRZ outliers	127900	4559 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	137	 4% 64% 21% • 12%
1	B	137	 2% 71% 15% • 12%

## 2 Entry composition

There is only 1 type of molecule in this entry. The entry contains 1852 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Putative uncharacterized protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	121	Total	C	N	O	S	35	0	0
			926	581	156	182	7			
1	B	121	Total	C	N	O	S	7	0	0
			926	581	156	182	7			


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q97NZ8
A	-1	ASN	-	expression tag	UNP Q97NZ8
A	0	ALA	-	expression tag	UNP Q97NZ8
B	-2	SER	-	expression tag	UNP Q97NZ8
B	-1	ASN	-	expression tag	UNP Q97NZ8
B	0	ALA	-	expression tag	UNP Q97NZ8


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Putative uncharacterized protein



- Molecule 1: Putative uncharacterized protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 2 2 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.26Å 118.47Å 87.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	24.77 – 2.50 24.77 – 2.50	Depositor EDS
% Data completeness (in resolution range)	93.5 (24.77-2.50) 93.6 (24.77-2.50)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.23 (at 2.50Å)	Xtrriage
Refinement program	REFMAC 5.7.0029	Depositor
R, $R_{free}$	0.238 , 0.286 0.243 , 0.293	Depositor DCC
$R_{free}$ test set	374 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	30.6	Xtrriage
Anisotropy	0.226	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.37 , 34.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.91	EDS
Total number of atoms	1852	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 40.16 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 2.8498e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	1.24	8/939 (0.9%)	1.12	10/1270 (0.8%)
1	B	0.81	1/939 (0.1%)	0.87	2/1270 (0.2%)
All	All	1.05	9/1878 (0.5%)	1.00	12/2540 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (9) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	7	ASP	CB-CG	-19.23	1.11	1.51
1	A	21	ARG	CG-CD	18.13	1.97	1.51
1	B	21	ARG	CD-NE	-16.92	1.17	1.46
1	A	19	LYS	CB-CG	-12.31	1.19	1.52
1	A	16	GLU	CG-CD	-9.75	1.37	1.51
1	A	77	LYS	CB-CG	-8.69	1.29	1.52
1	A	62	LEU	CA-CB	-6.34	1.39	1.53
1	A	125	GLN	CB-CG	6.18	1.69	1.52
1	A	114	CYS	CB-SG	-5.21	1.73	1.81

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	90	GLU	OE1-CD-OE2	-18.26	101.39	123.30
1	A	111	ARG	CD-NE-CZ	10.77	138.67	123.60
1	A	90	GLU	CG-CD-OE1	8.87	136.04	118.30
1	A	19	LYS	CA-CB-CG	8.19	131.41	113.40
1	B	21	ARG	CD-NE-CZ	6.54	132.76	123.60

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	21	ARG	CB-CG-CD	-6.44	94.86	111.60
1	A	62	LEU	CA-CB-CG	6.08	129.28	115.30
1	A	114	CYS	N-CA-CB	-5.49	100.72	110.60
1	A	77	LYS	CA-CB-CG	5.42	125.32	113.40
1	A	19	LYS	CB-CG-CD	5.33	125.47	111.60
1	A	125	GLN	CA-CB-CG	-5.17	102.01	113.40
1	B	21	ARG	CG-CD-NE	5.01	122.32	111.80

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	113	VAL	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	926	0	911	19	0
1	B	926	0	911	19	0
All	All	1852	0	1822	38	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 11.

All (38) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:CYS:HB2	1:A:101:MET:CE	2.16	0.76
1:A:105:ILE:HB	1:A:114:CYS:HB3	1.68	0.76
1:B:58:VAL:CG1	1:B:101:MET:HE2	2.21	0.70
1:A:13:GLU:OE2	1:A:34:SER:OG	2.09	0.70
1:A:24:HIS:HD2	1:A:90:GLU:HB3	1.57	0.70
1:B:58:VAL:CG1	1:B:101:MET:CE	2.74	0.65
1:B:58:VAL:HG11	1:B:101:MET:CE	2.27	0.63

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:96:ARG:HG3	1:B:96:ARG:HH11	1.64	0.62
1:A:24:HIS:CD2	1:A:90:GLU:HB3	2.35	0.61
1:A:43:HIS:HD2	1:A:45:GLY:H	1.50	0.58
1:A:99:CYS:CB	1:A:101:MET:HE1	2.34	0.57
1:B:17:ILE:HD13	1:B:20:MET:HE2	1.87	0.55
1:A:99:CYS:HB2	1:A:101:MET:HE1	1.84	0.55
1:B:58:VAL:HG12	1:B:101:MET:HE2	1.89	0.55
1:B:7:ASP:OD1	1:B:7:ASP:N	2.40	0.54
1:A:105:ILE:HD12	1:A:114:CYS:SG	2.49	0.53
1:B:96:ARG:HH11	1:B:96:ARG:CG	2.21	0.53
1:B:91:CYS:HB2	1:B:101:MET:HE1	1.89	0.53
1:A:38:TYR:CD1	1:A:38:TYR:N	2.78	0.52
1:A:11:ALA:O	1:A:12:PHE:HB2	2.11	0.51
1:B:58:VAL:HG11	1:B:101:MET:HE3	1.91	0.50
1:B:43:HIS:HD2	1:B:45:GLY:H	1.59	0.50
1:A:43:HIS:CD2	1:A:45:GLY:H	2.31	0.49
1:B:36:LEU:HA	1:B:41:ASN:O	2.12	0.49
1:B:43:HIS:HD2	1:B:45:GLY:N	2.12	0.48
1:A:99:CYS:CB	1:A:101:MET:CE	2.86	0.48
1:A:73:ILE:O	1:A:73:ILE:HG23	2.14	0.47
1:B:58:VAL:HG12	1:B:101:MET:CE	2.44	0.47
1:A:51:CYS:SG	1:A:105:ILE:HD11	2.56	0.46
1:A:99:CYS:HB2	1:A:101:MET:HE2	1.98	0.45
1:A:81:LEU:O	1:A:82:ASP:HB2	2.16	0.45
1:B:26:VAL:HG22	1:B:88:LYS:HG3	1.99	0.44
1:A:28:THR:HA	1:A:85:LEU:O	2.18	0.43
1:B:36:LEU:HD11	1:B:81:LEU:HD13	2.01	0.42
1:B:96:ARG:CG	1:B:96:ARG:NH1	2.82	0.42
1:B:28:THR:HG22	1:B:86:THR:HB	2.02	0.42
1:A:108:GLN:NE2	1:A:108:GLN:H	2.18	0.41
1:B:43:HIS:CD2	1:B:45:GLY:H	2.38	0.41

There are no symmetry-related clashes.

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/137 (87%)	111 (93%)	8 (7%)	0	100	100
1	B	119/137 (87%)	113 (95%)	6 (5%)	0	100	100
All	All	238/274 (87%)	224 (94%)	14 (6%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	104/119 (87%)	96 (92%)	8 (8%)	13	25
1	B	104/119 (87%)	93 (89%)	11 (11%)	6	13
All	All	208/238 (87%)	189 (91%)	19 (9%)	9	18

All (19) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	20	MET
1	A	38	TYR
1	A	68	THR
1	A	96	ARG
1	A	108	GLN
1	A	115	LYS
1	A	122	VAL
1	A	125	GLN
1	B	7	ASP
1	B	38	TYR
1	B	55	SER
1	B	67	VAL
1	B	80	LYS
1	B	86	THR
1	B	88	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	96	ARG
1	B	108	GLN
1	B	119	THR
1	B	122	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	24	HIS
1	A	43	HIS
1	A	94	GLN
1	A	108	GLN
1	B	43	HIS
1	B	94	GLN
1	B	108	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	121/137 (88%)	0.18	5 (4%) 37 40	17, 28, 49, 79	10 (8%)
1	B	121/137 (88%)	-0.03	3 (2%) 57 61	15, 23, 39, 58	2 (1%)
All	All	242/274 (88%)	0.07	8 (3%) 46 50	15, 26, 44, 79	12 (4%)

All (8) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	5	HIS	3.8
1	B	38	TYR	3.3
1	B	109	GLU	3.2
1	B	49	THR	2.7
1	A	110	GLY	2.3
1	A	109	GLU	2.1
1	A	49	THR	2.0
1	A	22	ASP	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.