



Full wwPDB X-ray Structure Validation Report ⓘ

Aug 20, 2023 – 08:06 PM EDT

PDB ID : 2IF8
Title : Crystal structure of Inositol Phosphate Multikinase Ipk2 in complex with ADP and Mn²⁺ from *S. cerevisiae*
Authors : Holmes, W.; Jogl, G.
Deposited on : 2006-09-20
Resolution : 2.40 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : 1.13
EDS : 2.35
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.35


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION



The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3907 (2.40-2.40)
Clashscore	141614	4398 (2.40-2.40)
Ramachandran outliers	138981	4318 (2.40-2.40)
Sidechain outliers	138945	4319 (2.40-2.40)
RSRZ outliers	127900	3811 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	363	
1	B	363	

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 4568 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Inositol polyphosphate multikinase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	255	2083	1338	335	400	10	0	0	0
1	B	256	2091	1342	336	403	10	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	356	LEU	-	cloning artifact	UNP P07250
A	357	GLU	-	cloning artifact	UNP P07250
A	358	HIS	-	expression tag	UNP P07250
A	359	HIS	-	expression tag	UNP P07250
A	360	HIS	-	expression tag	UNP P07250
A	361	HIS	-	expression tag	UNP P07250
A	362	HIS	-	expression tag	UNP P07250
A	363	HIS	-	expression tag	UNP P07250
B	356	LEU	-	cloning artifact	UNP P07250
B	357	GLU	-	cloning artifact	UNP P07250
B	358	HIS	-	expression tag	UNP P07250
B	359	HIS	-	expression tag	UNP P07250
B	360	HIS	-	expression tag	UNP P07250
B	361	HIS	-	expression tag	UNP P07250
B	362	HIS	-	expression tag	UNP P07250
B	363	HIS	-	expression tag	UNP P07250


- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	1	Total	Mn	0	0
			1	1		

- Molecule 3 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Ca 1 1	0	0

- Molecule 4 is ADENOSINE-5'-DIPHOSPHATE (three-letter code: ADP) (formula: $C_{10}H_{15}N_5O_{10}P_2$).



4 Data and refinement statistics

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	185.00Å 185.00Å 50.00Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	30.00 – 2.40 26.98 – 2.40	Depositor EDS
% Data completeness (in resolution range)	98.5 (30.00-2.40) 98.6 (26.98-2.40)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.80 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.193 , 0.246 0.190 , 0.238	Depositor DCC
R_{free} test set	1922 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	41.0	Xtrriage
Anisotropy	0.299	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 55.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.47$, $\langle L^2 \rangle = 0.29$	Xtrriage
Estimated twinning fraction	0.047 for h,-h-k,-l	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	4568	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.16% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: CA, ADP, MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	0.68	1/2122 (0.0%)	0.73	0/2854
1	B	0.58	0/2130	0.70	1/2865 (0.0%)
All	All	0.63	1/4252 (0.0%)	0.71	1/5719 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	168	CYS	CB-SG	-7.57	1.69	1.82

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	38	LEU	CA-CB-CG	5.38	127.68	115.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2083	0	2069	25	0
1	B	2091	0	2073	38	0
2	A	1	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
4	A	27	0	12	0	0
5	A	212	0	0	4	1
5	B	153	0	0	9	0
All	All	4568	0	4154	62	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (62) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:25:ASP:HB2	5:B:496:HOH:O	1.28	1.25
1:B:197:PHE:HB3	5:B:487:HOH:O	1.38	1.23
1:A:317:SER:N	5:A:561:HOH:O	1.87	1.05
1:A:147:LYS:HB3	5:A:520:HOH:O	1.70	0.91
1:A:174:LYS:HE2	1:A:189:GLU:OE1	1.95	0.67
1:B:62:LEU:HD13	1:B:247:MET:CE	2.24	0.67
1:B:269:ARG:NH2	1:B:317:SER:O	2.29	0.65
1:B:62:LEU:HD13	1:B:247:MET:HE1	1.79	0.63
1:A:62:LEU:HD13	1:A:247:MET:HE1	1.82	0.62
1:B:156:GLU:HG3	5:B:392:HOH:O	1.99	0.62
1:B:267:PRO:O	1:B:271:GLU:HG2	2.01	0.60
1:B:282:ASP:O	5:B:404:HOH:O	2.16	0.60
1:B:44:ILE:HG22	1:B:63:CYS:HB3	1.85	0.58
1:B:246:THR:HG22	1:B:247:MET:HE2	1.87	0.57
1:A:346:GLU:HG2	1:B:353:MET:CE	2.37	0.55
1:B:66:MET:HG2	1:B:326:PHE:CZ	2.41	0.55
1:A:134:LEU:HD11	1:A:259:LEU:HG	1.89	0.55
1:B:167:ILE:O	1:B:203:GLY:HA3	2.07	0.54
1:B:200:LYS:O	1:B:204:ARG:HB2	2.07	0.54
1:B:175:ASN:HB3	1:B:178:VAL:HB	1.89	0.53
1:B:72:VAL:HG22	1:B:73:LEU:H	1.74	0.53
1:A:163:LEU:HB3	1:A:347:THR:HG21	1.91	0.52
1:B:197:PHE:CB	5:B:487:HOH:O	2.21	0.52
1:A:37:GLU:HG2	1:A:115:LEU:HD11	1.91	0.51
1:A:62:LEU:HD13	1:A:247:MET:CE	2.40	0.51
1:B:72:VAL:HG22	1:B:73:LEU:N	2.26	0.51
1:B:29:ILE:HG22	1:B:117:LEU:HB2	1.91	0.51
1:A:145:LEU:HD23	1:A:148:ARG:HD2	1.93	0.50
1:B:62:LEU:HD13	1:B:247:MET:HE3	1.93	0.49
1:B:354:LYS:HG3	5:B:429:HOH:O	2.14	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:317:SER:N	5:B:493:HOH:O	2.46	0.48
1:A:190:GLU:HB2	1:A:192:ASP:OD1	2.15	0.47
1:B:357:GLU:O	5:B:482:HOH:O	2.20	0.47
1:A:215:ILE:O	1:A:218:TYR:HB3	2.14	0.46
1:B:237:LEU:HD23	1:B:356:LEU:CD1	2.46	0.46
1:B:178:VAL:HA	1:B:181:GLN:OE1	2.16	0.45
1:B:237:LEU:HD23	1:B:356:LEU:HD11	1.98	0.44
1:B:135:GLY:HA2	1:B:340:ASN:O	2.17	0.44
1:B:138:LEU:HB2	1:B:151:MET:HB3	1.98	0.44
1:B:134:LEU:HD11	1:B:259:LEU:HG	2.00	0.43
1:A:189:GLU:HG2	1:A:190:GLU:H	1.83	0.43
1:A:190:GLU:HG2	5:A:558:HOH:O	2.18	0.43
1:A:278:LYS:HE2	1:A:282:ASP:OD2	2.19	0.43
1:A:167:ILE:O	1:A:203:GLY:HA3	2.19	0.42
1:B:146:GLU:H	1:B:146:GLU:HG2	1.62	0.42
1:B:216:GLU:HG3	1:B:355:PHE:CE1	2.54	0.42
1:A:67:PRO:HG3	1:A:120:LEU:HD11	2.02	0.42
1:B:183:SER:HB3	1:B:185:GLU:OE1	2.20	0.41
1:B:38:LEU:HD11	1:B:72:VAL:HB	2.02	0.41
1:A:332:THR:HB	1:A:335:LYS:HD2	2.02	0.41
1:B:134:LEU:HD13	1:B:341:VAL:HG13	2.01	0.41
1:A:331:ILE:O	1:A:331:ILE:HG13	2.19	0.41
1:B:163:LEU:HB3	1:B:347:THR:HG21	2.03	0.41
1:A:121:LEU:HD23	1:A:262:ILE:HD13	2.03	0.41
1:A:178:VAL:HG12	1:A:179:LEU:N	2.36	0.41
1:B:192:ASP:CG	1:B:193:SER:N	2.74	0.41
1:B:337:TYR:CD1	1:B:337:TYR:C	2.95	0.41
1:A:31:LYS:O	1:A:114:TYR:HA	2.22	0.40
1:A:41:TYR:CD2	1:A:69:TYR:HB2	2.56	0.40
1:A:317:SER:N	5:A:536:HOH:O	2.54	0.40
1:B:133:LYS:NZ	5:B:478:HOH:O	2.50	0.40
1:A:44:ILE:HD12	1:A:62:LEU:HD23	2.04	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
5:A:562:HOH:O	5:A:590:HOH:O[4_665]	1.90	0.30

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	247/363 (68%)	241 (98%)	3 (1%)	3 (1%)	13	19
1	B	248/363 (68%)	241 (97%)	5 (2%)	2 (1%)	19	29
All	All	495/726 (68%)	482 (97%)	8 (2%)	5 (1%)	15	23

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	178	VAL
1	B	179	LEU
1	B	325	ASP
1	A	177	SER
1	A	325	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	234/327 (72%)	222 (95%)	12 (5%)	24	39
1	B	235/327 (72%)	217 (92%)	18 (8%)	13	20
All	All	469/654 (72%)	439 (94%)	30 (6%)	17	28

All (30) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	118	GLU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	141	SER
1	A	147	LYS
1	A	174	LYS
1	A	179	LEU
1	A	183	SER
1	A	209	GLN
1	A	234	LYS
1	A	282	ASP
1	A	339	GLU
1	A	342	ILE
1	A	343	GLU
1	B	25	ASP
1	B	27	LEU
1	B	31	LYS
1	B	45	GLN
1	B	73	LEU
1	B	113	GLN
1	B	141	SER
1	B	146	GLU
1	B	147	LYS
1	B	148	ARG
1	B	180	ASN
1	B	190	GLU
1	B	201	LEU
1	B	204	ARG
1	B	325	ASP
1	B	331	ILE
1	B	337	TYR
1	B	342	ILE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	45	GLN
1	B	45	GLN
1	B	223	HIS

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 2 are monoatomic - leaving 1 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	ADP	A	401	2	24,29,29	1.05	2 (8%)	29,45,45	1.20	3 (10%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	ADP	A	401	2	-	6/12/32/32	0/3/3/3

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	401	ADP	C5-C4	2.84	1.48	1.40
4	A	401	ADP	O4'-C1'	2.39	1.44	1.41

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	401	ADP	N3-C2-N1	-3.22	123.64	128.68
4	A	401	ADP	C3'-C2'-C1'	2.68	105.01	100.98

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	A	401	ADP	C4-C5-N7	-2.16	107.15	109.40

There are no chirality outliers.


All (6) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
4	A	401	ADP	C5'-O5'-PA-O2A
4	A	401	ADP	O4'-C4'-C5'-O5'
4	A	401	ADP	C3'-C4'-C5'-O5'
4	A	401	ADP	C5'-O5'-PA-O3A
4	A	401	ADP	C5'-O5'-PA-O1A
4	A	401	ADP	PB-O3A-PA-O1A

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	255/363 (70%)	0.07	21 (8%) 11 10	29, 39, 62, 97	0
1	B	256/363 (70%)	0.50	26 (10%) 6 6	34, 49, 77, 96	0
All	All	511/726 (70%)	0.28	47 (9%) 9 8	29, 45, 73, 97	0

All (47) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	111	SER	9.3
1	B	285	ILE	7.5
1	A	286	ASP	6.9
1	B	286	ASP	6.8
1	B	191	ALA	6.2
1	B	145	LEU	6.1
1	A	285	ILE	5.5
1	B	283	ASP	5.4
1	B	284	PHE	5.3
1	A	191	ALA	5.2
1	A	111	SER	4.6
1	B	190	GLU	4.5
1	B	192	ASP	4.0
1	B	112	LYS	3.9
1	A	147	LYS	3.8
1	B	144	SER	3.8
1	A	283	ASP	3.5
1	B	150	ARG	3.5
1	B	147	LYS	3.5
1	B	75	GLU	3.3
1	B	146	GLU	3.3
1	B	142	LYS	3.0
1	B	282	ASP	2.9
1	B	197	PHE	2.9

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	284	PHE	2.8
1	A	260	LEU	2.8
1	A	324	ILE	2.7
1	A	259	LEU	2.7
1	B	268	GLU	2.7
1	B	143	ALA	2.7
1	B	193	SER	2.6
1	B	201	LEU	2.5
1	B	334	GLY	2.5
1	A	192	ASP	2.5
1	B	156	GLU	2.5
1	A	282	ASP	2.4
1	A	261	PHE	2.4
1	A	145	LEU	2.4
1	A	258	SER	2.3
1	A	190	GLU	2.3
1	A	262	ILE	2.2
1	B	321	MET	2.2
1	A	112	LYS	2.2
1	A	130	LEU	2.2
1	B	45	GLN	2.1
1	A	146	GLU	2.1
1	A	134	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	ADP	A	401	27/27	0.83	0.18	82,86,95,96	0
2	MN	A	402	1/1	0.89	0.10	112,112,112,112	0
3	CA	A	403	1/1	0.99	0.12	50,50,50,50	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.