



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 10:46 am GMT

PDB ID : 2IG6
Title : Crystal structure of a nimc/nima family protein (ca_c2569) from clostridium acetobutylicum at 1.80 Å resolution
Authors : Joint Center for Structural Genomics (JCSG)
Deposited on : 2006-09-22
Resolution : 1.80 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.80 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	5253 (1.80-1.80)
Clashscore	122126	6077 (1.80-1.80)
Ramachandran outliers	120053	6011 (1.80-1.80)
Sidechain outliers	120020	6010 (1.80-1.80)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	150	88% 7% 5%
1	B	150	81% 10% 7%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	UNL	A	340	-	-	X	-

2 Entry composition i

There are 6 unique types of molecules in this entry. The entry contains 2631 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called NimC/NimA family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	143	1149	736	195	210	4	4	0	3	0
1	B	139	1139	732	188	211	4	4	0	6	0

There are 46 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-18	MSE	-	LEADER SEQUENCE	UNP Q97G05
A	-17	GLY	-	LEADER SEQUENCE	UNP Q97G05
A	-16	SER	-	LEADER SEQUENCE	UNP Q97G05
A	-15	ASP	-	LEADER SEQUENCE	UNP Q97G05
A	-14	LYS	-	LEADER SEQUENCE	UNP Q97G05
A	-13	ILE	-	LEADER SEQUENCE	UNP Q97G05
A	-12	HIS	-	LEADER SEQUENCE	UNP Q97G05
A	-11	HIS	-	LEADER SEQUENCE	UNP Q97G05
A	-10	HIS	-	LEADER SEQUENCE	UNP Q97G05
A	-9	HIS	-	LEADER SEQUENCE	UNP Q97G05
A	-8	HIS	-	LEADER SEQUENCE	UNP Q97G05
A	-7	HIS	-	LEADER SEQUENCE	UNP Q97G05
A	-6	GLU	-	LEADER SEQUENCE	UNP Q97G05
A	-5	ASN	-	LEADER SEQUENCE	UNP Q97G05
A	-4	LEU	-	LEADER SEQUENCE	UNP Q97G05
A	-3	TYR	-	LEADER SEQUENCE	UNP Q97G05
A	-2	PHE	-	LEADER SEQUENCE	UNP Q97G05
A	-1	GLN	-	LEADER SEQUENCE	UNP Q97G05
A	0	GLY	-	LEADER SEQUENCE	UNP Q97G05
A	1	MSE	MET	MODIFIED RESIDUE	UNP Q97G05
A	53	MSE	MET	MODIFIED RESIDUE	UNP Q97G05
A	64	MSE	MET	MODIFIED RESIDUE	UNP Q97G05
A	99	MSE	MET	MODIFIED RESIDUE	UNP Q97G05
B	-18	MSE	-	LEADER SEQUENCE	UNP Q97G05
B	-17	GLY	-	LEADER SEQUENCE	UNP Q97G05

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-16	SER	-	LEADER SEQUENCE	UNP Q97G05
B	-15	ASP	-	LEADER SEQUENCE	UNP Q97G05
B	-14	LYS	-	LEADER SEQUENCE	UNP Q97G05
B	-13	ILE	-	LEADER SEQUENCE	UNP Q97G05
B	-12	HIS	-	LEADER SEQUENCE	UNP Q97G05
B	-11	HIS	-	LEADER SEQUENCE	UNP Q97G05
B	-10	HIS	-	LEADER SEQUENCE	UNP Q97G05
B	-9	HIS	-	LEADER SEQUENCE	UNP Q97G05
B	-8	HIS	-	LEADER SEQUENCE	UNP Q97G05
B	-7	HIS	-	LEADER SEQUENCE	UNP Q97G05
B	-6	GLU	-	LEADER SEQUENCE	UNP Q97G05
B	-5	ASN	-	LEADER SEQUENCE	UNP Q97G05
B	-4	LEU	-	LEADER SEQUENCE	UNP Q97G05
B	-3	TYR	-	LEADER SEQUENCE	UNP Q97G05
B	-2	PHE	-	LEADER SEQUENCE	UNP Q97G05
B	-1	GLN	-	LEADER SEQUENCE	UNP Q97G05
B	0	GLY	-	LEADER SEQUENCE	UNP Q97G05
B	1	MSE	MET	MODIFIED RESIDUE	UNP Q97G05
B	53	MSE	MET	MODIFIED RESIDUE	UNP Q97G05
B	64	MSE	MET	MODIFIED RESIDUE	UNP Q97G05
B	99	MSE	MET	MODIFIED RESIDUE	UNP Q97G05

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0

- Molecule 3 is FLAVIN MONONUCLEOTIDE (three-letter code: FMN) (formula: C₁₇H₂₁N₄O₉P).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O P 31 17 4 9 1	0	0

- Molecule 4 is UNKNOWN LIGAND (three-letter code: UNL) (formula:).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total O 9 9	0	0

- Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	B	1	Total	C O	0	0
			4	2 2		


- Molecule 6 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	132	Total	O	0	2
			134	134		
6	B	143	Total	O	0	7
			150	150		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: NimC/NimA family protein

Chain A:  88% 7% 5%



- Molecule 1: NimC/NimA family protein

Chain B:  81% 10% 7%



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	52.43Å 65.31Å 108.79Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	26.86 – 1.80 26.86 – 1.80	Depositor EDS
% Data completeness (in resolution range)	95.9 (26.86-1.80) 95.9 (26.86-1.80)	Depositor EDS
R_{merge}	0.05	Depositor
R_{sym}	0.05	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.98 (at 1.80Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.168 , 0.198 0.181 , 0.207	Depositor DCC
R_{free} test set	1706 reflections (5.03%)	wwPDB-VP
Wilson B-factor (Å ²)	26.0	Xtrriage
Anisotropy	0.394	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 46.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.48$, $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2631	wwPDB-VP
Average B, all atoms (Å ²)	40.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.99% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FMN, UNL, SO4, EDO

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.56	0/1180	0.66	0/1581
1	B	0.66	1/1175 (0.1%)	0.72	0/1571
All	All	0.61	1/2355 (0.0%)	0.69	0/3152

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	1	MSE	SE-CE	-5.27	1.64	1.95

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1149	0	1115	11	0
1	B	1139	0	1130	17	0
2	A	5	0	0	0	0
2	B	10	0	0	0	0
3	A	31	0	19	1	0
4	A	9	0	0	4	0
5	B	4	0	6	1	0
6	A	134	0	0	1	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	150	0	0	0	0
All	All	2631	0	2270	28	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (28) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:340:UNL:O9	4:A:340:UNL:O4	1.64	1.15
4:A:340:UNL:O9	4:A:340:UNL:O8	1.74	1.03
1:A:19:ASN:ND2	1:B:19:ASN:OD1	1.94	0.99
1:B:5:LEU:HD12	1:B:32[B]:VAL:CG1	2.19	0.72
1:B:44:ASN:HB2	1:B:106[A]:ILE:HG22	1.77	0.65
1:B:32[A]:VAL:CG2	1:B:39:LEU:HD11	2.35	0.57
1:B:5:LEU:HD12	1:B:32[B]:VAL:HG12	1.89	0.54
1:A:35:TYR:CD2	1:A:88:LEU:HD22	2.42	0.54
1:A:87:GLU:HG2	1:A:102[B]:VAL:HG11	1.92	0.51
1:B:94:PRO:CB	1:B:97:LYS:HZ2	2.25	0.49
1:A:99:MSE:HE1	4:A:340:UNL:O2	2.13	0.48
1:B:94:PRO:CB	1:B:97:LYS:NZ	2.77	0.48
1:B:94:PRO:HB3	1:B:97:LYS:HZ2	1.80	0.47
1:A:102[A]:VAL:HG23	6:A:360:HOH:O	2.17	0.45
1:B:32[A]:VAL:HG21	1:B:39:LEU:HD11	1.98	0.45
1:A:99:MSE:HE1	4:A:340:UNL:O3	2.17	0.45
1:B:1:MSE:HG2	1:B:39:LEU:HB2	1.99	0.44
1:B:35:TYR:HD2	1:B:88:LEU:HD23	1.81	0.44
1:A:35:TYR:CD2	1:A:88:LEU:CD2	3.01	0.44
1:B:87:GLU:HG3	1:B:102:VAL:HG11	1.99	0.44
1:B:32[A]:VAL:HG12	5:B:134:EDO:H22	2.01	0.43
1:A:-9:HIS:O	1:A:-6:GLU:HB2	2.19	0.42
1:A:13[B]:VAL:HG22	3:A:339:FMN:HM72	2.00	0.42
1:A:19:ASN:HD21	1:B:19:ASN:CG	2.03	0.41
1:B:5:LEU:CD1	1:B:32[B]:VAL:CG1	2.96	0.41
1:B:64:MSE:HG2	1:B:70:TRP:HB3	2.03	0.41
1:A:99:MSE:HE2	1:A:100:TYR:CE2	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	144/150 (96%)	140 (97%)	4 (3%)	0	100	100
1	B	141/150 (94%)	136 (96%)	5 (4%)	0	100	100
All	All	285/300 (95%)	276 (97%)	9 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/125 (97%)	120 (99%)	1 (1%)	83	80
1	B	123/125 (98%)	121 (98%)	2 (2%)	65	57
All	All	244/250 (98%)	241 (99%)	3 (1%)	71	68

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	27	ARG
1	B	27	ARG
1	B	44	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	-11	HIS
1	A	19	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 1 is unknown - leaving 5 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	SO4	A	132	-	4,4,4	0.26	0	6,6,6	0.13	0
3	FMN	A	339	-	31,33,33	1.35	5 (16%)	39,50,50	2.04	6 (15%)
2	SO4	B	132	-	4,4,4	0.26	0	6,6,6	0.28	0
2	SO4	B	133	-	4,4,4	0.27	0	6,6,6	0.44	0
5	EDO	B	134	-	3,3,3	0.40	0	2,2,2	0.39	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	SO4	A	132	-	-	0/0/0/0	0/0/0/0
3	FMN	A	339	-	-	0/16/18/18	0/3/3/3
2	SO4	B	132	-	-	0/0/0/0	0/0/0/0
2	SO4	B	133	-	-	0/0/0/0	0/0/0/0
5	EDO	B	134	-	-	0/1/1/1	0/0/0/0

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	339	FMN	C1'-N10	2.07	1.50	1.48
3	A	339	FMN	C5A-N5	2.22	1.38	1.35
3	A	339	FMN	C4-N3	2.94	1.38	1.33
3	A	339	FMN	C10-N1	3.30	1.37	1.33
3	A	339	FMN	C4A-N5	4.06	1.39	1.33

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	339	FMN	C4A-C4-N3	-3.62	118.33	123.47
3	A	339	FMN	C9A-N10-C10	-2.21	118.83	121.77
3	A	339	FMN	C4A-N5-C5A	2.40	119.28	116.76
3	A	339	FMN	C5A-C9A-N10	4.39	121.06	117.71
3	A	339	FMN	C1'-N10-C9A	6.29	123.90	118.31
3	A	339	FMN	C4-N3-C2	7.42	121.46	115.14

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	339	FMN	1	0
5	B	134	EDO	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.