



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 12:08 pm GMT

PDB ID : 4IJ2
Title : Human methemoglobin in complex with the second and third NEAT domains of IsdH from *Staphylococcus aureus*
Authors : Dickson, C.F.; Jacques, D.A.; Guss, J.M.; Gell, D.A.
Deposited on : 2012-12-21
Resolution : 4.24 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 4.24 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.




Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	1017 (4.76-3.70)
Clashscore	122126	1086 (4.76-3.70)
Ramachandran outliers	120053	1039 (4.76-3.70)
Sidechain outliers	120020	1024 (4.76-3.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	141	92% 6% •
1	C	141	92% 6% •
2	B	146	94% 6%
2	D	146	94% 6%
3	E	336	83% 7% 11%
3	F	336	85% 7% 8%
3	G	336	84% 6% 11%

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	H	336	 56% . 41%

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 13547 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Hemoglobin subunit alpha.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	139	1044	670	181	190	3	0	0	0
1	C	139	1044	670	181	190	3	0	0	0

- Molecule 2 is a protein called Hemoglobin subunit beta.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	146	1123	724	195	201	3	0	0	0
2	D	146	1123	724	195	201	3	0	0	0

- Molecule 3 is a protein called Iron-regulated surface determinant protein H.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	E	300	2456	1577	401	474	4	0	0	0
3	F	309	2515	1612	412	487	4	0	0	0
3	G	300	2456	1577	401	474	4	0	0	0
3	H	197	1614	1035	264	313	2	0	0	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
E	325	SER	-	EXPRESSION TAG	UNP Q2FG07
E	642	ALA	TYR	ENGINEERED MUTATION	UNP Q2FG07
F	325	SER	-	EXPRESSION TAG	UNP Q2FG07
F	642	ALA	TYR	ENGINEERED MUTATION	UNP Q2FG07

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
G	325	SER	-	EXPRESSION TAG	UNP Q2FG07
G	642	ALA	TYR	ENGINEERED MUTATION	UNP Q2FG07
H	325	SER	-	EXPRESSION TAG	UNP Q2FG07
H	642	ALA	TYR	ENGINEERED MUTATION	UNP Q2FG07

- Molecule 4 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$).




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	B	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	C	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		
4	D	1	Total	C	Fe	N	O	0	0
			43	34	1	4	4		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Hemoglobin subunit alpha




- Molecule 1: Hemoglobin subunit alpha




- Molecule 2: Hemoglobin subunit beta



- Molecule 2: Hemoglobin subunit beta



- Molecule 3: Iron-regulated surface determinant protein H



- Molecule 3: Iron-regulated surface determinant protein H

4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	132.90Å 185.30Å 103.20Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	29.15 – 4.24 49.71 – 4.24	Depositor EDS
% Data completeness (in resolution range)	99.4 (29.15-4.24) 98.2 (49.71-4.24)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.30 (at 4.29Å)	Xtriage
Refinement program	BUSTER-TNT, BUSTER 2.10.0	Depositor
R, R_{free}	0.299 , 0.310 0.332 , 0.340	Depositor DCC
R_{free} test set	951 reflections (5.12%)	wwPDB-VP
Wilson B-factor (Å ²)	206.8	Xtriage
Anisotropy	0.106	Xtriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 107.1	EDS
L-test for twinning ²	$\langle L \rangle = 0.43$, $\langle L^2 \rangle = 0.25$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.84	EDS
Total number of atoms	13547	wwPDB-VP
Average B, all atoms (Å ²)	86.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.82% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.38	0/1071	0.50	0/1457
1	C	0.38	0/1071	0.50	0/1457
2	B	0.40	0/1153	0.50	0/1566
2	D	0.40	0/1153	0.50	0/1566
3	E	0.38	0/2509	0.51	0/3395
3	F	0.38	0/2569	0.51	0/3478
3	G	0.38	0/2509	0.51	0/3395
3	H	0.37	0/1649	0.50	0/2229
All	All	0.38	0/13684	0.50	0/18543

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1044	0	1047	5	0
1	C	1044	0	1047	5	0
2	B	1123	0	1118	5	0
2	D	1123	0	1118	6	0
3	E	2456	0	2425	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	F	2515	0	2482	8	0
3	G	2456	0	2425	5	0
3	H	1614	0	1595	3	0
4	A	43	0	30	0	0
4	B	43	0	30	1	0
4	C	43	0	30	0	0
4	D	43	0	30	1	0
All	All	13547	0	13377	40	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (40) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:97:HIS:HB3	1:C:41:THR:HG22	1.73	0.70
2:B:3:LEU:HD23	2:B:7:GLU:HB3	1.77	0.67
2:D:3:LEU:HD23	2:D:7:GLU:HB3	1.77	0.66
2:B:106:LEU:HD23	4:B:201:HEM:HBB2	1.82	0.61
3:F:570:GLU:HG2	3:F:589:LYS:HE3	1.86	0.58
3:E:570:GLU:HG2	3:E:589:LYS:HE3	1.86	0.58
3:G:570:GLU:HG2	3:G:589:LYS:HE3	1.86	0.57
3:F:604:VAL:HG12	3:F:620:PHE:HB3	1.92	0.52
3:E:604:VAL:HG12	3:E:620:PHE:HB3	1.92	0.51
3:G:604:VAL:HG12	3:G:620:PHE:HB3	1.92	0.51
3:H:432:THR:HB	3:H:462:ILE:HD12	1.94	0.49
3:E:432:THR:HB	3:E:462:ILE:HD12	1.95	0.49
3:G:432:THR:HB	3:G:462:ILE:HD12	1.95	0.49
2:D:106:LEU:HD23	4:D:201:HEM:HBB2	1.96	0.48
3:F:432:THR:HB	3:F:462:ILE:HD12	1.95	0.48
3:F:490:LEU:HA	3:F:493:GLN:HE21	1.79	0.47
1:A:43:PHE:HB3	1:A:46:PHE:HB2	1.95	0.47
1:A:41:THR:HB	2:D:97:HIS:HB2	1.96	0.47
3:H:490:LEU:HD13	3:H:526:GLN:HE22	1.80	0.46
1:C:113:LEU:HB3	1:C:116:GLU:HB2	1.98	0.46
1:A:113:LEU:HB3	1:A:116:GLU:HB2	1.98	0.46
1:C:37:PRO:HA	1:C:40:LYS:HG3	1.98	0.45
1:C:97:ASN:HA	1:C:100:LEU:HD12	1.98	0.45
3:E:483:PRO:HA	3:E:486:LYS:HG2	1.97	0.45
3:F:483:PRO:HA	3:F:486:LYS:HG2	1.99	0.45
3:F:487:ALA:HB1	3:F:492:ARG:HB3	1.98	0.44

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:41:THR:HB	2:D:97:HIS:CB	2.47	0.44
3:E:554:VAL:HG11	3:E:565:MET:HB3	2.00	0.43
3:F:554:VAL:HG11	3:F:565:MET:HB3	2.00	0.43
2:D:31:LEU:HD22	2:D:106:LEU:HD13	2.00	0.43
1:A:22:GLY:HA3	1:A:60:LYS:HD2	2.01	0.42
3:G:554:VAL:HG11	3:G:565:MET:HB3	2.00	0.42
1:C:22:GLY:HA3	1:C:60:LYS:HD2	2.02	0.42
3:E:396:TRP:HA	3:E:441:ILE:HG22	2.02	0.41
2:B:28:LEU:HD23	2:B:60:VAL:HG13	2.02	0.41
3:H:396:TRP:HA	3:H:441:ILE:HG22	2.02	0.41
3:G:396:TRP:HA	3:G:441:ILE:HG22	2.02	0.41
2:D:28:LEU:HD23	2:D:60:VAL:HG13	2.02	0.41
2:B:31:LEU:HD22	2:B:106:LEU:HD13	2.02	0.40
3:F:396:TRP:HA	3:F:441:ILE:HG22	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	137/141 (97%)	131 (96%)	5 (4%)	1 (1%)	24	66
1	C	137/141 (97%)	131 (96%)	6 (4%)	0	100	100
2	B	144/146 (99%)	139 (96%)	5 (4%)	0	100	100
2	D	144/146 (99%)	139 (96%)	5 (4%)	0	100	100
3	E	292/336 (87%)	281 (96%)	11 (4%)	0	100	100
3	F	303/336 (90%)	291 (96%)	12 (4%)	0	100	100
3	G	292/336 (87%)	281 (96%)	11 (4%)	0	100	100
3	H	193/336 (57%)	186 (96%)	7 (4%)	0	100	100
All	All	1642/1918 (86%)	1579 (96%)	62 (4%)	1 (0%)	53	87

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	44	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	110/113 (97%)	109 (99%)	1 (1%)	81	90
1	C	110/113 (97%)	110 (100%)	0	100	100
2	B	118/118 (100%)	116 (98%)	2 (2%)	63	83
2	D	118/118 (100%)	116 (98%)	2 (2%)	63	83
3	E	271/305 (89%)	261 (96%)	10 (4%)	37	66
3	F	276/305 (90%)	266 (96%)	10 (4%)	38	67
3	G	271/305 (89%)	262 (97%)	9 (3%)	41	68
3	H	175/305 (57%)	171 (98%)	4 (2%)	53	76
All	All	1449/1682 (86%)	1411 (97%)	38 (3%)	49	73

All (38) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	34	LEU
2	B	14	LEU
2	B	21	ASP
2	D	14	LEU
2	D	21	ASP
3	E	480	LEU
3	E	481	LEU
3	E	501	GLN
3	E	504	LEU
3	E	511	GLU
3	E	547	LEU
3	E	587	LYS
3	E	595	LYS

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
3	E	612	LYS
3	E	652	ASN
3	F	480	LEU
3	F	492	ARG
3	F	501	GLN
3	F	504	LEU
3	F	511	GLU
3	F	547	LEU
3	F	587	LYS
3	F	595	LYS
3	F	612	LYS
3	F	652	ASN
3	G	480	LEU
3	G	493	GLN
3	G	504	LEU
3	G	511	GLU
3	G	547	LEU
3	G	587	LYS
3	G	595	LYS
3	G	612	LYS
3	G	652	ASN
3	H	480	LEU
3	H	501	GLN
3	H	504	LEU
3	H	511	GLU

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (11) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	HIS
1	A	97	ASN
2	B	39	GLN
2	B	63	HIS
1	C	58	HIS
2	D	39	GLN
2	D	63	HIS
3	F	493	GLN
3	G	478	GLN
3	H	493	GLN
3	H	526	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

4 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	HEM	A	201	1	27,50,50	2.21	6 (22%)	17,82,82	1.23	1 (5%)
4	HEM	B	201	2	27,50,50	2.22	6 (22%)	17,82,82	1.23	1 (5%)
4	HEM	C	201	1	27,50,50	2.21	6 (22%)	17,82,82	1.26	2 (11%)
4	HEM	D	201	2	27,50,50	2.21	6 (22%)	17,82,82	1.35	2 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	HEM	A	201	1	-	0/6/54/54	0/0/8/8
4	HEM	B	201	2	-	0/6/54/54	0/0/8/8
4	HEM	C	201	1	-	0/6/54/54	0/0/8/8
4	HEM	D	201	2	-	0/6/54/54	0/0/8/8

All (24) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	A	201	HEM	C3C-C2C	-4.17	1.34	1.40
4	B	201	HEM	C3B-C2B	-4.12	1.34	1.40
4	B	201	HEM	C3C-C2C	-4.11	1.34	1.40
4	C	201	HEM	C3C-C2C	-4.11	1.34	1.40
4	D	201	HEM	C3C-C2C	-4.10	1.34	1.40
4	A	201	HEM	C3B-C2B	-3.93	1.34	1.40
4	D	201	HEM	C3B-C2B	-3.92	1.34	1.40
4	C	201	HEM	C3B-C2B	-3.88	1.35	1.40
4	B	201	HEM	CAA-C2A	2.27	1.55	1.52
4	C	201	HEM	CAA-C2A	2.27	1.55	1.52
4	D	201	HEM	CAA-C2A	2.27	1.55	1.52
4	A	201	HEM	CAA-C2A	2.30	1.55	1.52
4	C	201	HEM	C3C-CAC	3.81	1.55	1.47
4	A	201	HEM	C3C-CAC	3.86	1.55	1.47
4	B	201	HEM	C3C-CAC	3.89	1.55	1.47
4	D	201	HEM	C3C-CAC	3.93	1.55	1.47
4	D	201	HEM	C3B-CAB	3.97	1.55	1.47
4	A	201	HEM	C3B-CAB	4.05	1.55	1.47
4	C	201	HEM	C3B-CAB	4.10	1.56	1.47
4	B	201	HEM	C3B-CAB	4.11	1.56	1.47
4	B	201	HEM	C3D-C2D	5.55	1.54	1.37
4	A	201	HEM	C3D-C2D	5.57	1.54	1.37
4	D	201	HEM	C3D-C2D	5.60	1.54	1.37
4	C	201	HEM	C3D-C2D	5.65	1.54	1.37

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	B	201	HEM	C1D-C2D-C3D	-2.47	105.28	107.00
4	D	201	HEM	C1D-C2D-C3D	-2.42	105.31	107.00
4	D	201	HEM	CAD-CBD-CGD	-2.19	108.91	112.66
4	C	201	HEM	CBA-CAA-C2A	-2.05	108.56	112.48
4	C	201	HEM	C1D-C2D-C3D	-2.04	105.58	107.00
4	A	201	HEM	CMA-C3A-C4A	-2.03	125.34	128.46

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	B	201	HEM	1	0
4	D	201	HEM	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers [i](#)

Unable to reproduce the depositors R factor - this section is therefore empty.