



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 9, 2018 – 12:50 pm GMT

PDB ID : 1IKP  
Title : Pseudomonas Aeruginosa Exotoxin A, P201Q, W281A mutant  
Authors : McKay, D.B.; Wedekind, J.E.; Trame, C.B.  
Deposited on : 2001-05-04  
Resolution : 1.45 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.45 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1761 (1.48-1.44)
Clashscore	122126	1816 (1.48-1.44)
Ramachandran outliers	120053	1793 (1.48-1.44)
Sidechain outliers	120020	1793 (1.48-1.44)
RSRZ outliers	108989	1733 (1.48-1.44)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	613	

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 5185 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called EXOTOXIN A.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	599	4641	2898	837	892	14	0	9	0

There are 5 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	201	GLN	PRO	ENGINEERED	UNP P11439
A	281	ALA	TRP	ENGINEERED	UNP P11439
A	364	ASN	SER	SEE REMARK 999	UNP P11439
A	407	VAL	ILE	SEE REMARK 999	UNP P11439
A	515	SER	GLY	SEE REMARK 999	UNP P11439

- Molecule 2 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	2	Total	Cl	0	0
			2	2		

- Molecule 3 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	2	Total	Na	0	0
			2	2		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	540	Total	O	0	0
			540	540		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: EXOTOXIN A



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.91Å 99.07Å 58.63Å 90.00° 97.34° 90.00°	Depositor
Resolution (Å)	21.82 – 1.45 49.54 – 1.40	Depositor EDS
% Data completeness (in resolution range)	90.2 (21.82-1.45) 86.9 (49.54-1.40)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	0.03	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	4.11 (at 1.40Å)	Xtrriage
Refinement program	CNS 1.0	Depositor
R, $R_{free}$	0.205 , 0.228 0.209 , 0.233	Depositor DCC
$R_{free}$ test set	11675 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	16.0	Xtrriage
Anisotropy	0.316	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.015 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	5185	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.31% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	1/4740 (0.0%)	0.69	0/6447

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	84	PRO	N-CD	5.31	1.55	1.47

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4641	0	4506	70	0
2	A	2	0	0	0	0
3	A	2	0	0	0	0
4	A	540	0	0	10	0
All	All	5185	0	4506	70	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (70) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:37:THR:HG22	1:A:39:GLY:H	1.23	1.03
1:A:137:MET:CE	1:A:137:MET:HA	1.95	0.96
1:A:137:MET:HE2	1:A:141:LEU:HD23	1.57	0.85
1:A:137:MET:CE	1:A:141:LEU:HD23	2.13	0.79
1:A:137:MET:HE3	1:A:137:MET:HA	1.62	0.78
1:A:137:MET:HE2	1:A:137:MET:HA	1.66	0.74
1:A:348:GLU:HG2	1:A:352:ARG:HH11	1.54	0.72
1:A:262:HIS:HE1	1:A:391:GLU:OE2	1.74	0.71
1:A:348:GLU:HG2	1:A:352:ARG:NH1	2.06	0.70
1:A:212:GLN:HE22	1:A:432:ARG:HE	1.41	0.69
1:A:226:ARG:HD3	4:A:756:HOH:O	1.93	0.67
1:A:327:GLU:O	1:A:331:GLU:HG3	1.95	0.66
1:A:37:THR:HG21	1:A:176:MET:HE1	1.78	0.65
1:A:483:GLN:HE22	1:A:579:GLY:HA2	1.62	0.64
1:A:137:MET:CA	1:A:137:MET:HE2	2.28	0.63
1:A:488:ASP:OD1	1:A:492:ARG:HG2	1.98	0.63
1:A:279:ARG:HD2	1:A:279:ARG:N	2.14	0.63
1:A:353:GLN:HG3	4:A:786:HOH:O	1.98	0.63
1:A:308[B]:VAL:HG12	4:A:931:HOH:O	2.00	0.60
1:A:86:ARG:H	1:A:249:HIS:CD2	2.19	0.60
1:A:37:THR:HG21	1:A:176:MET:CE	2.32	0.58
1:A:18:ASP:OD1	1:A:20:LYS:HG2	2.03	0.58
1:A:37:THR:HG22	1:A:39:GLY:N	2.06	0.58
1:A:3:GLU:HG3	1:A:23:VAL:O	2.05	0.57
1:A:281:ALA:HB3	1:A:284:LEU:HB2	1.88	0.54
1:A:60:ILE:HD12	1:A:64:LEU:HD23	1.90	0.54
1:A:212:GLN:HE22	1:A:432:ARG:NE	2.06	0.53
1:A:137:MET:CA	1:A:137:MET:CE	2.72	0.53
1:A:289:TYR:O	1:A:293[B]:ARG:HG3	2.09	0.53
1:A:349:ARG:HH12	1:A:353:GLN:HG3	1.74	0.53
1:A:74[A]:ARG:HH11	1:A:74[A]:ARG:HB2	1.73	0.53
1:A:262:HIS:HD2	1:A:267:LEU:O	1.92	0.53
1:A:67:THR:HB	1:A:74[A]:ARG:HH11	1.74	0.52
1:A:276:ARG:HG2	4:A:1150:HOH:O	2.10	0.52
1:A:147:ARG:HD2	4:A:1005:HOH:O	2.08	0.52
1:A:420:GLU:HG3	4:A:1167:HOH:O	2.10	0.51
1:A:143:ALA:O	1:A:147:ARG:HG2	2.11	0.51
1:A:86:ARG:H	1:A:249:HIS:HD2	1.58	0.50
1:A:275:HIS:HD2	4:A:1169:HOH:O	1.94	0.49
1:A:137:MET:CE	1:A:141:LEU:CD2	2.89	0.49
1:A:262:HIS:CE1	1:A:391:GLU:OE2	2.61	0.49

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:524:ALA:O	1:A:528:GLU:HG3	2.12	0.49
1:A:118:HIS:HE1	1:A:202:LEU:HD21	1.76	0.49
1:A:485:GLN:HG3	1:A:486:GLU:HG2	1.95	0.48
1:A:212:GLN:NE2	1:A:432:ARG:NE	2.62	0.48
1:A:262:HIS:CD2	1:A:267:LEU:O	2.68	0.47
1:A:350:PHE:CE2	1:A:357:ASN:HB3	2.50	0.47
1:A:118:HIS:CE1	1:A:202:LEU:HD21	2.49	0.46
1:A:78:GLY:HA2	1:A:137:MET:HE3	1.99	0.45
1:A:306:ASN:ND2	1:A:307:GLN:HG3	2.31	0.45
1:A:348:GLU:O	1:A:352:ARG:HG3	2.17	0.45
1:A:254:GLY:H	1:A:364:ASN:HD22	1.65	0.44
1:A:522[A]:GLU:H	1:A:522[A]:GLU:CD	2.21	0.44
1:A:213:ARG:HH11	1:A:213:ARG:HG2	1.81	0.44
1:A:98:SER:HB3	1:A:118:HIS:HB2	1.99	0.43
1:A:136:GLU:O	1:A:137:MET:HE3	2.19	0.43
1:A:65[A]:SER:OG	4:A:867:HOH:O	2.20	0.43
1:A:74[A]:ARG:NH1	1:A:74[A]:ARG:HB2	2.34	0.43
1:A:80:GLU:OE2	1:A:275:HIS:ND1	2.50	0.42
1:A:240:LYS:HA	1:A:241:PRO:HD3	1.88	0.41
1:A:294:LEU:CD2	1:A:336:ALA:HA	2.50	0.41
1:A:603:GLN:HG2	4:A:1228:HOH:O	2.20	0.41
1:A:76:GLU:HA	1:A:85:VAL:O	2.21	0.41
1:A:349:ARG:NH1	1:A:353:GLN:HG3	2.34	0.41
1:A:284:LEU:HD22	1:A:289:TYR:CZ	2.56	0.41
1:A:208:TYR:OH	1:A:241:PRO:HG3	2.21	0.41
1:A:80:GLU:HG3	4:A:1072:HOH:O	2.20	0.41
1:A:175:VAL:HG21	1:A:223:LYS:HD2	2.03	0.40
1:A:462:LEU:HD12	1:A:463:ASP:O	2.21	0.40
1:A:498:LEU:HA	1:A:498:LEU:HD23	1.96	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	602/613 (98%)	586 (97%)	15 (2%)	1 (0%)	49 22

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	241	PRO

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	481/487 (99%)	474 (98%)	7 (2%)	67 35

All (7) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	80	GLU
1	A	124	ASN
1	A	137	MET
1	A	139	ASP
1	A	262	HIS
1	A	279	ARG
1	A	306	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	124	ASN
1	A	201	GLN
1	A	212	GLN
1	A	246	HIS
1	A	249	HIS
1	A	262	HIS
1	A	292	GLN
1	A	306	ASN

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	332	GLN
1	A	364	ASN
1	A	415[A]	GLN
1	A	483	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	599/613 (97%)	0.26	37 (6%) <span style="border: 1px solid red; padding: 2px;">20</span> <span style="border: 1px solid red; padding: 2px;">21</span>	9, 18, 48, 89	2 (0%)

All (37) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	279	ARG	9.3
1	A	549	GLY	6.9
1	A	489	ALA	6.8
1	A	33	ALA	5.7
1	A	492	ARG	5.7
1	A	355	THR	5.6
1	A	32	PRO	5.0
1	A	493	ILE	4.5
1	A	519	ALA	4.5
1	A	34	ILE	4.3
1	A	521	PRO	4.1
1	A	548	GLU	3.8
1	A	415[A]	GLN	3.6
1	A	490	ARG	3.5
1	A	487	PRO	3.5
1	A	38	ASN	3.2
1	A	518	LEU	3.1
1	A	322	GLY	2.9
1	A	280	GLY	2.8
1	A	35	ALA	2.8
1	A	302	ARG	2.8
1	A	491	GLY	2.8
1	A	323	GLY	2.7
1	A	37	THR	2.6
1	A	550	GLY	2.6
1	A	301	ALA	2.5
1	A	320	GLY	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	488	ASP	2.4
1	A	129	MET	2.2
1	A	516	LEU	2.1
1	A	36	ASP	2.1
1	A	354	GLY	2.1
1	A	83	LYS	2.1
1	A	2	GLU	2.0
1	A	356	GLY	2.0
1	A	522[A]	GLU	2.0
1	A	606	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	NA	A	704	1/1	0.94	0.18	30,30,30,30	0
2	CL	A	702	1/1	0.95	0.06	27,27,27,27	0
2	CL	A	701	1/1	0.98	0.04	17,17,17,17	0
3	NA	A	703	1/1	1.00	0.07	14,14,14,14	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.