



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 10, 2018 – 06:35 am GMT

PDB ID : 5IKV
Title : The Structure of Flufenamic Acid Bound to Human Cyclooxygenase-2
Authors : Orlando, B.J.; Malkowski, M.G.
Deposited on : 2016-03-03
Resolution : 2.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.7.3 (157068), CSD as539be (2018)
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	4155 (2.50-2.50)
Clashscore	122126	4827 (2.50-2.50)
Ramachandran outliers	120053	4735 (2.50-2.50)
Sidechain outliers	120020	4737 (2.50-2.50)
RSRZ outliers	108989	4058 (2.50-2.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	551	 3% 94% 6%
1	B	551	 3% 94% 5%

2 Entry composition [i](#)


There are 8 unique types of molecules in this entry. The entry contains 18566 atoms, of which 8946 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Prostaglandin G/H synthase 2.


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	551	8825	2883	4355	755	806	26	129	0	0
1	B	551	8825	2883	4355	755	806	26	37	0	0

- Molecule 2 is 2-[[3-(TRIFLUOROMETHYL)PHENYL]AMINO] BENZOIC ACID (three-letter code: FLF) (formula: C₁₄H₁₀F₃NO₂).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	F	H	N			O
2	A	1	29	14	3	9	1	2	0	0
2	B	1	29	14	3	9	1	2	0	0

- Molecule 3 is PROTOPORPHYRIN IX CONTAINING CO (three-letter code: COH) (formula: C₃₄H₃₂CoN₄O₄).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
			Total	C	Co	H	N			O
3	A	1	Total	C	Co	H	N	O	0	0
			73	34	1	30	4	4		
3	B	1	Total	C	Co	H	N	O	0	0
			73	34	1	30	4	4		

- Molecule 4 is N-ACETYL-D-GLUCOSAMINE (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	H	N	O		
4	A	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
4	A	1	Total	C	H	N	O	0	0
			26	8	12	1	5		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
4	A	1	Total	C	H	N	O	0	0
			27	8	13	1	5		
4	B	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
4	B	1	Total	C	H	N	O	0	0
			26	8	12	1	5		
4	B	1	Total	C	H	N	O	0	0
			27	8	13	1	5		

- Molecule 5 is ALPHA-D-MANNOSE (three-letter code: MAN) (formula: C₆H₁₂O₆).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	
5	A	1	Total	C	H	O	0	0
			21	6	10	5		
5	B	1	Total	C	H	O	0	0
			21	6	10	5		

- Molecule 6 is AMMONIUM ION (three-letter code: NH4) (formula: H₄N).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
6	A	1	Total	H	N	0	0
			5	4	1		
6	A	1	Total	H	N	0	0
			5	4	1		
6	A	1	Total	H	N	0	0
			5	4	1		
6	A	1	Total	H	N	0	0
			5	4	1		
6	A	1	Total	H	N	0	0
			5	4	1		
6	B	1	Total	H	N	0	0
			5	4	1		
6	B	1	Total	H	N	0	0
			5	4	1		
6	B	1	Total	H	N	0	0
			5	4	1		

- Molecule 7 is B-OCTYLGLUCOSIDE (three-letter code: BOG) (formula: C₁₄H₂₈O₆).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	H	O		
7	A	1	48	14	28	6	0	0


- Molecule 8 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
8	A	182	Total	O	0	0
			182	182		
8	B	237	Total	O	0	0
			237	237		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Prostaglandin G/H synthase 2



- Molecule 1: Prostaglandin G/H synthase 2



4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, α , β , γ	126.92Å 149.33Å 184.77Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	30.01 – 2.51 30.01 – 2.51	Depositor EDS
% Data completeness (in resolution range)	97.1 (30.01-2.51) 92.2 (30.01-2.51)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	3.09 (at 2.51Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, R_{free}	0.185 , 0.224 0.190 , 0.228	Depositor DCC
R_{free} test set	2953 reflections (5.04%)	wwPDB-VP
Wilson B-factor (Å ²)	34.2	Xtrriage
Anisotropy	0.068	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.41 , 48.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	18566	wwPDB-VP
Average B, all atoms (Å ²)	49.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 2.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN, NH4, COH, FLF, BOG

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.31	0/4602	0.42	0/6241
1	B	0.34	0/4602	0.43	0/6241
All	All	0.33	0/9204	0.43	0/12482

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4470	4355	4354	17	0
1	B	4470	4355	4354	14	0
2	A	20	9	9	0	0
2	B	20	9	9	3	0
3	A	43	30	30	4	0
3	B	43	30	30	2	0
4	A	42	37	37	0	0
4	B	42	37	36	0	0
5	A	11	10	10	0	0
5	B	11	10	10	0	0
6	A	5	20	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	B	4	16	0	0	0
7	A	20	28	28	2	0
8	A	182	0	0	0	0
8	B	237	0	0	1	0
All	All	9620	8946	8907	37	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (37) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:484:GLU:OE1	1:B:487:MET:N	1.95	0.99
1:B:482:THR:OG1	1:B:484:GLU:OE2	2.00	0.75
3:B:602:COH:HHD	3:B:602:COH:HBC1	1.70	0.72
3:A:602:COH:HBB1	3:A:602:COH:HHC	1.75	0.67
1:B:157:ASP:N	1:B:157:ASP:OD1	2.27	0.66
1:A:101:ASN:O	1:A:105:ASN:ND2	2.30	0.65
3:B:602:COH:HHC	3:B:602:COH:HBB1	1.78	0.65
1:A:567:LEU:HA	7:A:612:BOG:H8'1	1.83	0.60
1:B:514:PRO:O	8:B:701:HOH:O	2.18	0.56
1:A:35:PRO:HB2	1:A:55:TYR:HB3	1.88	0.56
3:A:602:COH:HMC1	3:A:602:COH:HBC1	1.88	0.54
2:B:601:FLF:C6'	2:B:601:FLF:H5	2.42	0.49
1:B:385:TYR:OH	2:B:601:FLF:O2	2.17	0.48
1:A:484:GLU:OE1	1:A:487:MET:N	2.38	0.46
1:A:479:GLU:O	1:A:483:GLY:N	2.42	0.45
1:A:178:VAL:HG11	1:A:445:GLN:HG3	1.99	0.45
1:A:388:HIS:N	1:A:389:PRO:CD	2.79	0.45
1:B:388:HIS:N	1:B:389:PRO:CD	2.80	0.45
1:B:352:LEU:HD11	1:B:518:PHE:CE2	2.52	0.45
1:B:403:ASN:ND2	1:B:406:GLN:HG2	2.32	0.45
1:A:463:PHE:CD1	1:A:506:ALA:HB3	2.52	0.44
1:B:504:TYR:HB3	1:B:505:PRO:HD3	1.99	0.44
1:A:212:THR:HG22	3:A:602:COH:O2D	2.18	0.44
1:A:83:LYS:HE3	1:A:471:MET:CE	2.47	0.44
1:B:463:PHE:CD1	1:B:506:ALA:HB3	2.53	0.44
1:A:91:TYR:CE1	1:A:95:HIS:CE1	3.06	0.44
1:B:414:LEU:HD11	1:B:419:ILE:HD13	2.00	0.43
1:B:181:LEU:HD23	1:B:487:MET:HG2	2.00	0.43
1:A:388:HIS:CE1	1:A:447:VAL:HG11	2.54	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:389:PRO:HB2	1:B:434:VAL:HA	2.01	0.42
1:A:83:LYS:HE3	1:A:471:MET:HE2	2.02	0.42
1:B:269:THR:O	1:B:270:GLN:HB2	2.19	0.42
2:B:601:FLF:H6'	2:B:601:FLF:H5	2.01	0.42
1:A:34:ASN:HA	1:A:35:PRO:HD3	1.83	0.42
1:A:504:TYR:HB3	1:A:505:PRO:HD3	2.03	0.40
1:A:212:THR:HG22	3:A:602:COH:CGD	2.51	0.40
1:A:566:SER:HB2	7:A:612:BOG:H3'1	2.04	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	549/551 (100%)	535 (97%)	14 (3%)	0	100	100
1	B	549/551 (100%)	534 (97%)	15 (3%)	0	100	100
All	All	1098/1102 (100%)	1069 (97%)	29 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	490/490 (100%)	480 (98%)	10 (2%)	58	82

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	B	490/490 (100%)	478 (98%)	12 (2%)	52	78
All	All	980/980 (100%)	958 (98%)	22 (2%)	55	80

All (22) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	50	VAL
1	A	75	LEU
1	A	173	ASP
1	A	212	THR
1	A	232	HIS
1	A	278	GLN
1	A	289	GLN
1	A	385	TYR
1	A	484	GLU
1	A	513	ARG
1	B	75	LEU
1	B	157	ASP
1	B	170	GLN
1	B	212	THR
1	B	232	HIS
1	B	278	GLN
1	B	289	GLN
1	B	385	TYR
1	B	398	HIS
1	B	484	GLU
1	B	513	ARG
1	B	573	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 22 ligands modelled in this entry, 9 are modelled with single atom - leaving 13 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	FLF	A	601	-	18,21,21	0.66	0	26,30,30	2.19	3 (11%)
3	COH	A	602	1	32,50,50	5.26	23 (71%)	23,82,82	4.53	9 (39%)
4	NAG	A	603	1,4	14,14,15	0.29	0	17,19,21	0.50	0
4	NAG	A	604	5,4	14,14,15	0.28	0	17,19,21	0.56	0
5	MAN	A	605	4	11,11,12	1.00	1 (9%)	15,15,17	1.79	2 (13%)
4	NAG	A	606	1	14,14,15	0.17	0	17,19,21	0.55	0
7	BOG	A	612	-	20,20,20	1.11	1 (5%)	25,25,25	0.84	1 (4%)
2	FLF	B	601	-	18,21,21	1.79	3 (16%)	26,30,30	1.72	2 (7%)
3	COH	B	602	1	32,50,50	5.23	23 (71%)	23,82,82	4.79	12 (52%)
4	NAG	B	603	1,4	14,14,15	0.30	0	17,19,21	0.58	0
4	NAG	B	604	5,4	14,14,15	0.31	0	17,19,21	0.73	0
5	MAN	B	605	4	11,11,12	1.01	2 (18%)	15,15,17	1.58	2 (13%)
4	NAG	B	606	1	14,14,15	0.32	0	17,19,21	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	FLF	A	601	-	-	0/10/14/14	0/2/2/2
3	COH	A	602	1	-	0/8/94/94	0/0/8/8
4	NAG	A	603	1,4	-	0/6/23/26	0/1/1/1
4	NAG	A	604	5,4	-	0/6/23/26	0/1/1/1

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	MAN	A	605	4	-	0/2/19/22	1/1/1/1
4	NAG	A	606	1	-	0/6/23/26	0/1/1/1
7	BOG	A	612	-	-	0/11/31/31	0/1/1/1
2	FLF	B	601	-	-	0/10/14/14	0/2/2/2
3	COH	B	602	1	-	0/8/94/94	0/0/8/8
4	NAG	B	603	1,4	-	0/6/23/26	0/1/1/1
4	NAG	B	604	5,4	-	0/6/23/26	0/1/1/1
5	MAN	B	605	4	-	0/2/19/22	1/1/1/1
4	NAG	B	606	1	-	0/6/23/26	0/1/1/1

All (53) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	A	602	COH	CHD-C4C	-8.40	1.37	1.53
3	A	602	COH	CHA-C1A	-8.25	1.37	1.53
3	B	602	COH	CHA-C1A	-8.14	1.37	1.53
3	B	602	COH	CHD-C4C	-8.11	1.37	1.53
3	B	602	COH	CHB-C4A	-8.07	1.37	1.53
3	A	602	COH	CHB-C4A	-7.82	1.38	1.53
3	A	602	COH	CHC-C1C	-7.74	1.38	1.53
3	B	602	COH	CHC-C1C	-7.59	1.38	1.53
3	A	602	COH	CHD-C1D	-5.57	1.37	1.51
3	B	602	COH	CHD-C1D	-5.54	1.37	1.51
3	A	602	COH	CHA-C4D	-5.29	1.38	1.51
3	B	602	COH	CHA-C4D	-5.29	1.38	1.51
3	B	602	COH	CHB-C1B	-5.20	1.38	1.51
3	A	602	COH	CHB-C1B	-4.96	1.38	1.51
3	B	602	COH	CHC-C4B	-4.90	1.38	1.51
3	A	602	COH	CHC-C4B	-4.80	1.39	1.51
3	B	602	COH	C3B-C2B	-4.37	1.34	1.40
3	A	602	COH	C3B-C2B	-4.04	1.34	1.40
3	A	602	COH	C4C-C3C	-4.04	1.44	1.51
2	B	601	FLF	C1-C6	-3.72	1.35	1.40
3	B	602	COH	C4C-C3C	-3.53	1.45	1.51
2	B	601	FLF	C5-C6	-2.47	1.35	1.39
2	B	601	FLF	C6'-C1'	-2.32	1.35	1.39
3	A	602	COH	CO-NB	2.00	2.07	1.97
5	B	605	MAN	C2-C3	2.03	1.55	1.52
5	B	605	MAN	C1-C2	2.16	1.57	1.52
5	A	605	MAN	O5-C5	2.27	1.48	1.43
3	A	602	COH	CMA-C3A	2.53	1.54	1.50
3	B	602	COH	CAA-C2A	2.54	1.54	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	602	COH	CMA-C3A	2.56	1.54	1.50
3	B	602	COH	CO-NB	2.64	2.11	1.97
3	B	602	COH	CMC-C2C	2.68	1.55	1.50
3	A	602	COH	CAA-C2A	2.73	1.55	1.51
7	A	612	BOG	O5-C1	2.77	1.49	1.41
3	A	602	COH	CMC-C2C	2.85	1.55	1.50
3	A	602	COH	C3B-CAB	4.02	1.55	1.47
3	B	602	COH	C3B-CAB	4.04	1.55	1.47
3	A	602	COH	C4D-C3D	4.08	1.43	1.38
3	B	602	COH	C4D-C3D	4.36	1.44	1.38
3	A	602	COH	C1D-C2D	4.46	1.44	1.38
3	B	602	COH	C1D-C2D	4.76	1.44	1.38
3	B	602	COH	C3D-C2D	4.98	1.52	1.37
3	B	602	COH	C1B-C2B	5.06	1.45	1.38
3	A	602	COH	C3D-C2D	5.13	1.52	1.37
3	A	602	COH	C1B-C2B	5.20	1.45	1.38
3	B	602	COH	C1B-NB	7.98	1.38	1.34
3	A	602	COH	C1B-NB	8.41	1.38	1.34
3	B	602	COH	C4D-ND	8.83	1.39	1.34
3	B	602	COH	C4B-NB	9.02	1.39	1.34
3	A	602	COH	C4D-ND	9.14	1.39	1.34
3	A	602	COH	C4B-NB	9.26	1.39	1.34
3	A	602	COH	C1D-ND	10.24	1.39	1.34
3	B	602	COH	C1D-ND	10.57	1.40	1.34

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	A	601	FLF	C2-C1-C7	-9.22	106.05	120.23
2	B	601	FLF	C2-C1-C7	-7.13	109.26	120.23
3	B	602	COH	CBD-CAD-C3D	-4.27	104.32	112.48
3	A	602	COH	CBD-CAD-C3D	-4.12	104.62	112.48
3	B	602	COH	CHD-C1D-C2D	-3.38	123.59	129.45
2	A	601	FLF	C1-C6-N	-3.34	115.79	119.64
3	B	602	COH	CAD-C3D-C4D	-3.30	124.98	127.30
3	B	602	COH	CBA-CAA-C2A	-3.26	108.46	114.28
2	B	601	FLF	C1-C6-N	-3.03	116.15	119.64
3	A	602	COH	CAD-CBD-CGD	-2.84	107.81	112.66
3	B	602	COH	CAD-CBD-CGD	-2.69	108.06	112.66
3	A	602	COH	CHD-C1D-C2D	-2.69	124.80	129.45
5	B	605	MAN	O2-C2-C3	-2.65	105.03	110.19
7	A	612	BOG	C1-O5-C5	-2.12	109.53	113.71

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	602	COH	CHB-C1B-C2B	-2.12	125.79	129.45
5	A	605	MAN	O2-C2-C3	-2.03	106.23	110.19
3	A	602	COH	CMA-C3A-C4A	2.07	126.30	120.24
3	B	602	COH	CMA-C3A-C4A	2.10	126.38	120.24
3	B	602	COH	C4B-C3B-C2B	2.30	107.57	104.13
3	A	602	COH	C4B-C3B-C2B	2.35	107.64	104.13
2	A	601	FLF	C2-C1-C6	3.16	121.06	117.97
5	B	605	MAN	C1-O5-C5	4.77	118.75	112.19
5	A	605	MAN	C1-O5-C5	6.23	120.76	112.19
3	A	602	COH	CHB-C4A-NA	9.50	124.10	110.12
3	B	602	COH	CHD-C4C-NC	9.79	124.52	110.12
3	B	602	COH	CHB-C4A-NA	10.07	124.94	110.12
3	A	602	COH	CHC-C1C-NC	10.20	125.13	110.12
3	A	602	COH	CHD-C4C-NC	10.28	125.25	110.12
3	A	602	COH	CHA-C1A-NA	10.37	125.38	110.12
3	B	602	COH	CHA-C1A-NA	10.52	125.60	110.12
3	B	602	COH	CHC-C1C-NC	11.28	126.72	110.12

There are no chirality outliers.

There are no torsion outliers.

All (2) ring outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	A	605	MAN	C1-C2-C3-C4-C5-O5
5	B	605	MAN	C1-C2-C3-C4-C5-O5

4 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	602	COH	4	0
7	A	612	BOG	2	0
2	B	601	FLF	3	0
3	B	602	COH	2	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	551/551 (100%)	0.02	17 (3%) 49 52	26, 44, 68, 80	14 (2%)
1	B	551/551 (100%)	-0.02	15 (2%) 54 58	26, 41, 61, 83	4 (0%)
All	All	1102/1102 (100%)	-0.00	32 (2%) 51 55	26, 42, 65, 83	18 (1%)

All (32) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	74	PHE	3.5
1	A	277	PRO	3.2
1	A	160	PRO	3.2
1	B	107	PHE	3.0
1	B	75	LEU	2.9
1	A	165	VAL	2.9
1	A	278	GLN	2.7
1	B	78	ILE	2.6
1	B	277	PRO	2.6
1	A	272	GLU	2.5
1	A	270	GLN	2.5
1	B	278	GLN	2.4
1	A	368	ASN	2.3
1	B	80	LEU	2.3
1	A	216	ARG	2.3
1	A	506	ALA	2.2
1	A	72	PRO	2.2
1	A	82	LEU	2.2
1	B	65	TYR	2.1
1	A	35	PRO	2.1
1	A	493	ALA	2.1
1	B	446	LYS	2.1
1	B	368	ASN	2.1
1	B	299	MET	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	158	ASP	2.1
1	B	54	GLN	2.1
1	A	507	LEU	2.0
1	B	158	ASP	2.0
1	B	270	GLN	2.0
1	B	52	PHE	2.0
1	A	496	GLY	2.0
1	A	80	LEU	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
5	MAN	B	605	11/12	0.73	0.28	70,79,95,95	0
7	BOG	A	612	20/20	0.74	0.38	28,96,132,143	0
5	MAN	A	605	11/12	0.77	0.32	58,127,164,179	0
4	NAG	B	606	14/15	0.79	0.20	54,76,94,95	0
6	NH4	A	608	1/1	0.83	0.27	36,44,44,44	0
6	NH4	B	609	1/1	0.87	0.22	36,43,43,43	0
6	NH4	A	610	1/1	0.89	0.09	32,39,39,39	0
6	NH4	B	608	1/1	0.90	0.12	45,54,54,54	0
3	COH	A	602	43/43	0.92	0.21	50,68,102,114	0
4	NAG	B	604	14/15	0.93	0.22	51,63,80,82	0
6	NH4	B	610	1/1	0.93	0.46	13,16,16,16	0
3	COH	B	602	43/43	0.94	0.19	48,73,106,116	0
6	NH4	A	607	1/1	0.94	0.19	23,28,28,28	0
4	NAG	A	606	14/15	0.94	0.14	47,55,66,67	0
6	NH4	B	607	1/1	0.94	0.27	26,31,31,31	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	NAG	A	604	14/15	0.94	0.30	42,54,63,70	0
4	NAG	A	603	14/15	0.95	0.17	31,40,55,55	0
2	FLF	A	601	20/20	0.96	0.15	31,47,58,60	0
6	NH4	A	609	1/1	0.97	0.18	33,40,40,40	0
2	FLF	B	601	20/20	0.97	0.16	28,33,40,44	0
6	NH4	A	611	1/1	0.97	0.31	13,16,16,16	0
4	NAG	B	603	14/15	0.97	0.12	30,41,47,52	0

6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.