



Full wwPDB X-ray Structure Validation Report ⓘ

May 25, 2020 – 07:07 am BST

PDB ID : 2ILK
Title : CRYSTAL STRUCTURE OF HUMAN INTERLEUKIN-10 AT 1.6
ANGSTROMS RESOLUTION
Authors : Zdanov, A.; Schalk-Hihi, C.; Wlodawer, A.
Deposited on : 1996-07-01
Resolution : 1.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	3665 (1.60-1.60)
Ramachandran outliers	138981	3564 (1.60-1.60)
Sidechain outliers	138945	3563 (1.60-1.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	160	 64% 21% 9% . .

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 1598 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called INTERLEUKIN-10.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	155	1280	809	225	235	11	0	4	0

- Molecule 2 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	S		
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0
2	A	1	5	4	1	0	0

- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	303	Total 303	O 303	0	0

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: INTERLEUKIN-10

Chain A:  64% 21% 9%



4 Data and refinement statistics

Property	Value	Source
Space group	P 32 2 1	Depositor
Cell constants a, b, c, α , β , γ	69.53Å 69.53Å 70.54Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	10.00 – 1.60 9.95 – 1.60	Depositor EDS
% Data completeness (in resolution range)	(Not available) (10.00-1.60) 81.6 (9.95-1.60)	Depositor EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$	-	Xtrriage
Refinement program	PROFFT	Depositor
R, R_{free}	(Not available) , (Not available) 0.160 , (Not available)	Depositor DCC
R_{free} test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å ²)	16.0	Xtrriage
Anisotropy	0.016	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.16 , 98.1	EDS
L-test for twinning ¹	$\langle L \rangle = 0.44$, $\langle L^2 \rangle = 0.26$	Xtrriage
Estimated twinning fraction	0.048 for -h,-k,l	Xtrriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	1598	wwPDB-VP
Average B, all atoms (Å ²)	27.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.59% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.78	0/1320	1.98	36/1765 (2.0%)

There are no bond length outliers.

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	102	ARG	NE-CZ-NH1	15.33	127.97	120.30
1	A	102	ARG	NE-CZ-NH2	-10.82	114.89	120.30
1	A	59	TYR	CB-CG-CD2	-10.55	114.67	121.00
1	A	104	ARG	NE-CZ-NH2	-10.44	115.08	120.30
1	A	106[A]	ARG	NE-CZ-NH2	9.60	125.10	120.30
1	A	106[B]	ARG	NE-CZ-NH2	9.60	125.10	120.30
1	A	159	ARG	CD-NE-CZ	8.93	136.10	123.60
1	A	104	ARG	NE-CZ-NH1	8.67	124.64	120.30
1	A	55	ASP	CB-CG-OD1	8.51	125.96	118.30
1	A	59	TYR	CB-CG-CD1	8.16	125.89	121.00
1	A	10	ASN	N-CA-C	7.89	132.29	111.00
1	A	137	TYR	CB-CG-CD2	-7.83	116.30	121.00
1	A	159	ARG	NE-CZ-NH2	7.76	124.18	120.30
1	A	7	GLN	C-N-CA	7.68	140.89	121.70
1	A	84	ASP	CB-CG-OD1	7.60	125.14	118.30
1	A	72	TYR	CB-CG-CD2	-7.35	116.59	121.00
1	A	7	GLN	CA-C-O	7.30	135.43	120.10
1	A	91	VAL	CG1-CB-CG2	-7.21	99.36	110.90
1	A	156	MET	CA-CB-CG	-7.18	101.09	113.30
1	A	27	ARG	NE-CZ-NH2	-6.92	116.84	120.30
1	A	12	CYS	N-CA-CB	-6.65	98.64	110.60
1	A	106[A]	ARG	NE-CZ-NH1	-6.54	117.03	120.30
1	A	106[B]	ARG	NE-CZ-NH1	-6.54	117.03	120.30
1	A	50	GLU	CB-CA-C	6.27	122.93	110.40

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	9	GLU	C-N-CA	6.09	136.92	121.70
1	A	42	GLN	CB-CG-CD	6.00	127.20	111.60
1	A	36	PHE	CB-CG-CD2	5.92	124.94	120.80
1	A	142	GLU	OE1-CD-OE2	-5.75	116.40	123.30
1	A	54	GLU	OE1-CD-OE2	-5.74	116.42	123.30
1	A	10	ASN	CA-C-O	5.71	132.08	120.10
1	A	32	ARG	CD-NE-CZ	5.69	131.56	123.60
1	A	10	ASN	N-CA-CB	-5.63	100.46	110.60
1	A	87	ILE	N-CA-CB	5.41	123.23	110.80
1	A	72	TYR	CB-CG-CD1	5.34	124.21	121.00
1	A	38	GLN	CB-CA-C	-5.34	99.72	110.40
1	A	77	MET	CA-CB-CG	-5.18	104.48	113.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1280	0	1271	46	2
2	A	15	0	0	2	0
3	A	303	0	0	26	7
All	All	1598	0	1271	46	9

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 18.

All (46) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:126:ASN:HB3	3:A:452:HOH:O	1.29	1.28
1:A:75:GLU:HG2	3:A:423:HOH:O	1.42	1.16
1:A:10:ASN:CB	1:A:108:CYS:HA	1.85	1.07
1:A:28:ASP:HB3	3:A:501:HOH:O	1.62	0.99
1:A:9:GLU:HA	1:A:109:HIS:HA	1.57	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:79:GLN:NE2	3:A:454:HOH:O	2.11	0.84
1:A:117:LYS:HG2	2:A:601:SO4:O1	1.79	0.82
1:A:54:GLU:OE1	3:A:247:HOH:O	2.03	0.76
1:A:121:VAL:HG13	3:A:496:HOH:O	1.86	0.76
1:A:54:GLU:HG2	3:A:281:HOH:O	1.85	0.74
1:A:63:GLN:OE1	3:A:399:HOH:O	2.06	0.73
1:A:158:ILE:CD1	3:A:500:HOH:O	2.39	0.70
1:A:81:GLU:OE2	1:A:88[A]:LYS:HE2	1.93	0.69
1:A:158:ILE:HD13	3:A:500:HOH:O	1.95	0.66
1:A:8:SER:CA	1:A:110:ARG:HE	2.11	0.63
1:A:42:GLN:NE2	3:A:307:HOH:O	2.32	0.62
1:A:7:GLN:HG3	3:A:261:HOH:O	1.99	0.61
1:A:160:ASN:ND2	2:A:603:SO4:O1	2.35	0.57
1:A:75:GLU:OE1	3:A:459:HOH:O	2.17	0.56
1:A:18:ASN:HD21	1:A:104:ARG:HE	1.53	0.56
1:A:39:MET:HB3	3:A:353:HOH:O	2.09	0.53
1:A:10:ASN:CB	1:A:12:CYS:SG	2.97	0.52
1:A:28:ASP:OD1	3:A:419:HOH:O	2.19	0.51
1:A:45:ASN:ND2	1:A:45:ASN:H	2.10	0.50
1:A:74:GLU:HA	3:A:469:HOH:O	2.10	0.50
1:A:159:ARG:HB2	3:A:499:HOH:O	2.11	0.50
1:A:157:LYS:CE	3:A:245:HOH:O	2.60	0.49
1:A:159:ARG:CB	3:A:499:HOH:O	2.59	0.48
1:A:87:ILE:C	1:A:87:ILE:HD12	2.34	0.48
1:A:40:LYS:NZ	3:A:373:HOH:O	2.47	0.48
1:A:73:LEU:O	3:A:469:HOH:O	2.20	0.47
1:A:6:THR:O	1:A:8:SER:N	2.40	0.47
1:A:18:ASN:ND2	1:A:104:ARG:HH21	2.13	0.47
1:A:126:ASN:CG	3:A:470:HOH:O	2.54	0.47
1:A:84:ASP:HB3	1:A:87:ILE:HG13	1.97	0.47
1:A:10:ASN:CB	1:A:109:HIS:H	2.29	0.46
1:A:10:ASN:H	1:A:109:HIS:H	1.65	0.45
1:A:7:GLN:C	1:A:110:ARG:HH21	2.21	0.44
1:A:9:GLU:CB	3:A:210:HOH:O	2.66	0.44
1:A:7:GLN:HG2	3:A:290:HOH:O	2.18	0.44
1:A:157:LYS:HE3	3:A:245:HOH:O	2.18	0.43
1:A:103:LEU:HD21	1:A:107:ARG:CZ	2.47	0.43
1:A:117:LYS:HE2	1:A:117:LYS:HB3	1.39	0.42
1:A:79:GLN:HE21	1:A:79:GLN:HB2	1.58	0.42
1:A:39:MET:CB	3:A:353:HOH:O	2.69	0.41
1:A:134:LYS:HB2	1:A:134:LYS:HE3	1.82	0.40

All (9) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
3:A:235:HOH:O	3:A:235:HOH:O[4_556]	1.92	0.28
3:A:245:HOH:O	3:A:414:HOH:O[4_556]	2.05	0.15
3:A:253:HOH:O	3:A:406:HOH:O[5_676]	2.06	0.14
3:A:345:HOH:O	3:A:351:HOH:O[3_565]	2.08	0.12
3:A:276:HOH:O	3:A:438:HOH:O[6_655]	2.09	0.11
3:A:229:HOH:O	3:A:386:HOH:O[4_556]	2.11	0.09
1:A:6:THR:O	1:A:59:TYR:OH[4_556]	2.15	0.05
1:A:7:GLN:NE2	1:A:38:GLN:O[3_565]	2.17	0.03
3:A:428:HOH:O	3:A:428:HOH:O[6_766]	2.19	0.01

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	157/160 (98%)	151 (96%)	2 (1%)	4 (2%)	5 0

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	8	SER
1	A	10	ASN
1	A	9	GLU
1	A	11	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	142/145 (98%)	134 (94%)	8 (6%)	21 5

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	7	GLN
1	A	11	SER
1	A	18	ASN
1	A	42	GLN
1	A	45	ASN
1	A	117	LYS
1	A	156	MET
1	A	157	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	18	ASN
1	A	38	GLN
1	A	45	ASN
1	A	63	GLN
1	A	70	GLN
1	A	79	GLN
1	A	83	GLN
1	A	92	ASN
1	A	97	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

3 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	SO4	A	602	-	4,4,4	0.68	0	6,6,6	0.60	0
2	SO4	A	601	-	4,4,4	0.67	0	6,6,6	0.50	0
2	SO4	A	603	-	4,4,4	0.62	0	6,6,6	0.31	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

2 monomers are involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	601	SO4	1	0
2	A	603	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.