



Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:08 am BST

PDB ID : 4ILT
Title : Structure of the dioxygenase domain of SACTE_2871, a novel dioxygenase carbohydrate-binding protein fusion from the cellulolytic bacterium *Streptomyces* sp. SirexAA-E
Authors : Bianchetti, C.M.; Takasuka, T.E.; Bergeman, L.F.; Harmann, C.H.; Fox, B.G.
Deposited on : 2012-12-31
Resolution : 2.55 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.55 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1284 (2.56-2.52)
Clashscore	141614	1332 (2.56-2.52)
Ramachandran outliers	138981	1315 (2.56-2.52)
Sidechain outliers	138945	1315 (2.56-2.52)
RSRZ outliers	127900	1272 (2.56-2.52)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	155	
1	B	155	
1	C	155	
1	D	155	

2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 4899 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Intradiol ring-cleavage dioxygenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	154	1172	749	201	218	4	0	0	0
1	B	154	1172	749	201	218	4	0	0	0
1	C	155	1178	752	202	220	4	0	0	0
1	D	154	1173	749	201	219	4	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	76	SER	-	EXPRESSION TAG	UNP G2NEL6
B	76	SER	-	EXPRESSION TAG	UNP G2NEL6
C	76	SER	-	EXPRESSION TAG	UNP G2NEL6
D	76	SER	-	EXPRESSION TAG	UNP G2NEL6

- Molecule 2 is FE (III) ION (three-letter code: FE) (formula: Fe).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	B	1	Total 1	Fe 1	0	0
2	A	1	Total 1	Fe 1	0	0
2	D	1	Total 1	Fe 1	0	0
2	C	1	Total 1	Fe 1	0	0

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total Cl 1 1	0	0
3	D	1	Total Cl 1 1	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	51	Total O 51 51	0	0
4	B	72	Total O 72 72	0	0
4	C	45	Total O 45 45	0	0
4	D	30	Total O 30 30	0	0

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Intradiol ring-cleavage dioxygenase

Chain A: 




- Molecule 1: Intradiol ring-cleavage dioxygenase

Chain B: 




- Molecule 1: Intradiol ring-cleavage dioxygenase

Chain C: 



- Molecule 1: Intradiol ring-cleavage dioxygenase

Chain D: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	49.19Å 85.12Å 70.26Å 90.00° 95.53° 90.00°	Depositor
Resolution (Å)	42.56 – 2.55 42.56 – 2.55	Depositor EDS
% Data completeness (in resolution range)	98.5 (42.56-2.55) 92.4 (42.56-2.55)	Depositor EDS
R_{merge}	0.15	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.36 (at 2.54Å)	Xtrriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.203 , 0.268 0.205 , 0.270	Depositor DCC
R_{free} test set	931 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å ²)	30.8	Xtrriage
Anisotropy	0.135	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.30 , 34.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.92	EDS
Total number of atoms	4899	wwPDB-VP
Average B, all atoms (Å ²)	36.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.12% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: FE, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.50	0/1209	0.66	0/1653
1	B	0.51	0/1209	0.68	0/1653
1	C	0.49	0/1215	0.66	0/1661
1	D	0.44	0/1210	0.63	0/1654
All	All	0.49	0/4843	0.66	0/6621

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1172	0	1134	19	0
1	B	1172	0	1133	9	0
1	C	1178	0	1138	16	0
1	D	1173	0	1134	7	0
2	A	1	0	0	0	0
2	B	1	0	0	0	0
2	C	1	0	0	0	0
2	D	1	0	0	0	0
3	A	1	0	0	0	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	D	1	0	0	0	0
4	A	51	0	0	0	0
4	B	72	0	0	2	0
4	C	45	0	0	0	0
4	D	30	0	0	1	0
All	All	4899	0	4539	48	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 5.

All (48) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:87:TYR:O	1:D:146:ARG:NH2	2.06	0.89
1:A:87:TYR:O	1:A:146:ARG:NH2	2.12	0.81
1:C:87:TYR:O	1:C:146:ARG:NH2	2.12	0.80
1:B:87:TYR:O	1:B:146:ARG:NH2	2.17	0.78
1:C:95:ARG:NH2	1:C:143:PHE:O	2.28	0.66
1:A:104:PRO:HB3	1:C:121:PRO:HG2	1.79	0.65
1:A:98:LEU:HD13	1:A:159:LEU:HD11	1.84	0.60
1:D:139:ASP:O	1:D:146:ARG:NH1	2.35	0.60
1:A:95:ARG:HD2	1:A:97:SER:O	2.07	0.55
1:A:104:PRO:HB3	1:C:121:PRO:CG	2.37	0.54
1:D:183:ARG:NH2	4:D:428:HOH:O	2.41	0.53
1:B:99:VAL:HG12	1:B:103:THR:HG21	1.93	0.51
1:D:114:VAL:HG22	1:D:225:PHE:HB2	1.93	0.50
1:B:77:ASP:N	1:C:76:SER:HG	2.10	0.50
1:C:98:LEU:HD13	1:C:159:LEU:HD11	1.94	0.49
1:A:170:ARG:CZ	1:A:203:LEU:HD12	2.42	0.48
1:A:211:ASN:O	1:A:221:GLY:HA2	2.14	0.48
1:B:95:ARG:NH2	1:B:143:PHE:O	2.48	0.47
1:C:149:GLN:NE2	1:C:158:THR:O	2.46	0.46
1:B:211:ASN:O	1:B:221:GLY:HA2	2.15	0.46
1:C:114:VAL:HG22	1:C:225:PHE:HB2	1.98	0.46
1:A:183:ARG:HG3	1:A:229:VAL:HG21	1.97	0.45
1:B:114:VAL:HG22	1:B:225:PHE:HB2	1.98	0.45
1:A:151:THR:HG22	1:A:157:PHE:HB3	1.99	0.44
1:A:96:THR:HG22	1:A:149:GLN:HB3	1.99	0.43
1:C:96:THR:HG22	1:C:149:GLN:HB3	2.00	0.43
1:D:79:THR:HA	1:D:80:PRO:HD3	1.82	0.43
1:A:107:PRO:HB2	1:A:218:GLY:HA3	1.99	0.43

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:93:PRO:HA	1:D:94:PRO:HD3	1.87	0.43
1:B:203:LEU:HD23	1:B:203:LEU:HA	1.79	0.43
1:C:113:TYR:HB2	1:C:115:PHE:CE2	2.53	0.43
1:C:189:GLN:O	1:C:208:LEU:HB3	2.19	0.42
1:A:144:ALA:O	1:A:146:ARG:HG3	2.20	0.42
1:A:89:LYS:HB2	1:A:146:ARG:NE	2.35	0.42
1:C:177:LYS:HA	1:C:186:LEU:O	2.19	0.42
1:C:77:ASP:HA	1:C:78:PRO:HD3	1.86	0.42
1:D:172:ARG:HE	1:D:195:GLU:CD	2.24	0.41
1:B:79:THR:HA	1:B:80:PRO:HD3	1.85	0.41
1:A:122:LEU:O	1:A:155:GLY:HA2	2.21	0.41
1:C:108:LEU:HA	1:C:219:ARG:O	2.20	0.41
1:A:93:PRO:HA	1:A:94:PRO:HD3	1.94	0.41
1:C:89:LYS:HB2	1:C:146:ARG:NE	2.36	0.41
1:A:142:GLY:HA3	4:B:456:HOH:O	2.21	0.41
1:C:170:ARG:CZ	1:C:203:LEU:HD12	2.51	0.41
1:A:114:VAL:HG12	1:A:122:LEU:HD12	2.03	0.40
1:B:211:ASN:HA	4:B:442:HOH:O	2.22	0.40
1:A:126:LEU:O	1:A:178:ALA:HA	2.21	0.40
1:A:212:VAL:HG11	1:A:219:ARG:NH1	2.37	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	152/155 (98%)	149 (98%)	3 (2%)	0	100	100
1	B	152/155 (98%)	149 (98%)	3 (2%)	0	100	100
1	C	153/155 (99%)	150 (98%)	3 (2%)	0	100	100
1	D	152/155 (98%)	149 (98%)	3 (2%)	0	100	100
All	All	609/620 (98%)	597 (98%)	12 (2%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	125/126 (99%)	123 (98%)	2 (2%)	62	77
1	B	125/126 (99%)	123 (98%)	2 (2%)	62	77
1	C	126/126 (100%)	123 (98%)	3 (2%)	49	64
1	D	126/126 (100%)	122 (97%)	4 (3%)	39	53
All	All	502/504 (100%)	491 (98%)	11 (2%)	52	66

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	77	ASP
1	A	229	VAL
1	B	77	ASP
1	B	229	VAL
1	C	77	ASP
1	C	101	SER
1	C	229	VAL
1	D	76	SER
1	D	77	ASP
1	D	95	ARG
1	D	229	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	211	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

Of 6 ligands modelled in this entry, 6 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	154/155 (99%)	0.14	0 100 100	23, 31, 45, 53	0
1	B	154/155 (99%)	0.16	1 (0%) 89 92	24, 33, 47, 54	0
1	C	155/155 (100%)	0.36	7 (4%) 33 40	25, 35, 54, 64	0
1	D	154/155 (99%)	0.65	11 (7%) 16 19	29, 43, 60, 68	0
All	All	617/620 (99%)	0.33	19 (3%) 49 56	23, 35, 54, 68	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	139	ASP	5.1
1	D	217	PRO	4.8
1	C	76	SER	3.7
1	C	83	MET	3.6
1	D	152	GLY	3.2
1	D	101	SER	3.2
1	D	143	PHE	3.0
1	C	217	PRO	2.7
1	B	139	ASP	2.7
1	D	227	LEU	2.7
1	D	76	SER	2.6
1	C	215	ALA	2.5
1	D	229	VAL	2.5
1	D	215	ALA	2.4
1	D	127	LEU	2.4
1	C	214	SER	2.3
1	D	122	LEU	2.3
1	C	230	ALA	2.0
1	D	91	ASP	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
3	CL	D	302	1/1	0.96	0.07	31,31,31,31	0
3	CL	A	302	1/1	0.96	0.09	32,32,32,32	0
2	FE	D	301	1/1	0.98	0.09	43,43,43,43	0
2	FE	A	301	1/1	0.98	0.10	25,25,25,25	0
2	FE	B	301	1/1	0.99	0.10	29,29,29,29	0
2	FE	C	301	1/1	0.99	0.09	28,28,28,28	0

6.5 Other polymers [i](#)

There are no such residues in this entry.