



# Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 09:36 pm GMT

PDB ID : 1IX9  
Title : Crystal Structure of the E. coli Manganase(III) superoxide dismutase mutant Y174F at 0.90 angstroms resolution.  
Authors : Anderson, B.F.; Edwards, R.A.; Whittaker, M.M.; Whittaker, J.W.; Baker, E.N.; Jameson, G.B.  
Deposited on : 2002-06-17  
Resolution : 0.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : trunk30967  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : trunk30967


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 0.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1087 (1.06-0.74)
Clashscore	122126	1156 (1.06-0.74)
Ramachandran outliers	120053	1069 (1.06-0.74)
Sidechain outliers	120020	1071 (1.06-0.74)
RSRZ outliers	108989	1054 (1.06-0.74)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	205	 87% 12% •
1	B	205	 2% 81% 17% ••

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4379 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Superoxide Dismutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	205	1741	1119	299	320	3	0	21	0
1	B	205	1736	1116	299	318	3	0	20	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	174	PHE	TYR	ENGINEERED	UNP P00448
B	174	PHE	TYR	ENGINEERED	UNP P00448

- Molecule 2 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	Mn		
2	B	1	2	2	0	1
2	A	1	2	2	0	1


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	470	470	470	0	9
3	B	428	428	428	0	3

### 3 Residue-property plots


These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Superoxide Dismutase

Chain A:  87% 12%



- Molecule 1: Superoxide Dismutase

Chain B:  81% 17% 2%



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	46.89Å 46.02Å 95.99Å 90.00° 98.40° 90.00°	Depositor
Resolution (Å)	30.00 – 0.90 47.48 – 0.90	Depositor EDS
% Data completeness (in resolution range)	95.2 (30.00-0.90) 96.2 (47.48-0.90)	Depositor EDS
$R_{merge}$	0.03	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.95 (at 0.90Å)	Xtrriage
Refinement program	SHELXL-97	Depositor
R, $R_{free}$	0.107 , 0.127 0.115 , 0.133	Depositor DCC
$R_{free}$ test set	2940 reflections (1.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	7.9	Xtrriage
Anisotropy	0.519	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 69.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	4379	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	13.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.00% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: MN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.74	1/1867 (0.1%)	1.15	23/2514 (0.9%)
1	B	0.84	1/1843 (0.1%)	1.34	29/2486 (1.2%)
All	All	0.79	2/3710 (0.1%)	1.25	52/5000 (1.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	2

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	6	SER	CB-OG	-6.23	1.34	1.42
1	B	142	SER	CB-OG	-5.54	1.35	1.42

All (52) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	204	LYS	CA-C-O	-9.87	99.38	120.10
1	B	90	LYS	O-C-N	8.93	138.38	123.20
1	B	204	LYS	O-C-N	8.32	136.01	122.70
1	A	136[A]	ASP	CB-CG-OD1	8.08	125.57	118.30
1	A	136[B]	ASP	CB-CG-OD1	8.08	125.57	118.30
1	A	180[A]	ARG	NE-CZ-NH2	7.96	124.28	120.30
1	A	180[B]	ARG	NE-CZ-NH2	7.96	124.28	120.30
1	B	180[A]	ARG	NE-CZ-NH1	7.64	124.12	120.30
1	B	180[B]	ARG	NE-CZ-NH1	7.64	124.12	120.30

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	90	LYS	CA-C-N	-7.63	100.95	116.20
1	B	204	LYS	C-N-CA	-7.53	102.86	121.70
1	B	200	ARG	NE-CZ-NH2	-6.88	116.86	120.30
1	B	180[A]	ARG	NE-CZ-NH2	-6.67	116.96	120.30
1	B	180[B]	ARG	NE-CZ-NH2	-6.67	116.96	120.30
1	A	104	ARG	NE-CZ-NH1	6.50	123.55	120.30
1	B	136[A]	ASP	N-CA-CB	-6.07	99.68	110.60
1	B	136[B]	ASP	N-CA-CB	-6.07	99.68	110.60
1	B	136[A]	ASP	N-CA-C	6.01	127.24	111.00
1	B	136[B]	ASP	N-CA-C	6.01	127.24	111.00
1	B	203	ALA	O-C-N	5.91	132.15	122.70
1	A	135[A]	GLY	CA-C-O	5.86	131.15	120.60
1	A	135[B]	GLY	CA-C-O	5.86	131.15	120.60
1	A	66	ASP	CB-CG-OD1	-5.80	113.08	118.30
1	B	205	LYS	N-CA-C	-5.76	95.46	111.00
1	A	58	THR	O-C-N	-5.75	113.50	122.70
1	B	135[A]	GLY	CA-C-O	5.61	130.70	120.60
1	B	135[B]	GLY	CA-C-O	5.61	130.70	120.60
1	A	110[A]	ASP	CB-CG-OD2	-5.56	113.30	118.30
1	A	110[B]	ASP	CB-CG-OD2	-5.56	113.30	118.30
1	B	136[A]	ASP	CA-C-O	5.52	131.70	120.10
1	B	136[B]	ASP	CA-C-O	5.52	131.70	120.10
1	B	136[A]	ASP	CB-CG-OD2	-5.48	113.37	118.30
1	B	136[B]	ASP	CB-CG-OD2	-5.48	113.37	118.30
1	A	186[A]	LYS	CG-CD-CE	5.48	128.33	111.90
1	A	186[B]	LYS	CG-CD-CE	5.48	128.33	111.90
1	A	186[C]	LYS	CG-CD-CE	5.48	128.33	111.90
1	B	59[A]	LYS	N-CA-CB	5.45	120.41	110.60
1	B	59[B]	LYS	N-CA-CB	5.45	120.41	110.60
1	B	112	PHE	CB-CG-CD2	-5.42	117.00	120.80
1	B	2	TYR	CB-CG-CD1	5.36	124.22	121.00
1	A	135[A]	GLY	CA-C-N	-5.32	105.50	117.20
1	A	135[B]	GLY	CA-C-N	-5.32	105.50	117.20
1	B	65	ALA	CB-CA-C	5.32	118.07	110.10
1	A	47	GLU	CB-CG-CD	5.27	128.44	114.20
1	B	203	ALA	CB-CA-C	5.25	117.98	110.10
1	B	203	ALA	N-CA-CB	-5.21	102.81	110.10
1	A	2	TYR	CB-CG-CD1	5.18	124.11	121.00
1	A	1[A]	SER	N-CA-CB	-5.11	102.83	110.50
1	A	1[B]	SER	N-CA-CB	-5.11	102.83	110.50
1	A	167	ASP	CB-CG-OD2	-5.11	113.70	118.30
1	A	161	PHE	CB-CG-CD1	5.02	124.31	120.80

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	142	SER	N-CA-CB	5.01	118.01	110.50

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	135[A]	GLY	Peptide
1	B	136[A]	ASP	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1741	0	1731	36	0
1	B	1736	0	1723	39	1
2	A	2	0	0	0	0
2	B	2	0	0	0	0
3	A	470	0	0	26	0
3	B	428	0	0	12	0
All	All	4379	0	3454	71	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (71) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:68[B]:LYS:HE3	3:B:487:HOH:O	1.55	1.05
1:B:135[B]:GLY:O	1:B:136[B]:ASP:OD1	1.80	0.99
1:A:186[C]:LYS:CE	3:A:402:HOH:O	2.11	0.98
1:A:186[A]:LYS:HE2	3:A:530:HOH:O	1.65	0.97
1:B:20[B]:LYS:HE2	3:B:442:HOH:O	1.68	0.94
1:B:20[C]:LYS:HG2	3:B:339:HOH:O	1.67	0.94
1:B:60:LEU:O	1:B:68[B]:LYS:NZ	2.01	0.92
1:A:68[C]:LYS:CG	3:A:297:HOH:O	2.19	0.89

Continued on next page...



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:186[C]:LYS:CD	3:A:402:HOH:O	2.22	0.85
1:A:179[B]:ASN:OD1	3:A:304:HOH:O	1.97	0.83
1:A:68[C]:LYS:HG3	3:A:297:HOH:O	1.81	0.80
1:B:20[B]:LYS:HG2	3:B:339:HOH:O	1.81	0.79
1:A:186[C]:LYS:HE3	3:A:402:HOH:O	1.81	0.77
1:A:205[A]:LYS:HE2	3:A:405:HOH:O	1.85	0.77
1:A:50:ASN:HB2	3:A:376:HOH:O	1.91	0.69
1:A:68[C]:LYS:HG2	3:A:297:HOH:O	1.88	0.68
1:A:29[B]:LYS:NZ	1:B:178:GLN:HE22	1.96	0.63
1:B:1[B]:SER:HB2	3:B:311:HOH:O	2.00	0.61
1:A:205[A]:LYS:CE	3:A:405:HOH:O	2.48	0.59
1:A:186[C]:LYS:HD2	3:A:402:HOH:O	1.96	0.58
1:B:133:LEU:CD1	1:B:136[A]:ASP:HA	2.34	0.57
1:A:68[B]:LYS:HD2	3:A:498:HOH:O	2.04	0.57
1:B:201:PHE:O	1:B:204:LYS:HG2	2.06	0.55
1:A:29[A]:LYS:HE3	3:A:613:HOH:O	2.08	0.54
1:B:135[B]:GLY:O	1:B:136[B]:ASP:HB2	2.08	0.53
1:B:133:LEU:HG	1:B:136[A]:ASP:HA	1.91	0.53
1:A:68[C]:LYS:HZ1	1:A:155:ILE:HD11	1.75	0.51
1:A:50:ASN:ND2	3:A:542:HOH:O	2.29	0.51
1:B:153:GLU:HG2	1:B:159:SER:HB3	1.93	0.51
1:B:135[B]:GLY:C	1:B:136[B]:ASP:OD1	2.47	0.51
1:A:68[C]:LYS:NZ	1:A:155:ILE:HD11	2.26	0.51
1:B:39:ASN:O	1:B:43[B]:GLU:HG3	2.11	0.50
1:B:135[B]:GLY:O	1:B:136[B]:ASP:CB	2.58	0.50
1:B:1[B]:SER:N	1:B:39:ASN:OD1	2.40	0.50
1:A:115[B]:GLU:CG	3:A:565:HOH:O	2.58	0.50
1:A:68[A]:LYS:CD	3:A:498:HOH:O	2.59	0.49
1:A:115[B]:GLU:OE2	3:A:453:HOH:O	2.18	0.49
1:B:135[B]:GLY:O	1:B:136[B]:ASP:CG	2.50	0.49
1:B:149[B]:PRO:HG2	1:B:162:PRO:HG3	1.93	0.49
1:B:61:ASP:HA	1:B:68[B]:LYS:NZ	2.27	0.49
1:B:68[B]:LYS:HE2	3:B:494:HOH:O	2.13	0.48
1:B:43[B]:GLU:CD	3:B:271:HOH:O	2.52	0.48
1:B:179[B]:ASN:ND2	3:B:411:HOH:O	2.46	0.48
1:B:91[A]:GLY:C	3:B:475:HOH:O	2.51	0.48
1:A:68[A]:LYS:HD2	3:A:498:HOH:O	2.14	0.47
1:B:99[B]:LYS:NZ	1:B:103:GLU:OE2	2.38	0.47
1:B:61:ASP:HA	1:B:68[B]:LYS:CE	2.45	0.47
1:A:65:ALA:HA	1:A:68[A]:LYS:HE3	1.97	0.46
1:A:68[A]:LYS:CE	3:A:498:HOH:O	2.63	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:68[C]:LYS:NZ	3:A:531:HOH:O	2.46	0.46
1:A:186[B]:LYS:HD3	3:A:402:HOH:O	2.14	0.46
1:B:133:LEU:HD11	1:B:136[A]:ASP:CA	2.46	0.46
1:A:186[A]:LYS:CE	3:A:530:HOH:O	2.42	0.46
1:B:132:VAL:CG2	1:B:134[B]:LYS:HD2	2.46	0.45
1:B:135[B]:GLY:C	1:B:136[B]:ASP:CG	2.70	0.45
1:A:29[B]:LYS:HZ3	1:B:178:GLN:HE22	1.65	0.45
1:B:115[B]:GLU:HG2	3:B:268:HOH:O	2.17	0.44
1:A:99[B]:LYS:NZ	3:A:456:HOH:O	2.44	0.44
1:A:179[B]:ASN:HB3	1:B:30:HIS:CD2	2.53	0.44
1:B:132:VAL:HG23	1:B:134[B]:LYS:HD2	2.01	0.42
1:A:117[A]:GLU:OE1	3:A:638:HOH:O	2.22	0.42
1:B:9:TYR:HB2	1:B:13:ALA:CB	2.50	0.41
1:A:68[C]:LYS:HZ3	1:A:155:ILE:CD1	2.33	0.41
1:A:29[B]:LYS:CE	1:B:178:GLN:HE22	2.33	0.41
1:A:63:LEU:O	1:A:68[B]:LYS:NZ	2.54	0.41
1:B:180[B]:ARG:NH2	3:B:413:HOH:O	2.53	0.41
1:B:133:LEU:HD21	1:B:204:LYS:HD3	2.03	0.41
1:B:132:VAL:HG23	1:B:134[B]:LYS:CD	2.51	0.41
1:B:134[B]:LYS:CE	3:B:397:HOH:O	2.69	0.41
1:A:179[A]:ASN:ND2	3:A:229:HOH:O	2.54	0.40
1:A:68[C]:LYS:NZ	1:A:155:ILE:CD1	2.85	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:6:SER:OG	1:B:136[A]:ASP:OD2[2_646]	2.03	0.17

## 5.3 Torsion angles

### 5.3.1 Protein backbone

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	224/205 (109%)	219 (98%)	5 (2%)	0	100	100
1	B	223/205 (109%)	217 (97%)	4 (2%)	2 (1%)	19	2
All	All	447/410 (109%)	436 (98%)	9 (2%)	2 (0%)	49	11

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	136[A]	ASP
1	B	136[B]	ASP

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	188/166 (113%)	185 (98%)	3 (2%)	65	29
1	B	185/166 (111%)	182 (98%)	3 (2%)	65	29
All	All	373/332 (112%)	367 (98%)	6 (2%)	74	29

All (6) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	1[A]	SER
1	A	1[B]	SER
1	A	164	MET
1	B	136[A]	ASP
1	B	136[B]	ASP
1	B	164	MET

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	50	ASN
1	A	62	GLN
1	B	178	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

### 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled '#RSRZ > 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q < 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	205/205 (100%)	-0.71	0 <b>100</b> <b>100</b>	5, 8, 19, 29	0
1	B	205/205 (100%)	-0.57	4 (1%) 65 48	6, 11, 21, 32	2 (0%)
All	All	410/410 (100%)	-0.64	4 (0%) <b>82</b> <b>68</b>	5, 10, 21, 32	2 (0%)

All (4) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	204	LYS	3.7
1	B	136[A]	ASP	2.8
1	B	205	LYS	2.6
1	B	135[A]	GLY	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q < 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MN	B	206[A]	1/1	1.00	0.02	6,6,6,6	1

*Continued on next page...*

*Continued from previous page...*

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	MN	B	206[B]	1/1	1.00	0.02	6,6,6,6	1
2	MN	A	206[A]	1/1	1.00	0.03	5,5,5,5	1
2	MN	A	206[B]	1/1	1.00	0.03	5,5,5,5	1

## 6.5 Other polymers [i](#)

There are no such residues in this entry.