

Full wwPDB X-ray Structure Validation Report (i)

Jan 29, 2024 – 04:10 PM JST

PDB ID : 8IY1

Title: Structure of Acb2 complexed with cAAG

Authors: Cao, X.L.; Xiao, Y.; Feng, Y.

Deposited on : 2023-04-03

Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.36

buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.36

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$egin{aligned} ext{Similar resolution} \ (\# ext{Entries, resolution range}(ext{Å})) \end{aligned}$		
R_{free}	130704	5197 (2.10-2.10)		
Clashscore	141614	5710 (2.10-2.10)		
Ramachandran outliers	138981	5647 (2.10-2.10)		
Sidechain outliers	138945	5648 (2.10-2.10)		
RSRZ outliers	127900	5083 (2.10-2.10)		

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	A	92	93%	7%
1	Λ	32	16%	7 %
1	В	92	96%	•
1	С	92	93%	5% •
1	D	92	91%	8% •
1	Е	92	96%	
1	F	92	16%	:



2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 5037 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called p26.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	92	Total	С	N	О	S	0	0	0
1	A	92	740	456	138	142	4	U	U	U
1	В	92	Total	С	N	О	S	0	0	0
1	Ъ	92	740	456	138	142	4	0	U	
1	C	91	Total	С	N	О	S	0	0	0
1		91	732	452	137	139	4	0		
1	D	91	Total	С	N	О	S	0	0	0
1	D	91	732	452	137	139	4	0	U	
1	Е	91	Total	С	N	О	S	0	0	0
1	ш	91	732	452	137	139	4	U	U	U
1	F	91	Total	С	N	О	S	0	0	0
1	I.	91	732	452	137	139	4	U	U	U

• Molecule 2 is [(2R,3S,4R,5R)-5-(6-aminopurin-9-yl)-4-hydroxy-2-(hydroxymethyl)oxolan -3-yl] dihydrogen phosphate (three-letter code: 3AM) (formula: $C_{10}H_{14}N_5O_7P$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues		Ato	ms			ZeroOcc	AltConf
2	A	1	Total	С	N	О	Р	0	0
2	A	1	22	10	5	6	1	U	0
2	٨	1	Total	С	N	О	Р	0	0
2	A	1	22	10	5	6	1	0	U
2	В	1	Total	С	N	О	Р	0	0
2	D	1	22	10	5	6	1	0	0
2	D	1	Total	С	N	О	Р	0	0
2	В	1	22	10	5	6	1	0	0
0	С	1	Total	С	N	О	Р	0	0
2		1	22	10	5	6	1	0	0
2	С	1	Total	С	N	О	Р	0	0
2		1	22	10	5	6	1	U	
2	D	1	Total	С	N	О	Р	0	0
2	D	1	22	10	5	6	1	U	0
2	D	1	Total	С	N	О	Р	0	0
2	D	1	22	10	5	6	1	U	0
2	E	1	Total	С	N	О	Р	0	0
2	E	1	22	10	5	6	1	U	0
2	E	1	Total	С	N	О	Р	0	0
2	E	1	22	10	5	6	1	U	0
2	F	1	Total	С	N	О	Р	0	0
	Г	1	22	10	5	6	1	0	0
2	F	1	Total	С	N	О	Р	0	0
	I.	1	22	10	5	6	1	U	0

 \bullet Molecule 3 is GUANOSINE-3'-MONOPHOSPHATE (three-letter code: 3GP) (formula: $C_{10}H_{14}N_5O_8P)$ (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues		Ato	ms			ZeroOcc	AltConf
3	A	1	Total	С	N	О	Р	0	0
3	A	1	23	10	5	7	1	U	0
3	В	1	Total	С	N	О	Р	0	0
3	Б	1	23	10	5	7	1	0	0
3	С	1	Total	С	N	О	Р	0	0
3			23	10	5	7	1	0	
3	D	1	Total	С	N	О	Р	0	0
3	D	1	23	10	5	7	1	0	0
3	Е	1	Total	С	N	О	Р	0	0
3	ت ا	1	23	10	5	7	1	0	0
3	F	1	Total	С	N	О	Р	0	0
3	l L	1	23	10	5	7	1	U	


• Molecule 4 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	52	Total O 52 52	0	0
4	В	38	Total O 38 38	0	0
4	С	52	Total O 52 52	0	0
4	D	45	Total O 45 45	0	0
4	E	19	Total O 19 19	0	0
4	F	21	Total O 21 21	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.









4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 3 2 1	Depositor
Cell constants	103.59Å 103.59Å 101.44Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Resolution (Å)	28.68 - 2.10	Depositor
Resolution (A)	28.68 - 2.10	EDS
% Data completeness	97.7 (28.68-2.10)	Depositor
(in resolution range)	97.7 (28.68-2.10)	EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	3.66 (at 2.10Å)	Xtriage
Refinement program	PHENIX (1.20_4459: ???)	Depositor
D D	0.221 , 0.248	Depositor
R, R_{free}	0.222 , 0.233	DCC
R_{free} test set	1805 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å ²)	31.8	Xtriage
Anisotropy	0.243	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.39, 60.9	EDS
L-test for twinning ²	$< L > = 0.50, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	0.076 for -h,-k,l	Xtriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	5037	wwPDB-VP
Average B, all atoms (Å ²)	45.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 11.44% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: 3GP, 3AM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond	lengths	Bond angles		
WIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.51	0/750	0.64	0/1005	
1	В	0.74	0/750	0.68	0/1005	
1	С	0.67	0/742	0.63	0/994	
1	D	0.69	0/742	0.70	1/994 (0.1%)	
1	Е	0.71	0/742	0.59	0/994	
1	F	0.72	0/742	0.59	0/994	
All	All	0.68	0/4468	0.64	1/5986 (0.0%)	

There are no bond length outliers.

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(^{o})$	$\operatorname{Ideal}({}^{o})$
1	D	11	TYR	CB-CA-C	-5.44	99.51	110.40

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	740	0	732	2	0
1	В	740	0	732	1	0
1	С	732	0	728	4	0
1	D	732	0	728	2	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	$\mathbf{H}(\mathbf{added})$	Clashes	Symm-Clashes
1	Е	732	0	728	1	0
1	F	732	0	728	0	0
2	A	44	0	22	1	0
2	В	44	0	22	2	0
2	С	44	0	22	0	0
2	D	44	0	22	0	0
2	Ε	44	0	22	0	0
2	F	44	0	22	0	0
3	A	23	0	11	1	0
3	В	23	0	11	5	0
3	С	23	0	11	0	0
3	D	23	0	11	0	0
3	Ε	23	0	11	0	0
3	F	23	0	11	0	0
4	A	52	0	0	1	3
4	В	38	0	0	3	0
4	С	52	0	0	2	1
4	D	45	0	0	0	1
4	Е	19	0	0	0	0
4	F	21	0	0	0	0
All	All	5037	0	4574	15	4

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
1:C:20:ASP:OD1	4:C:201:HOH:O	2.05	0.72
1:C:17:GLU:OE2	4:C:202:HOH:O	2.11	0.66
2:B:101:3AM:H5'	3:B:103:3GP:H3'	1.82	0.62
2:B:101:3AM:H8	3:B:103:3GP:O2P	1.99	0.62
2:A:101:3AM:H8	3:A:103:3GP:O2P	2.06	0.56
1:E:5:HIS:HA	1:E:8:ILE:HD12	1.88	0.56
1:A:44:ARG:NE	4:A:201:HOH:O	2.38	0.54
1:C:8:ILE:HG12	1:D:19:ILE:HG23	1.91	0.52
3:B:103:3GP:O2'	4:B:201:HOH:O	2.14	0.52
3:B:103:3GP:O2P	4:B:201:HOH:O	2.19	0.51
1:B:5:HIS:HB3	1:B:11:TYR:CD2	2.47	0.50
1:D:22:MET:HE3	1:D:84:ILE:HG23	1.98	0.45
3:B:103:3GP:O3P	4:B:202:HOH:O	2.21	0.43



Continued from previous page...

Atom-1	Atom-1 Atom-2		$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:C:4:GLN:HA	1:C:7:LYS:HE3	2.01	0.43
1:A:12:ARG:O	1:A:14:LEU:HD22	2.20	0.41

All (4) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	Clash overlap (Å)
4:A:238:HOH:O	4:A:250:HOH:O[5_556]	2.05	0.15
	4:D:224:HOH:O[3_545]	2.06	0.14
4:A:235:HOH:O	4:A:237:HOH:O[5_556]	2.16	0.04
4:A:222:HOH:O	4:A:223:HOH:O[2_555]	2.18	0.02

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	90/92~(98%)	90 (100%)	0	0	100	100
1	В	90/92 (98%)	89 (99%)	0	1 (1%)	14	9
1	С	$89/92\ (97\%)$	89 (100%)	0	0	100	100
1	D	89/92 (97%)	88 (99%)	1 (1%)	0	100	100
1	E	$89/92\ (97\%)$	89 (100%)	0	0	100	100
1	F	89/92 (97%)	89 (100%)	0	0	100	100
All	All	536/552 (97%)	534 (100%)	1 (0%)	1 (0%)	47	49

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	6	LYS



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	79/79 (100%)	76 (96%)	3 (4%)	33 34
1	В	79/79 (100%)	78 (99%)	1 (1%)	69 75
1	С	78/79 (99%)	78 (100%)	0	100 100
1	D	78/79 (99%)	75 (96%)	3 (4%)	33 34
1	E	78/79 (99%)	77 (99%)	1 (1%)	69 75
1	F	78/79 (99%)	77 (99%)	1 (1%)	69 75
All	All	470/474 (99%)	461 (98%)	9 (2%)	57 63

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	2	ASP
1	A	13	ASP
1	A	32	PHE
1	В	15	SER
1	D	16	GLN
1	D	17	GLU
1	D	32	PHE
1	Е	13	ASP
1	F	37	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	23	ASN
1	Е	31	GLN
1	F	5	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.



5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

18 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Trunc	Chain	Res	Link	Во	ond leng	ths	В	Bond angles		
Mol	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
3	3GP	Е	103	-	18,25,26	0.98	1 (5%)	19,37,40	1.24	2 (10%)	
2	3AM	В	102	-	18,24,25	1.02	1 (5%)	18,35,38	1.41	2 (11%)	
2	3AM	В	101	-	18,24,25	1.07	2 (11%)	18,35,38	2.15	7 (38%)	
2	3AM	D	102	-	18,24,25	1.06	1 (5%)	18,35,38	1.31	2 (11%)	
2	3AM	Е	101	-	18,24,25	1.01	1 (5%)	18,35,38	1.39	3 (16%)	
2	3AM	D	101	-	18,24,25	1.01	1 (5%)	18,35,38	1.47	2 (11%)	
2	3AM	A	101	-	18,24,25	1.27	2 (11%)	18,35,38	2.05	6 (33%)	
3	3GP	В	103	-	18,25,26	0.97	1 (5%)	19,37,40	1.28	3 (15%)	
3	3GP	С	103	-	18,25,26	1.04	1 (5%)	19,37,40	1.23	2 (10%)	
2	3AM	Е	102	-	18,24,25	1.07	1 (5%)	18,35,38	1.39	3 (16%)	
3	3GP	F	103	-	18,25,26	0.98	1 (5%)	19,37,40	1.25	2 (10%)	
2	3AM	A	102	-	18,24,25	1.05	1 (5%)	18,35,38	1.47	3 (16%)	
3	3GP	A	103	-	18,25,26	1.06	1 (5%)	19,37,40	1.42	3 (15%)	
3	3GP	D	103	-	18,25,26	1.04	1 (5%)	19,37,40	1.17	3 (15%)	
2	3AM	С	101	_	18,24,25	1.02	1 (5%)	18,35,38	1.33	2 (11%)	
2	3AM	F	102	-	18,24,25	1.03	1 (5%)	18,35,38	1.36	3 (16%)	
2	3AM	F	101	-	18,24,25	0.98	1 (5%)	18,35,38	1.37	3 (16%)	
2	3AM	С	102	-	18,24,25	1.05	1 (5%)	18,35,38	1.29	2 (11%)	



In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	3GP	Ε	103	-	-	2/4/26/27	0/3/3/3
2	3AM	В	102	-	-	0/4/26/27	0/3/3/3
2	3AM	В	101	-	-	2/4/26/27	0/3/3/3
2	3AM	D	102	-	-	0/4/26/27	0/3/3/3
2	3AM	${ m E}$	101	-	-	2/4/26/27	0/3/3/3
2	3AM	D	101	-	-	2/4/26/27	0/3/3/3
2	3AM	A	101	-	-	2/4/26/27	0/3/3/3
3	3GP	В	103	-	-	2/4/26/27	0/3/3/3
3	3GP	С	103	-	-	0/4/26/27	0/3/3/3
2	3AM	Е	102	-	-	0/4/26/27	0/3/3/3
3	3GP	F	103	-	-	2/4/26/27	0/3/3/3
2	3AM	A	102	-	-	0/4/26/27	0/3/3/3
3	3GP	A	103	-	-	2/4/26/27	0/3/3/3
3	3GP	D	103	-	-	0/4/26/27	0/3/3/3
2	3AM	С	101	-	-	0/4/26/27	0/3/3/3
2	3AM	F	102	-	-	0/4/26/27	0/3/3/3
2	3AM	F	101	-	-	2/4/26/27	0/3/3/3
2	3AM	С	102	-	-	0/4/26/27	0/3/3/3

All (20) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	Ideal(A)
2	A	101	3AM	O4'-C1'	3.29	1.45	1.41
3	С	103	3GP	C6-N1	-2.73	1.33	1.37
3	A	103	3GP	C6-N1	-2.65	1.33	1.37
3	D	103	3GP	C6-N1	-2.48	1.34	1.37
3	В	103	3GP	C6-N1	-2.44	1.34	1.37
2	Е	102	3AM	C5-C4	2.42	1.47	1.40
2	Е	101	3AM	C5-C4	2.42	1.47	1.40
2	F	102	3AM	C5-C4	2.41	1.47	1.40
2	С	101	3AM	C5-C4	2.37	1.47	1.40
2	D	101	3AM	C5-C4	2.36	1.47	1.40
2	С	102	3AM	C5-C4	2.35	1.47	1.40
2	D	102	3AM	C5-C4	2.34	1.47	1.40
2	F	101	3AM	C5-C4	2.34	1.47	1.40
2	A	101	3AM	C5-C4	2.32	1.47	1.40
3	F	103	3GP	C6-N1	-2.28	1.34	1.37



Continued from previous page...

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\text{\AA})$	$\operatorname{Ideal}(\text{\AA})$
2	В	102	3AM	C5-C4	2.23	1.46	1.40
2	В	101	3AM	C5-C4	2.21	1.46	1.40
2	A	102	3AM	C5-C4	2.18	1.46	1.40
2	В	101	3AM	O4'-C1'	2.15	1.44	1.41
3	Е	103	3GP	C6-N1	-2.15	1.34	1.37

All (53) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$\mathbf{Observed}(^{o})$	$Ideal(^{o})$
2	A	101	3AM	O4'-C4'-C5'	4.16	118.21	109.21
2	В	101	3AM	O5'-C5'-C4'	4.02	125.09	111.29
2	В	101	3AM	N3-C2-N1	-3.75	122.81	128.68
2	В	101	3AM	C5'-C4'-C3'	-3.65	103.20	114.85
2	A	102	3AM	N3-C2-N1	-3.60	123.06	128.68
2	Е	101	3AM	N3-C2-N1	-3.56	123.11	128.68
2	F	101	3AM	N3-C2-N1	-3.56	123.12	128.68
2	В	102	3AM	N3-C2-N1	-3.54	123.14	128.68
2	A	101	3AM	O4'-C1'-C2'	-3.53	101.77	106.93
2	Е	102	3AM	N3-C2-N1	-3.51	123.19	128.68
2	F	102	3AM	N3-C2-N1	-3.48	123.24	128.68
2	В	101	3AM	O4'-C4'-C5'	3.44	116.65	109.21
2	A	101	3AM	N3-C2-N1	-3.44	123.30	128.68
2	D	102	3AM	N3-C2-N1	-3.24	123.61	128.68
2	С	101	3AM	N3-C2-N1	-3.24	123.62	128.68
2	D	101	3AM	N3-C2-N1	-3.22	123.65	128.68
2	A	101	3AM	C4-C5-N7	-3.17	106.09	109.40
2	С	102	3AM	N3-C2-N1	-3.16	123.74	128.68
2	A	101	3AM	C1'-N9-C4	-2.71	121.89	126.64
2	В	101	3AM	C4-C5-N7	-2.67	106.62	109.40
2	С	101	3AM	C4-C5-N7	-2.65	106.64	109.40
2	A	102	3AM	C4-C5-N7	-2.61	106.68	109.40
2	С	102	3AM	C4-C5-N7	-2.60	106.69	109.40
2	A	101	3AM	C5'-C4'-C3'	-2.57	106.64	114.85
3	A	103	3GP	C5-C6-N1	2.53	118.42	113.95
3	С	103	3GP	C5-C6-N1	2.52	118.40	113.95
2	В	101	3AM	O4'-C1'-C2'	-2.50	103.28	106.93
2	F	102	3AM	C4-C5-N7	-2.48	106.82	109.40
3	В	103	3GP	C5-C6-N1	2.48	118.33	113.95
2	В	101	3AM	C1'-N9-C4	-2.42	122.40	126.64
2	Е	102	3AM	C4-C5-N7	-2.40	106.90	109.40
3	С	103	3GP	C8-N7-C5	2.39	107.54	102.99
2	D	102	3AM	C4-C5-N7	-2.36	106.94	109.40
	1	l	1	1	1	Continued on n	ert nage



 $Continued\ from\ previous\ page...$

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
3	В	103	3GP	O6-C6-C5	-2.36	119.77	124.37
3	F	103	3GP	C8-N7-C5	2.35	107.47	102.99
2	D	101	3AM	C4-C5-N7	-2.32	106.98	109.40
2	A	102	3AM	C1'-N9-C4	-2.31	122.58	126.64
3	Е	103	3GP	C8-N7-C5	2.31	107.39	102.99
3	D	103	3GP	C5-C6-N1	2.30	118.02	113.95
3	A	103	3GP	C8-N7-C5	2.27	107.32	102.99
3	D	103	3GP	C8-N7-C5	2.26	107.30	102.99
2	F	101	3AM	C4-C5-N7	-2.26	107.05	109.40
3	В	103	3GP	C8-N7-C5	2.24	107.26	102.99
2	Е	101	3AM	C4-C5-N7	-2.24	107.07	109.40
3	A	103	3GP	O3'-C3'-C2'	2.19	119.52	111.53
2	F	101	3AM	C2-N1-C6	2.18	122.49	118.75
3	Е	103	3GP	C5-C6-N1	2.15	117.75	113.95
3	D	103	3GP	C2'-C3'-C4'	2.14	107.01	103.22
2	Е	101	3AM	C2-N1-C6	2.13	122.40	118.75
3	F	103	3GP	C5-C6-N1	2.13	117.72	113.95
2	В	102	3AM	C4-C5-N7	-2.11	107.20	109.40
2	F	102	3AM	C2-N1-C6	2.08	122.32	118.75
2	Е	102	3AM	C2-N1-C6	2.08	122.31	118.75

There are no chirality outliers.

All (18) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	103	3GP	C4'-C3'-O3'-P
3	A	103	3GP	C2'-C3'-O3'-P
3	В	103	3GP	C4'-C3'-O3'-P
3	В	103	3GP	C2'-C3'-O3'-P
3	Е	103	3GP	C4'-C3'-O3'-P
3	Е	103	3GP	C2'-C3'-O3'-P
3	F	103	3GP	C4'-C3'-O3'-P
3	F	103	3GP	C2'-C3'-O3'-P
2	A	101	3AM	C3'-C4'-C5'-O5'
2	A	101	3AM	O4'-C4'-C5'-O5'
2	Е	101	3AM	O4'-C4'-C5'-O5'
2	F	101	3AM	O4'-C4'-C5'-O5'
2	Е	101	3AM	C3'-C4'-C5'-O5'
2	D	101	3AM	O4'-C4'-C5'-O5'
2	В	101	3AM	C3'-C4'-C5'-O5'
2	F	101	3AM	C3'-C4'-C5'-O5'
2	В	101	3AM	O4'-C4'-C5'-O5'



Continued from previous page...

\mathbf{Mol}	Chain	Res	Type	Atoms
2	D	101	3AM	C3'-C4'-C5'-O5'


There are no ring outliers.

4 monomers are involved in 6 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	В	101	3AM	2	0
2	A	101	3AM	1	0
3	В	103	3GP	5	0
3	A	103	3GP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





















































































5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	$\langle { m RSRZ} \rangle$	# RSRZ > 2	$OWAB(A^2)$	Q<0.9
1	A	92/92 (100%)	0.42	3 (3%) 46 53	16, 30, 56, 84	0
1	В	92/92 (100%)	1.07	15 (16%) 1 2	27, 43, 80, 103	0
1	С	91/92 (98%)	0.57	7 (7%) 13 17	20, 36, 72, 102	0
1	D	91/92 (98%)	0.68	9 (9%) 7 9	21, 37, 72, 93	0
1	E	91/92 (98%)	1.01	13 (14%) 2 3	31, 52, 93, 107	0
1	F	91/92 (98%)	0.98	15 (16%) 1 2	30, 50, 88, 107	0
All	All	548/552 (99%)	0.79	62 (11%) 5 6	16, 42, 83, 107	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	Е	5	HIS	9.8
1	В	2	ASP	7.6
1	Е	4	GLN	7.0
1	В	11	TYR	6.5
1	В	5	HIS	6.4
1	D	11	TYR	6.0
1	Е	11	TYR	5.9
1	F	3	ASN	5.8
1	В	4	GLN	5.4
1	В	6	LYS	5.1
1	В	3	ASN	4.9
1	Е	19	ILE	4.9
1	Е	8	ILE	4.9
1	F	6	LYS	4.8
1	Е	54	ARG	4.6
1	В	7	LYS	4.4
1	D	52	HIS	4.3
1	F	19	ILE	4.3
1	F	7	LYS	4.3



Continued from previous page...

Mol	nued fron Chain	Res	Type	RSRZ
1	В	51	LEU	3.9
1	F	51	LEU	3.6
1	Е	6	LYS	3.6
1	A	2	ASP	3.6
1	D	5	HIS	3.6
1	F	11	TYR	3.5
1	F	4	GLN	3.5
1	С	54	ARG	3.5
1	F	54	ARG	3.5
1	Е	3	ASN	3.4
1	F	5	HIS	3.3
1	A	52	HIS	3.3
1	F	55	ASP	3.2
1	A	51	LEU	3.2
1	C	11	TYR	3.1
1	C C F	3	ASN	3.0
1	С	5	HIS	2.9
1		52	HIS	2.9
1	В	18	GLU	2.8
1	F	57	ILE	2.7
1	D	54	ARG	2.7
1	Е	52	HIS	2.7
1	В	52	HIS	2.7
1	С	4	GLN	2.7
1	В	76	ILE	2.7
1	Е	7	LYS	2.6
1	F	17	GLU	2.5
1	С	55	ASP	2.5
1	D	83	ILE	2.5
1	Е	20	ASP	2.4
1	В	14	LEU	2.4
1	F	8	ILE	2.3
1	D	16	GLN	2.2
1	В	83	ILE	2.2
1	E	57	ILE	2.2
1	F	49	ALA	2.2
1	D	3	ASN	2.2
1	В	16	GLN	2.1
1	D	82	ALA	2.1
1	Е	16	GLN	2.1
1	С	6	LYS	2.1
1	D	81	MET	2.0



Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	В	57	ILE	2.0

Non-standard residues in protein, DNA, RNA chains (i) 6.2

There are no non-standard protein/DNA/RNA residues in this entry.

Carbohydrates (i) 6.3

There are no monosaccharides in this entry.


Ligands (i) 6.4

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B\text{-}factors}({f \AA}^2)$	Q<0.9
2	3AM	A	101	22/23	0.96	0.14	12,21,26,32	22
2	3AM	A	102	22/23	0.96	0.13	13,20,26,32	22
2	3AM	В	101	22/23	0.96	0.11	18,33,37,39	22
2	3AM	В	102	22/23	0.96	0.12	18,31,37,39	22
2	3AM	D	102	22/23	0.96	0.11	20,25,32,33	22
2	3AM	D	101	22/23	0.97	0.11	20,24,32,33	22
2	3AM	E	101	22/23	0.97	0.12	32,40,46,49	22
2	3AM	Е	102	22/23	0.97	0.12	33,41,46,49	22
2	3AM	F	101	22/23	0.97	0.11	32,37,41,47	22
2	3AM	F	102	22/23	0.97	0.13	31,36,41,48	22
3	3GP	A	103	23/24	0.97	0.14	16,22,28,32	23
3	3GP	В	103	23/24	0.97	0.13	25,33,37,41	23
3	3GP	D	103	23/24	0.97	0.12	20,25,33,33	23
3	3GP	Е	103	23/24	0.97	0.14	33,41,46,49	23
3	3GP	F	103	23/24	0.97	0.13	31,36,41,48	23
2	3AM	С	101	22/23	0.98	0.10	21,27,33,36	22
2	3AM	С	102	22/23	0.98	0.10	22,27,33,37	22
3	3GP	С	103	23/24	0.98	0.11	21,27,33,38	23

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.





































Electron density around 3AM F 101:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)































Electron density around 3GP E 103:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 3GP F 103:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around 3AM C 101:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around 3AM C 102:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

