



Full wwPDB X-ray Structure Validation Report ⓘ

May 17, 2020 – 03:41 am BST

PDB ID : 2J0R
Title : Structure of the haem-chaperone Proteobacteria-protein HemS
Authors : Schneider, S.; Sharp, K.H.; Barker, P.D.; Paoli, M.
Deposited on : 2006-08-04
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	345	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	PEG	A	1350	-	-	-	X
4	PGE	A	1349	-	-	-	X

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 2901 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called HEMIN TRANSPORT PROTEIN HEMS.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	331	2681	1683	483	508	7	13	5	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	162	ASN	ILE	conflict	UNP P31517
A	334	GLN	GLU	conflict	UNP P31517

- Molecule 2 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: C₂H₆O₂).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		
2	A	1	Total	C	O	0	0
			4	2	2		

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C₄H₁₀O₃).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		
3	A	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: C₆H₁₄O₄).




Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	C	O	0	0
			10	6	4		

- Molecule 5 is PENTAETHYLENE GLYCOL (three-letter code: 1PE) (formula: $C_{10}H_{22}O_6$).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
5	A	1	Total	C	O	0	0
			16	10	6		

- Molecule 6 is DODECAETHYLENE GLYCOL (three-letter code: 12P) (formula: $C_{24}H_{50}O_{13}$).



Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
6	A	1	Total	C O	0	0
			19	12 7		


- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	134	Total	O	0	0
			134	134		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: HEMIN TRANSPORT PROTEIN HEMS



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	62.02Å 68.27Å 73.62Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	50.06 – 1.90 29.90 – 1.90	Depositor EDS
% Data completeness (in resolution range)	99.9 (50.06-1.90) 100.0 (29.90-1.90)	Depositor EDS
R_{merge}	0.09	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.23 (at 1.91Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, R_{free}	0.189 , 0.233 0.201 , 0.243	Depositor DCC
R_{free} test set	1289 reflections (5.10%)	wwPDB-VP
Wilson B-factor (Å ²)	20.6	Xtrriage
Anisotropy	0.102	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 51.6	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.94	EDS
Total number of atoms	2901	wwPDB-VP
Average B, all atoms (Å ²)	23.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.60% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: 12P, PGE, PEG, EDO, 1PE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z > 5$	RMSZ	# $ Z > 5$
1	A	1.11	5/2742 (0.2%)	0.80	6/3715 (0.2%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (5) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	45	LYS	CE-NZ	-6.55	1.32	1.49
1	A	168	GLU	CG-CD	-6.29	1.42	1.51
1	A	239	GLU	CD-OE2	-5.40	1.19	1.25
1	A	116	GLU	CD-OE2	-5.23	1.19	1.25
1	A	224	VAL	CB-CG1	-5.16	1.42	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	49	GLY	N-CA-C	6.16	128.49	113.10
1	A	88	GLY	N-CA-C	5.71	127.38	113.10
1	A	86	LEU	C-N-CA	5.66	135.86	121.70
1	A	293	ARG	NE-CZ-NH1	5.59	123.09	120.30
1	A	146	ASP	CB-CG-OD1	5.13	122.91	118.30
1	A	23	ASP	CB-CG-OD2	5.01	122.81	118.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	118	THR	Peptide

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2681	0	2620	35	1
2	A	20	0	30	1	0
3	A	21	0	30	3	0
4	A	10	0	14	2	0
5	A	16	0	22	1	0
6	A	19	0	25	3	0
7	A	134	0	0	1	0
All	All	2901	0	2741	36	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (36) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:104:PHE:HB2	3:A:1347:PEG:H41	1.65	0.78
1:A:188:GLU:HG2	1:A:204:ARG:HH22	1.54	0.71
1:A:65:LYS:HZ2	6:A:1352:12P:H241	1.56	0.69
1:A:241:ASN:H	1:A:241:ASN:HD22	1.41	0.69
1:A:65:LYS:HE2	6:A:1352:12P:H351	1.74	0.68
1:A:241:ASN:HD21	1:A:334:GLN:HE22	1.42	0.66
1:A:332:ARG:HH22	3:A:1348:PEG:H21	1.66	0.61
1:A:47:LEU:C	1:A:49:GLY:H	2.08	0.56
1:A:81:TYR:CZ	1:A:99:LEU:HD21	2.41	0.56
1:A:241:ASN:ND2	1:A:334:GLN:HE22	2.04	0.55
1:A:52:ARG:NH2	7:A:2024:HOH:O	2.42	0.53
1:A:131:HIS:HB3	1:A:168:GLU:HG3	1.93	0.51
1:A:65:LYS:HD2	6:A:1352:12P:H242	1.93	0.49
1:A:118:THR:O	1:A:120:HIS:N	2.45	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:148:PRO:HG2	4:A:1349:PGE:H1	1.94	0.48
1:A:118:THR:C	1:A:119:ARG:HG3	2.34	0.48
1:A:136:LEU:HD22	1:A:251:GLY:HA3	1.96	0.48
1:A:317:LEU:N	1:A:317:LEU:HD12	2.29	0.48
1:A:47:LEU:O	1:A:49:GLY:N	2.47	0.47
1:A:316:GLN:C	1:A:317:LEU:HD12	2.35	0.47
1:A:241:ASN:H	1:A:241:ASN:ND2	2.10	0.46
3:A:1347:PEG:H22	3:A:1347:PEG:H42	1.57	0.46
1:A:330:GLN:O	1:A:334:GLN:HG3	2.15	0.46
1:A:170:LEU:HA	1:A:171:SER:C	2.35	0.45
1:A:96:PRO:O	1:A:97:ARG:HB2	2.17	0.45
1:A:189:TRP:CE3	1:A:216:VAL:HG11	2.53	0.44
1:A:210:GLN:HB3	2:A:1343:EDO:H11	2.01	0.43
1:A:47:LEU:C	1:A:49:GLY:N	2.73	0.42
1:A:122:VAL:HG13	1:A:122:VAL:O	2.18	0.42
1:A:19:LYS:NZ	1:A:27:LEU:HD11	2.34	0.41
1:A:318:TYR:CE1	5:A:1351:1PE:H131	2.56	0.41
1:A:200:GLN:HA	1:A:203:LYS:HG2	2.02	0.41
1:A:208:THR:O	1:A:209:ARG:C	2.56	0.41
1:A:147:MET:HG2	4:A:1349:PGE:H62	2.03	0.41
1:A:9:TYR:CD1	1:A:35:LEU:HD11	2.56	0.41
1:A:49:GLY:O	1:A:50:ASP:C	2.58	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:143:GLU:OE2	1:A:211:GLN:NE2[2_444]	1.28	0.92

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	330/345 (96%)	317 (96%)	11 (3%)	2 (1%)	25 15

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	48	LYS
1	A	49	GLY

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	284/292 (97%)	267 (94%)	17 (6%)	19 9

All (17) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	10	LEU
1	A	71	THR
1	A	77	GLN
1	A	104	PHE
1	A	117	GLU
1	A	143	GLU
1	A	168	GLU
1	A	171	SER
1	A	188	GLU
1	A	193	THR
1	A	207	LEU
1	A	237	GLN
1	A	241	ASN
1	A	267	GLN
1	A	293	ARG
1	A	295	PRO
1	A	320	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	84	GLN
1	A	127	GLN
1	A	226	ASN
1	A	237	GLN
1	A	241	ASN
1	A	320	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
2	EDO	A	1346	-	3,3,3	0.49	0	2,2,2	0.31	0
2	EDO	A	1342	-	3,3,3	0.44	0	2,2,2	0.37	0
2	EDO	A	1345	-	3,3,3	0.50	0	2,2,2	0.30	0
2	EDO	A	1344	-	3,3,3	0.36	0	2,2,2	0.48	0
3	PEG	A	1348	-	6,6,6	0.44	0	5,5,5	0.33	0
5	1PE	A	1351	-	15,15,15	0.49	0	14,14,14	0.27	0
4	PGE	A	1349	-	9,9,9	0.61	0	8,8,8	0.44	0
3	PEG	A	1350	-	6,6,6	0.38	0	5,5,5	0.21	0
6	12P	A	1352	-	18,18,36	0.48	0	17,17,35	0.32	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	EDO	A	1343	-	3,3,3	0.51	0	2,2,2	0.24	0
3	PEG	A	1347	-	6,6,6	0.51	0	5,5,5	0.49	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	EDO	A	1346	-	-	1/1/1/1	-
2	EDO	A	1342	-	-	1/1/1/1	-
2	EDO	A	1345	-	-	1/1/1/1	-
2	EDO	A	1344	-	-	1/1/1/1	-
3	PEG	A	1348	-	-	2/4/4/4	-
5	1PE	A	1351	-	-	8/13/13/13	-
4	PGE	A	1349	-	-	7/7/7/7	-
3	PEG	A	1350	-	-	1/4/4/4	-
6	12P	A	1352	-	-	9/16/16/34	-
2	EDO	A	1343	-	-	1/1/1/1	-
3	PEG	A	1347	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (34) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	1347	PEG	C4-C3-O2-C2
3	A	1350	PEG	C4-C3-O2-C2
5	A	1351	1PE	OH4-C13-C23-OH3
6	A	1352	12P	O28-C29-C30-O31
6	A	1352	12P	O25-C26-C27-O28
5	A	1351	1PE	OH2-C12-C22-OH3
4	A	1349	PGE	O3-C5-C6-O4
5	A	1351	1PE	OH6-C15-C25-OH5
4	A	1349	PGE	O1-C1-C2-O2
3	A	1348	PEG	O2-C3-C4-O4
4	A	1349	PGE	O2-C3-C4-O3
5	A	1351	1PE	OH7-C16-C26-OH6
2	A	1342	EDO	O1-C1-C2-O2

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms
2	A	1345	EDO	O1-C1-C2-O2
6	A	1352	12P	O31-C32-C33-O34
2	A	1344	EDO	O1-C1-C2-O2
6	A	1352	12P	C23-C24-O25-C26
4	A	1349	PGE	C3-C4-O3-C5
5	A	1351	1PE	C25-C15-OH6-C26
5	A	1351	1PE	C24-C14-OH5-C25
3	A	1347	PEG	C1-C2-O2-C3
2	A	1346	EDO	O1-C1-C2-O2
4	A	1349	PGE	C6-C5-O3-C4
6	A	1352	12P	C33-C32-O31-C30
6	A	1352	12P	C27-C26-O25-C24
2	A	1343	EDO	O1-C1-C2-O2
6	A	1352	12P	C36-C35-O34-C33
5	A	1351	1PE	C13-C23-OH3-C22
6	A	1352	12P	C24-C23-O22-C21
3	A	1348	PEG	O1-C1-C2-O2
5	A	1351	1PE	C23-C13-OH4-C24
4	A	1349	PGE	C4-C3-O2-C2
4	A	1349	PGE	C1-C2-O2-C3
6	A	1352	12P	O22-C23-C24-O25


There are no ring outliers.

6 monomers are involved in 10 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	A	1348	PEG	1	0
5	A	1351	1PE	1	0
4	A	1349	PGE	2	0
6	A	1352	12P	3	0
2	A	1343	EDO	1	0
3	A	1347	PEG	2	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and

any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	331/345 (95%)	0.71	24 (7%) 15 16	10, 20, 45, 60	3 (0%)

All (24) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	49	GLY	10.3
1	A	118	THR	7.2
1	A	120	HIS	6.2
1	A	199	PHE	5.8
1	A	119	ARG	5.1
1	A	340	ASN	3.9
1	A	196	HIS	3.1
1	A	204	ARG	3.0
1	A	297	LYS	2.9
1	A	341	LYS	2.9
1	A	195	VAL	2.8
1	A	171	SER	2.7
1	A	122	VAL	2.6
1	A	198	PHE	2.5
1	A	185	VAL	2.5
1	A	170	LEU	2.3
1	A	216	VAL	2.3
1	A	11	GLN	2.3
1	A	203	LYS	2.3
1	A	180	ALA	2.3
1	A	202	LEU	2.1
1	A	333	ASP	2.1
1	A	193	THR	2.1
1	A	339	ASN	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)


There are no carbohydrates in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	PGE	A	1349	10/10	0.67	0.46	30,41,44,44	0
3	PEG	A	1348	7/7	0.69	0.37	59,59,60,61	0
2	EDO	A	1342	4/4	0.72	0.23	41,42,43,44	0
3	PEG	A	1347	7/7	0.73	0.23	35,37,40,42	0
3	PEG	A	1350	7/7	0.76	0.41	30,33,39,42	0
5	1PE	A	1351	16/16	0.79	0.20	48,49,52,52	0
6	12P	A	1352	19/37	0.80	0.25	37,45,55,55	0
2	EDO	A	1345	4/4	0.81	0.13	43,43,44,44	0
2	EDO	A	1343	4/4	0.83	0.42	40,40,40,41	0
2	EDO	A	1346	4/4	0.84	0.10	44,44,44,45	0
2	EDO	A	1344	4/4	0.86	0.17	34,37,39,41	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



6.5 Other polymers [i](#)

There are no such residues in this entry.