



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 27, 2024 – 11:18 pm BST

PDB ID : 2J1D  
Title : Crystallization of hDaam1 C-terminal Fragment  
Authors : Lu, J.; Meng, W.; Poy, F.; Eck, M.J.  
Deposited on : 2006-08-10  
Resolution : 2.25 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.4, CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.25 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1377 (2.26-2.26)
Clashscore	141614	1487 (2.26-2.26)
Ramachandran outliers	138981	1449 (2.26-2.26)
Sidechain outliers	138945	1450 (2.26-2.26)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	G	483	 66% 13% • 18%

## 2 Entry composition [i](#)


There are 4 unique types of molecules in this entry. The entry contains 3288 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called DISHEVELED-ASSOCIATED ACTIVATOR OF MORPHOGENESIS 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	G	396	3202	2024	551	617	10	0	0	1

- Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O<sub>4</sub>P).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
			Total	O	P		
2	G	1	5	4	1	0	0

- Molecule 3 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	G	1	Total C O 6 3 3	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	G	75	Total O 75 75	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: DISHEVELED-ASSOCIATED ACTIVATOR OF MORPHOGENESIS 1

Chain G:  66% 13% 18%



## 4 Data and refinement statistics

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	100.90Å 100.34Å 148.40Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.83 – 2.25 44.42 – 2.55	Depositor EDS
% Data completeness (in resolution range)	100.0 (19.83-2.25) 98.4 (44.42-2.55)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.96 (at 2.54Å)	Xtrriage
Refinement program	REFMAC 5.2.0019	Depositor
R, $R_{free}$	0.218 , 0.250 0.262 , 0.267	Depositor DCC
$R_{free}$ test set	1294 reflections (5.27%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	51.5	Xtrriage
Anisotropy	0.190	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.36 , 75.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	0.000 for -1/2*h-1/2*k-1/2*l,-1/2*h-1/2*k+1/2*l,-h+k 0.000 for -1/2*h-1/2*k+1/2*l,-1/2*h-1/2*k-1/2*l,h-k 0.000 for -1/2*h+1/2*k+1/2*l,1/2*h-1/2*k+1/2*l,h+k 0.015 for -1/2*h+1/2*k-1/2*l,1/2*h-1/2*k-1/2*l,-h-k 0.000 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.92	EDS
Total number of atoms	3288	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	68.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 12.82% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, PO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	G	0.47	0/3250	0.67	4/4358 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	G	0	1

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	G	602	SER	N-CA-C	7.99	132.56	111.00
1	G	874	LEU	CA-CB-CG	6.68	130.66	115.30
1	G	602	SER	CB-CA-C	-6.25	98.23	110.10
1	G	759	ARG	NE-CZ-NH2	-5.15	117.72	120.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	G	830	GLY	Peptide

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	3202	0	3225	50	0
2	G	5	0	0	1	0
3	G	6	0	8	0	0
4	G	75	0	0	5	1
All	All	3288	0	3233	50	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 8.

All (50) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:752:ASP:OD2	4:G:2025:HOH:O	1.55	1.21
1:G:644:ASP:O	1:G:648:THR:HG23	1.65	0.95
1:G:644:ASP:O	1:G:648:THR:CG2	2.20	0.89
1:G:751:LEU:HA	1:G:754:MET:HE2	1.55	0.88
1:G:894:LYS:O	1:G:898:THR:HG23	1.90	0.71
1:G:923:LYS:HE3	4:G:2068:HOH:O	1.91	0.70
1:G:924:PHE:CE2	1:G:928:VAL:HG21	2.28	0.69
1:G:625:GLY:O	4:G:2005:HOH:O	2.11	0.68
1:G:741:ILE:HD11	4:G:2022:HOH:O	1.94	0.66
1:G:804:VAL:HG13	1:G:960:PHE:HE2	1.62	0.65
1:G:804:VAL:HG13	1:G:960:PHE:CE2	2.33	0.63
1:G:687:VAL:CG1	1:G:743:LEU:HD21	2.30	0.61
1:G:797:ILE:HD11	1:G:887:VAL:HG11	1.81	0.61
1:G:687:VAL:HG11	1:G:743:LEU:HD11	1.84	0.60
1:G:910:LEU:HD22	1:G:928:VAL:HG12	1.84	0.59
1:G:829:ARG:NE	1:G:829:ARG:HA	2.17	0.58
1:G:828:GLN:O	1:G:829:ARG:HB2	2.04	0.57
1:G:924:PHE:CZ	1:G:928:VAL:HG21	2.40	0.57
1:G:798:ARG:NH2	1:G:802:GLU:OE2	2.39	0.55
1:G:644:ASP:O	1:G:648:THR:HG22	2.06	0.55
1:G:957:VAL:HG13	1:G:962:GLU:HB2	1.88	0.55
1:G:602:SER:O	1:G:602:SER:OG	2.15	0.54
1:G:687:VAL:HG12	1:G:743:LEU:HD21	1.88	0.54
1:G:864:VAL:HG22	1:G:868:TYR:HB2	1.91	0.52
1:G:699:LEU:HD13	1:G:730:GLN:HB3	1.93	0.50
1:G:754:MET:HE3	1:G:759:ARG:HA	1.93	0.50
1:G:902:GLY:O	1:G:906:VAL:HG23	2.12	0.50

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:G:743:LEU:O	1:G:746:GLU:HB2	2.11	0.49
1:G:636:VAL:HG12	1:G:640:LEU:HD22	1.93	0.49
1:G:843:ILE:HG23	1:G:856:LEU:HD13	1.95	0.48
1:G:751:LEU:CA	1:G:754:MET:HE2	2.33	0.47
1:G:858:HIS:ND1	1:G:990:ASN:ND2	2.63	0.47
1:G:910:LEU:HD13	1:G:928:VAL:HB	1.96	0.46
1:G:687:VAL:HG11	1:G:743:LEU:CD1	2.46	0.46
1:G:751:LEU:HD22	1:G:754:MET:CE	2.45	0.46
1:G:782:LYS:NZ	2:G:3026:PO4:O1	2.33	0.46
1:G:848:SER:HB3	1:G:851:ASP:O	2.16	0.46
1:G:683:LYS:NZ	1:G:753:ARG:HH22	2.14	0.45
1:G:683:LYS:NZ	1:G:753:ARG:NH2	2.64	0.45
1:G:683:LYS:HZ1	1:G:753:ARG:HH22	1.65	0.45
1:G:636:VAL:HG13	1:G:640:LEU:HD13	2.00	0.44
1:G:906:VAL:HG12	1:G:932:ILE:HD11	2.00	0.43
1:G:623:LEU:O	1:G:624:GLU:HB3	2.19	0.43
1:G:687:VAL:CG1	1:G:743:LEU:HD11	2.47	0.43
1:G:603:ILE:HD12	1:G:604:PRO:O	2.18	0.43
1:G:925:VAL:HG23	4:G:2067:HOH:O	2.20	0.42
1:G:828:GLN:OE1	1:G:829:ARG:N	2.43	0.41
1:G:751:LEU:HD22	1:G:754:MET:HE1	2.03	0.40
1:G:855:THR:H	1:G:858:HIS:CD2	2.39	0.40
1:G:858:HIS:HB3	1:G:990:ASN:HD21	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:G:2054:HOH:O	4:G:2068:HOH:O[6_555]	1.65	0.55

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	G	392/483 (81%)	383 (98%)	9 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	G	352/438 (80%)	318 (90%)	34 (10%)	8	6

All (34) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	610	LEU
1	G	615	TRP
1	G	624	GLU
1	G	627	VAL
1	G	635	LYS
1	G	645	LEU
1	G	648	THR
1	G	683	LYS
1	G	685	LEU
1	G	686	SER
1	G	703	LEU
1	G	741	ILE
1	G	745	GLU
1	G	759	ARG
1	G	772	GLN
1	G	790	VAL
1	G	795	GLU
1	G	798	ARG
1	G	804	VAL
1	G	806	ARG
1	G	828	GLN
1	G	856	LEU
1	G	857	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	G	860	LEU
1	G	863	ILE
1	G	874	LEU
1	G	892	LEU
1	G	898	THR
1	G	899	LEU
1	G	930	GLN
1	G	934	VAL
1	G	951	ASP
1	G	970	ASP
1	G	1001	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	G	697	ASN
1	G	883	GLN
1	G	888	ASN
1	G	919	GLN
1	G	990	ASN
1	G	992	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	PO4	G	3026	-	4,4,4	0.84	0	6,6,6	0.77	0
3	GOL	G	3027	-	5,5,5	0.42	0	5,5,5	0.63	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	GOL	G	3027	-	-	2/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	G	3027	GOL	O1-C1-C2-C3
3	G	3027	GOL	O1-C1-C2-O2

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	G	3026	PO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.2 Non-standard residues in protein, DNA, RNA chains

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.3 Carbohydrates

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.4 Ligands

Unable to reproduce the depositors R factor - this section is therefore empty.

### 6.5 Other polymers

Unable to reproduce the depositors R factor - this section is therefore empty.