

wwPDB X-ray Structure Validation Summary Report (i)

Sep 10, 2023 – 03:48 PM EDT

PDB ID : 4J1R

Title: Crystal Structure of GSK3b in complex with inhibitor 15R

Authors: Zhan, C.; Wang, Y.; Wach, J.; Sheehan, P.; Zhong, C.; Harris, R.; Patskovsky,

Y.; Bishop, J.; Haggarty, S.; Ramek, A.; Berry, K.; O'Herin, C.; Koehler, A.N.; Hung, A.W.; Young, D.W.; Almo, S.C.; New York Structural Genomics

Research Consortium (NYSGRC)

Deposited on : 2013-02-01

Resolution : 2.70 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.13

EDS : 2.35.1buster-report : 1.1.7 (2018)

Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)

Refmac : 5.8.0158

CCP4 : 7.0.044 (Gargrove)

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.35.1




1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}(\mathring{\rm A})) \end{array}$
R_{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	424	66%	13%		19%
1	В	424	67%	14%	•	18%
1	С	424	67%	13%	·	19%
1	D	424	69%	12%		18%



2 Entry composition (i)

There are 5 unique types of molecules in this entry. The entry contains 11248 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Glycogen synthase kinase-3 beta.


Mol	Chain	Residues		Atoms				ZeroOcc	AltConf	Trace	
1	A	344	Total	С	N	О	Р	S	0	0	0
1	A	344	2753	1770	469	502	1	11	0		0
1	В	347	Total	С	N	О	Р	S	0	0	0
1	Б		2770	1780	475	503	1	11	0	0	
1	С	345	Total	С	N	О	Р	S	0	0	0
1		340	2754	1770	472	500	1	11	0	U	
1	D	246	Total	С	N	О	Р	S	0	0	0
1		346	2768	1780	474	502	1	11	U	U	U

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-3	GLY	-	expression tag	UNP P49841
A	-2	SER	-	expression tag	UNP P49841
A	-1	PRO	-	expression tag	UNP P49841
A	0	GLY	-	expression tag	UNP P49841
В	-3	GLY	-	expression tag	UNP P49841
В	-2	SER	-	expression tag	UNP P49841
В	-1	PRO	_	expression tag	UNP P49841
В	0	GLY	-	expression tag	UNP P49841
С	-3	GLY	-	expression tag	UNP P49841
С	-2	SER	-	expression tag	UNP P49841
С	-1	PRO	-	expression tag	UNP P49841
С	0	GLY	-	expression tag	UNP P49841
D	-3	GLY	-	expression tag	UNP P49841
D	-2	SER	-	expression tag	UNP P49841
D	-1	PRO	-	expression tag	UNP P49841
D	0	GLY	_	expression tag	UNP P49841


• Molecule 2 is PHOSPHATE ION (three-letter code: PO4) (formula: O₄P).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O P	0	0
		_	5 4 1		<u> </u>
2	В	1	Total O P	0	0
2	Ъ	1	5 4 1	U	U
2	C	1	Total O P	al O P	0
		1	5 4 1	0	U
9	D	1	Total O P	0	0
	ע	1	5 4 1	0	U

• Molecule 3 is (2R)-2-(1H-indol-3-ylmethyl)-1,4-dihydropyrido[2,3-b]pyrazin-3(2H)-one (three-letter code: I5R) (formula: $C_{16}H_{14}N_4O$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	
3	Δ	1	Total	С	N	О	0	0	
	11	1	21	16	4	1		U	
3	В	1	1 Total	\mathbf{C}	N	Ο	0	0	
	D	1	21	16	4	1		U	
3	C	1	Total	С	Ν	O	0	0	
J	O	1	21	16	4	1	U	0	
3	D	1	Total	С	N	О	0	0	
	ט	1	21	16	4	1			

• Molecule 4 is POTASSIUM ION (three-letter code: K) (formula: K).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	
1	Λ	1	1 Total K		0	
4	Λ	1	1 1		0	
1	B	1	Total K	0	0	
4	4 D	1	1 1	0	U	
1	С	1	Total K	0	0	
4		1	1 1	0	U	
4	D	1	Total K	0	0	
4	ט	1	1 1	0	U	

• Molecule 5 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	20	Total O 20 20	0	0
5	В	27	Total O 27 27	0	0
5	С	23	Total O 23 23	0	0
5	D	25	Total O 25 25	0	0




3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


• Molecule 1: Glycogen synthase kinase-3 beta







• Molecule 1: Glycogen synthase kinase-3 beta





4 Data and refinement statistics (i)

Property	Value	Source	
Space group	P 1	Depositor	
Cell constants	67.52Å 67.44Å 116.42Å	Donositon	
a, b, c, α , β , γ	90.13° 90.08° 81.11°	Depositor	
Resolution (Å)	46.96 - 2.70	Depositor	
Resolution (A)	46.96 - 2.70	EDS	
% Data completeness	97.2 (46.96-2.70)	Depositor	
(in resolution range)	96.8 (46.96-2.70)	EDS	
R_{merge}	0.10	Depositor	
R_{sym}	(Not available)	Depositor	
$< I/\sigma(I) > 1$	1.64 (at 2.69Å)	Xtriage	
Refinement program	PHENIX 1.7.3_928	Depositor	
P.P.	0.173 , 0.193	Depositor	
R, R_{free}	0.170 , 0.186	DCC	
R_{free} test set	2756 reflections (5.10%)	wwPDB-VP	
Wilson B-factor (Å ²)	53.4	Xtriage	
Anisotropy	0.318	Xtriage	
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	$0.33 \; , 34.7$	EDS	
L-test for twinning ²	$< L > = 0.45, < L^2> = 0.28$	Xtriage	
	0.440 for -h,-k,l		
Estimated twinning fraction	0.438 for k,h,-l	Xtriage	
	0.428 for -k,-h,-l		
Reported twinning fraction	0.469 for k,h,-l	Depositor	
Outliers	0 of 54014 reflections	Xtriage	
F_o, F_c correlation	0.95	EDS	
Total number of atoms	11248	wwPDB-VP	
Average B, all atoms (\mathring{A}^2)	64.0	wwPDB-VP	

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 6.22% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: K, PO4, PTR, I5R

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bo	nd lengths	Bond angles		
Moi Chain		RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.54	1/2805~(0.0%)	0.59	0/3818	
1	В	0.57	$2/2821 \ (0.1\%)$	0.61	$2/3837 \ (0.1\%)$	
1	С	0.52	0/2805	0.58	1/3817 (0.0%)	
1	D	0.55	0/2820	0.61	0/3838	
All	All	0.55	3/11251 (0.0%)	0.60	3/15310 (0.0%)	

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(A)	$\operatorname{Ideal}(\text{\AA})$
1	A	324	THR	CB-CG2	-5.99	1.32	1.52
1	В	218	CYS	CB-SG	-5.24	1.73	1.81
1	В	43	THR	CB-OG1	-5.23	1.32	1.43

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	В	180	ARG	NE-CZ-NH1	5.98	123.29	120.30
1	С	180	ARG	NE-CZ-NH1	5.17	122.89	120.30
1	В	180	ARG	NE-CZ-NH2	-5.13	117.73	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.



Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2753	0	2757	36	0
1	В	2770	0	2784	41	0
1	С	2754	0	2767	36	0
1	D	2768	0	2777	33	0
2	A	5	0	0	1	0
2	В	5	0	0	1	0
2	С	5	0	0	1	0
2	D	5	0	0	1	0
3	A	21	0	14	0	0
3	В	21	0	14	0	0
3	С	21	0	14	1	0
3	D	21	0	14	2	0
4	A	1	0	0	0	0
4	В	1	0	0	0	0
4	С	1	0	0	0	0
4	D	1	0	0	0	0
5	A	20	0	0	0	0
5	В	27	0	0	1	0
5	С	23	0	0	4	0
5	D	25	0	0	2	0
All	All	11248	0	11141	144	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

The worst 5 of 144 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$egin{aligned} & ext{Interatomic} \ & ext{distance} \ & ext{(Å)} \end{aligned}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:C:297:LYS:HD2	1:C:298:ALA:N	1.46	1.27
1:C:260:ASP:OD2	1:D:144:ARG:NH1	1.84	1.09
1:C:297:LYS:CD	1:C:298:ALA:H	1.64	1.09
1:B:141:ARG:HH21	1:B:141:ARG:HG2	1.16	1.06
1:C:297:LYS:HD2	1:C:298:ALA:H	0.85	0.99

There are no symmetry-related clashes.



5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	339/424~(80%)	325 (96%)	14 (4%)	0	100	100
1	В	342/424~(81%)	328 (96%)	13 (4%)	1 (0%)	41	66
1	C	340/424~(80%)	327 (96%)	13 (4%)	0	100	100
1	D	341/424~(80%)	327 (96%)	14 (4%)	0	100	100
All	All	$1362/1696\ (80\%)$	1307 (96%)	54 (4%)	1 (0%)	51	78

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	В	285	ASN

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	304/365 (83%)	288 (95%)	16 (5%)	22	48	
1	В	305/365~(84%)	292 (96%)	13 (4%)	29	57	
1	С	303/365 (83%)	289 (95%)	14 (5%)	27	54	
1	D	305/365 (84%)	292 (96%)	13 (4%)	29	57	
All	All	1217/1460 (83%)	1161 (95%)	56 (5%)	27	54	

5 of 56 residues with a non-rotameric sidechain are listed below:



Mol	Chain	Res	Type
1	В	369	SER
1	D	329	LEU
1	С	233	ASP
1	D	306	ARG
1	D	233	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	В	129	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Trme	Chain	Res	Link	Bo	ond leng	ths	Bond angles		
MIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	PTR	В	216	1	15,16,17	1.11	1 (6%)	19,22,24	1.13	1 (5%)
1	PTR	С	216	1	15,16,17	0.93	1 (6%)	19,22,24	0.86	1 (5%)
1	PTR	D	216	1	15,16,17	0.97	1 (6%)	19,22,24	0.98	0
1	PTR	A	216	1	15,16,17	1.06	0	19,22,24	1.25	3 (15%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	В	216	1	-	5/10/11/13	0/1/1/1

Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	PTR	С	216	1	-	1/10/11/13	0/1/1/1
1	PTR	D	216	1	-	1/10/11/13	0/1/1/1
1	PTR	A	216	1	-	0/10/11/13	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(A)
1	С	216	PTR	OH-CZ	-2.92	1.34	1.40
1	D	216	PTR	OH-CZ	-2.92	1.34	1.40
1	В	216	PTR	P-O1P	2.09	1.57	1.50

All (5) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
1	A	216	PTR	CB-CA-C	-2.43	106.92	111.47
1	A	216	PTR	CE2-CZ-CE1	2.31	123.74	120.18
1	В	216	PTR	CE2-CZ-CE1	2.28	123.69	120.18
1	С	216	PTR	O3P-P-OH	2.16	111.99	105.24
1	A	216	PTR	O2P-P-OH	2.16	111.99	105.24

There are no chirality outliers.

5 of 7 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
1	В	216	PTR	O-C-CA-CB
1	В	216	PTR	C-CA-CB-CG
1	В	216	PTR	CZ-OH-P-O1P
1	С	216	PTR	O-C-CA-CB
1	D	216	PTR	O-C-CA-CB

There are no ring outliers.

No monomer is involved in short contacts.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

Of 12 ligands modelled in this entry, 4 are monoatomic - leaving 8 for Mogul analysis.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	В	ond leng	$_{ m gths}$	В	ond ang	gles
MIOI	Type	Chain	nes	Lilik	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	PO4	D	501	-	4,4,4	0.81	0	6,6,6	0.71	0
2	PO4	A	501	-	4,4,4	1.08	0	6,6,6	0.27	0
3	I5R	A	502	_	21,24,24	2.59	6 (28%)	29,34,34	3.56	11 (37%)
2	PO4	С	501	-	4,4,4	0.79	0	6,6,6	0.95	0
3	I5R	D	502	-	21,24,24	2.68	8 (38%)	29,34,34	3.57	13 (44%)
3	I5R	В	502	-	21,24,24	2.67	8 (38%)	29,34,34	3.61	14 (48%)
3	I5R	С	502	-	21,24,24	3.31	10 (47%)	29,34,34	3.74	15 (51%)
2	PO4	В	501	-	4,4,4	0.86	0	6,6,6	0.52	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

I	Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
	3	I5R	В	502	-	-	0/3/16/16	0/4/4/4
	3	I5R	A	502	-	-	0/3/16/16	0/4/4/4
	3	I5R	С	502	-	-	0/3/16/16	0/4/4/4
	3	I5R	D	502	-	-	0/3/16/16	0/4/4/4

The worst 5 of 32 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\operatorname{Observed}(\operatorname{\AA})$	Ideal(Å)
3	С	502	I5R	CAR-NAK	-8.20	1.22	1.35
3	A	502	I5R	CAR-NAK	-7.38	1.23	1.35
3	В	502	I5R	CAR-NAK	-6.83	1.24	1.35
3	D	502	I5R	CAR-NAK	-6.67	1.24	1.35
3	С	502	I5R	O-C	5.43	1.34	1.23

The worst 5 of 53 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	D	502	I5R	CB-CA-N	-10.18	98.24	111.33

Continued on next page...



Continued from previous page...

Mol	Chain	Res	Type	Atoms	${f Z}$	$\mathbf{Observed}(^o)$	$\operatorname{Ideal}({}^{o})$
3	A	502	I5R	CB-CA-N	-9.96	98.52	111.33
3	В	502	I5R	CB-CA-N	-9.93	98.56	111.33
3	С	502	I5R	CB-CA-N	-9.31	99.36	111.33
3	A	502	I5R	NAL-CAR-NAK	8.38	125.88	116.83

There are no chirality outliers.

There are no torsion outliers.


There are no ring outliers.


6 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	501	PO4	1	0
2	A	501	PO4	1	0
2	С	501	PO4	1	0
3	D	502	I5R	2	0
3	С	502	I5R	1	0
2	В	501	PO4	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.














5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2		$OWAB(A^2)$	Q < 0.9	
1	A	343/424 (80%)	0.00	1 (0%)	94	95	25, 59, 99, 128	0
1	В	346/424 (81%)	0.07	3 (0%)	84	85	26, 59, 101, 142	0
1	С	344/424 (81%)	0.02	3 (0%)	84	85	37, 59, 99, 132	0
1	D	345/424 (81%)	0.07	2 (0%)	89	91	38, 59, 100, 137	0
All	All	1378/1696 (81%)	0.04	9 (0%)	87	89	25, 59, 101, 142	0

The worst 5 of 9 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	384	ILE	6.6
1	D	92	ARG	3.8
1	С	104	LEU	2.7
1	В	123	LYS	2.7
1	С	384	ILE	2.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	PTR	A	216	16/17	0.97	0.16	33,48,56,73	0
1	PTR	С	216	16/17	0.97	0.16	20,43,61,66	0
1	PTR	В	216	16/17	0.98	0.14	41,53,67,70	0
1	PTR	D	216	16/17	0.99	0.15	21,44,58,60	0



6.3 Carbohydrates (i)

There are no monosaccharides in this entry.


6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	K	D	503	1/1	0.97	0.19	79,79,79,79	0
3	I5R	A	502	21/21	0.98	0.20	44,59,67,77	0
3	I5R	В	502	21/21	0.98	0.24	43,65,78,87	0
3	I5R	С	502	21/21	0.98	0.21	41,56,64,75	0
3	I5R	D	502	21/21	0.98	0.20	39,62,79,84	0
4	K	В	503	1/1	0.98	0.10	84,84,84,84	0
2	PO4	С	501	5/5	0.98	0.12	39,52,71,88	0
2	PO4	В	501	5/5	0.99	0.14	50,51,62,67	0
4	K	A	503	1/1	0.99	0.14	68,68,68,68	0
2	PO4	A	501	5/5	0.99	0.15	46,59,74,88	0
4	K	С	503	1/1	0.99	0.07	77,77,77,77	0
2	PO4	D	501	5/5	0.99	0.16	41,46,55,85	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









Electron density around I5R B 502: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)









6.5 Other polymers (i)

There are no such residues in this entry.

