



wwPDB NMR Structure Validation Summary Report ⓘ

Feb 14, 2022 – 09:36 AM EST

PDB ID : 1J5N
Title : Solution Structure of the Non-Sequence-Specific HMGB protein NHP6A in complex with SRY DNA
Authors : Masse, J.E.; Wong, B.; Yen, Y.-M.; Allain, F.H.-T.; Johnson, R.C.; Feigon, J.
Deposited on : 2002-05-15

This is a wwPDB NMR Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.26
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.26


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	B	15	33% (yellow) 67% (orange)
2	C	15	13% (yellow) 87% (orange)
3	A	93	• 51% (yellow) 22% (orange) • 23% (cyan)

2 Ensemble composition and analysis

This entry contains 20 models. Model 1 is the overall representative, medoid model (most similar to other models).

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:17-A:88 (72)	0.26	1

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 2 single-model clusters were found.

Cluster number	Models
1	1, 3, 4, 5, 6, 7, 9, 10, 11, 13, 17, 19, 20
2	2, 14, 18
3	12, 15
Single-model clusters	8; 16

3 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 2489 atoms, of which 1119 are hydrogens and 0 are deuteriums.

- Molecule 1 is a DNA chain called 5'-D(*GP*GP*GP*GP*TP*GP*AP*TP*TP*GP*TP*TP*CP*AP*G)-3'.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
1	B	15	484	149	172	58	91	14	0

- Molecule 2 is a DNA chain called 5'-D(*CP*TP*GP*AP*AP*CP*AP*AP*TP*CP*AP*CP*CP*CP*C)-3'.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		P
2	C	15	466	143	169	55	85	14	0

- Molecule 3 is a protein called Nonhistone chromosomal protein 6A.

Mol	Chain	Residues	Atoms					Trace	
			Total	C	H	N	O		S
3	A	93	1539	478	778	138	143	2	0

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: 5'-D(*GP*GP*GP*GP*TP*GP*AP*TP*TP*GP*TP*TP*CP*AP*G)-3'

Chain B:  33% 67%

G101
G102
G103
G104
T105
G106
A107
T108
T109
G110
T111
T112
C113
A114
G115

- Molecule 2: 5'-D(*CP*TP*GP*AP*AP*CP*AP*AP*TP*CP*AP*CP*CP*CP*C)-3'

Chain C:  13% 87%

C116
T117
G118
A119
A120
C121
A122
A123
T124
C125
A126
C127
C128
C129
C130

- Molecule 3: Nonhistone chromosomal protein 6A

Chain A:  51% 22% 23%

H1
N2
T3
F4
H5
E6
P7
K8
R9
R10
T11
T12
B13
K14
K15
K16
D17
P18
N19
A20
F21
K22
K23
A24
L25
S26
A27
Y28
M29
F30
F31
A32
N33
E34
N35
R36
D37
L38
V39
R40
S41
E42
N43
P44
D45
T46
T47
F48
G49
G50
N51
G52
K53
K54
L55
G56
E57
K58
W59
K60

A61
L62
T63
P64
E65
E66
K67
Q68
P69
Y70
E71
A72
K73
A74
Q75
A76
D77
K78
K79
R80
R81
E82
S83
E84
K85
E86
L87
Y88
M89
A90
T91
L92
A93

4.2 Residue scores for the representative (medoid) model from the NMR ensemble

The representative model is number 1. Colouring as in section 4.1 above.


- Molecule 1: 5'-D(*GP*GP*GP*GP*TP*GP*AP*TP*TP*GP*TP*TP*CP*AP*G)-3'

Chain B:  33% 67%

G101
G102
G103
G104
T105
G106
A107
T108
T109
G110
T111
T112
C113
A114
G115


- Molecule 2: 5'-D(*CP*TP*GP*AP*AP*CP*AP*AP*TP*CP*AP*CP*CP*CP*C)-3'

Chain C:  7% 93%



- Molecule 3: Nonhistone chromosomal protein 6A

Chain A:  6% 51% 18% • 23%



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with acceptable covalent geometry*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.1

No chemical shift data was provided.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	B	1.08±0.01	0±0/350 (0.0± 0.1%)	2.24±0.02	35±1/541 (6.5± 0.2%)
2	C	1.09±0.02	0±0/332 (0.0± 0.0%)	2.22±0.02	25±1/508 (5.0± 0.1%)
3	A	0.21±0.00	0±0/604 (0.0± 0.0%)	0.32±0.00	0±0/812 (0.0± 0.0%)
All	All	0.80	2/25720 (0.0%)	1.69	1208/37220 (3.2%)

All unique bond outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)	Models	
								Worst	Total
1	B	109	DT	C5-C7	5.12	1.53	1.50	3	2

5 of 71 unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
2	C	124	DT	O4'-C1'-N1	10.30	115.21	108.00	16	20
1	B	109	DT	O4'-C1'-N1	10.20	115.14	108.00	15	20
1	B	108	DT	O4'-C1'-N1	10.02	115.02	108.00	13	20
1	B	103	DG	O4'-C1'-N9	9.68	114.78	108.00	20	20
1	B	107	DA	O4'-C1'-N9	9.39	114.57	108.00	16	20

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	B	312	172	172	28±4
2	C	297	169	169	38±7
3	A	590	579	579	170±9
All	All	23980	18400	18400	4125

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 97.

5 of 778 unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
3:A:47:THR:O	3:A:51:VAL:HG12	0.93	1.63	19	11
3:A:59:TRP:O	3:A:62:LEU:HD12	0.93	1.64	5	19
2:C:116:DC:H4'	2:C:117:DT:OP1	0.90	1.66	14	3
3:A:23:ARG:NH2	3:A:24:ALA:HB3	0.90	1.80	8	1
2:C:126:DA:H4'	2:C:127:DC:OP1	0.89	1.67	13	6

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
3	A	72/93 (77%)	53±2 (73±3%)	11±2 (15±3%)	8±2 (12±3%)	1 7
All	All	1440/1860 (77%)	1055 (73%)	219 (15%)	166 (12%)	1 7

5 of 17 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
3	A	23	ARG	20
3	A	63	THR	20
3	A	17	ASP	18
3	A	62	LEU	18
3	A	22	LYS	16

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
3	A	61/80 (76%)	41±2 (66±3%)	20±2 (34±3%)	1 11
All	All	1220/1600 (76%)	811 (66%)	409 (34%)	1 11

5 of 48 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
3	A	70	TYR	20
3	A	80	ARG	20
3	A	81	TYR	20
3	A	23	ARG	19
3	A	87	LEU	19

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation

No chemical shift data were provided