



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 1, 2023 – 02:52 PM JST

PDB ID : 5J5N  
Title : Crystal structure of the R39W mutant of Populus trichocarpa glutathione transferase PtGSTU30 in complex with glutathione  
Authors : Yang, Q.; Gu, J.; Yang, M.; Zeng, Q.  
Deposited on : 2016-04-03  
Resolution : 2.63 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.63 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	3797 (2.64-2.60)
Clashscore	141614	4168 (2.64-2.60)
Ramachandran outliers	138981	4093 (2.64-2.60)
Sidechain outliers	138945	4093 (2.64-2.60)
RSRZ outliers	127900	3731 (2.64-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	231	
1	B	231	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 3607 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Glutathione transferase family protein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	218	1782	1163	281	328	10	0	0	0
1	B	218	1782	1163	281	328	10	0	0	0

There are 24 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-10	MET	-	expression tag	UNP B9I0G5
A	-9	HIS	-	expression tag	UNP B9I0G5
A	-8	HIS	-	expression tag	UNP B9I0G5
A	-7	HIS	-	expression tag	UNP B9I0G5
A	-6	HIS	-	expression tag	UNP B9I0G5
A	-5	HIS	-	expression tag	UNP B9I0G5
A	-4	HIS	-	expression tag	UNP B9I0G5
A	-3	GLY	-	expression tag	UNP B9I0G5
A	-2	SER	-	expression tag	UNP B9I0G5
A	-1	GLU	-	expression tag	UNP B9I0G5
A	0	PHE	-	expression tag	UNP B9I0G5
A	39	TRP	ARG	engineered mutation	UNP B9I0G5
B	-10	MET	-	expression tag	UNP B9I0G5
B	-9	HIS	-	expression tag	UNP B9I0G5
B	-8	HIS	-	expression tag	UNP B9I0G5
B	-7	HIS	-	expression tag	UNP B9I0G5
B	-6	HIS	-	expression tag	UNP B9I0G5
B	-5	HIS	-	expression tag	UNP B9I0G5
B	-4	HIS	-	expression tag	UNP B9I0G5
B	-3	GLY	-	expression tag	UNP B9I0G5
B	-2	SER	-	expression tag	UNP B9I0G5
B	-1	GLU	-	expression tag	UNP B9I0G5
B	0	PHE	-	expression tag	UNP B9I0G5
B	39	TRP	ARG	engineered mutation	UNP B9I0G5

- Molecule 2 is GLUTATHIONE (three-letter code: GSH) (formula: C<sub>10</sub>H<sub>17</sub>N<sub>3</sub>O<sub>6</sub>S).



Mol	Chain	Residues	Atoms					ZeroOcc	AltConf
			Total	C	N	O	S		
2	A	1	20	10	3	6	1	0	0
2	B	1	20	10	3	6	1	0	0


- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
			Total	O		
3	A	2	2	2	0	0
3	B	1	1	1	0	0

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Glutathione transferase family protein



- Molecule 1: Glutathione transferase family protein



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	97.36Å 59.22Å 79.97Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	37.60 – 2.63 37.60 – 2.63	Depositor EDS
% Data completeness (in resolution range)	98.0 (37.60-2.63) 93.1 (37.60-2.63)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	0.25 (at 2.61Å)	Xtrriage
Refinement program	PHENIX 1.9_1692	Depositor
R, $R_{free}$	0.188 , 0.269 0.195 , 0.266	Depositor DCC
$R_{free}$ test set	682 reflections (4.86%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	63.0	Xtrriage
Anisotropy	0.289	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.27 , 28.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.47$ , $\langle L^2 \rangle = 0.31$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	3607	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	71.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 5.21% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GSH

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.48	0/1829	0.57	0/2472
1	B	0.48	0/1829	0.59	0/2472
All	All	0.48	0/3658	0.58	0/4944

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1782	0	1771	35	0
1	B	1782	0	1771	38	1
2	A	20	0	15	4	0
2	B	20	0	15	0	0
3	A	2	0	0	0	0
3	B	1	0	0	0	0
All	All	3607	0	3572	69	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash

magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:GLN:HG2	2:A:301:GSH:HA31	1.30	1.14
1:A:105:LYS:NZ	1:B:67:GLU:OE1	2.00	0.94
1:A:6:VAL:CG2	1:A:31:TYR:HB3	2.02	0.89
1:A:6:VAL:HG23	1:A:31:TYR:HB2	1.56	0.88
1:A:6:VAL:HG23	1:A:31:TYR:CB	2.08	0.83
1:A:6:VAL:CG2	1:A:31:TYR:CB	2.59	0.80
1:A:4:ASP:OD1	1:A:5:GLN:N	2.16	0.79
1:A:4:ASP:O	1:A:5:GLN:HG3	1.84	0.75
1:B:27:LYS:HE2	1:B:85:LEU:HD13	1.70	0.73
1:B:36:GLU:OE2	1:B:42:SER:OG	2.07	0.71
1:B:73:GLN:HG2	1:B:86:LEU:HD11	1.78	0.65
1:A:3:SER:O	1:A:4:ASP:HB3	1.98	0.62
1:B:220:GLU:HG2	1:B:220:GLU:O	1.98	0.61
1:B:214:MET:HA	1:B:219:ILE:HG12	1.83	0.61
1:B:123:GLU:HA	1:B:126:LYS:HE2	1.83	0.59
1:A:12:TRP:CD1	1:A:13:PRO:HD3	2.37	0.58
1:B:210:VAL:O	1:B:214:MET:HG3	2.02	0.58
1:B:218:GLY:O	1:B:219:ILE:HG22	2.04	0.58
1:B:194:GLU:O	1:B:198:LYS:HG3	2.04	0.58
1:B:12:TRP:CD1	1:B:13:PRO:HD3	2.40	0.57
1:A:135:LEU:HD13	1:B:51:VAL:HG13	1.86	0.56
1:B:66:CYS:O	1:B:67:GLU:HB2	2.05	0.56
1:B:101:ASP:OD2	1:B:105:LYS:HE3	2.08	0.54
1:B:36:GLU:HG2	1:B:45:LEU:HD22	1.90	0.53
1:A:12:TRP:CG	1:A:13:PRO:HD3	2.44	0.52
1:A:14:SER:HB2	2:A:301:GSH:SG2	2.50	0.52
1:B:218:GLY:C	1:B:219:ILE:CG2	2.79	0.51
1:B:38:LEU:HD22	1:B:39:TRP:CE2	2.46	0.50
1:A:68:SER:O	1:A:72:VAL:HG23	2.10	0.50
1:B:30:LYS:HD2	1:B:30:LYS:H	1.77	0.50
1:A:91:TYR:OH	1:B:64:PRO:HD2	2.13	0.49
1:B:67:GLU:O	1:B:71:ILE:HG13	2.12	0.48
1:B:27:LYS:HE3	1:B:76:ASP:HA	1.95	0.48
1:A:63:LYS:HB3	1:B:91:TYR:CZ	2.48	0.48
1:A:14:SER:O	1:A:18:MET:HG3	2.13	0.47
1:B:219:ILE:O	1:B:219:ILE:HG13	2.14	0.47
1:B:118:LYS:NZ	1:B:219:ILE:O	2.25	0.47
1:A:116:THR:HA	1:A:219:ILE:HD12	1.97	0.47
1:B:12:TRP:CG	1:B:13:PRO:HD3	2.50	0.47
1:B:177:GLU:HA	1:B:184:ILE:HG13	1.97	0.46
1:A:18:MET:HB2	1:A:160:PRO:HB3	1.98	0.46

*Continued on next page...*



Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:54:GLN:HG2	2:A:301:GSH:CA3	2.21	0.46
1:B:171:ILE:HD13	1:B:214:MET:SD	2.56	0.45
1:A:85:LEU:HD13	1:A:156:ILE:HD13	1.98	0.45
1:A:215:LYS:HE3	1:A:215:LYS:HB2	1.73	0.44
1:A:102:PHE:CE1	1:A:106:LYS:HG3	2.53	0.44
1:A:67:GLU:HA	2:A:301:GSH:O11	2.18	0.43
1:B:159:VAL:HG12	1:B:160:PRO:HD3	2.00	0.43
1:B:218:GLY:O	1:B:219:ILE:CG2	2.66	0.43
1:A:144:PRO:HG2	1:A:149:GLU:HA	2.01	0.42
1:B:5:GLN:N	1:B:5:GLN:OE1	2.53	0.42
1:B:202:ASP:HA	1:B:203:PRO:HD3	1.88	0.42
1:A:6:VAL:HG22	1:A:31:TYR:CB	2.48	0.42
1:A:112:ARG:HE	1:A:112:ARG:HB2	1.63	0.42
1:A:213:LEU:HD23	1:A:213:LEU:HA	1.89	0.41
1:A:29:VAL:HG22	1:A:79:TRP:CD1	2.55	0.41
1:B:11:PHE:CD2	1:B:13:PRO:HD2	2.55	0.41
1:B:77:GLU:O	1:B:80:LYS:HE3	2.20	0.41
1:A:133:LEU:HD23	1:A:133:LEU:HA	1.88	0.41
1:B:22:LEU:HD23	1:B:156:ILE:HG23	2.02	0.41
1:B:27:LYS:HG3	1:B:75:ILE:HG22	2.01	0.41
1:B:44:LEU:HA	1:B:44:LEU:HD12	1.85	0.41
1:A:6:VAL:HG22	1:A:31:TYR:HB3	1.94	0.41
1:B:11:PHE:CE2	1:B:38:LEU:HD12	2.56	0.41
1:A:44:LEU:HD12	1:A:44:LEU:HA	1.79	0.40
1:A:211:LEU:HD23	1:A:211:LEU:HA	1.86	0.40
1:A:49:ASN:ND2	1:A:54:GLN:H	2.20	0.40
1:A:110:LEU:O	1:A:114:ILE:HG13	2.22	0.40
1:B:202:ASP:HB3	1:B:205:LYS:HD2	2.02	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:3:SER:OG	1:B:220:GLU:OE1[4_455]	1.77	0.43

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	216/231 (94%)	208 (96%)	6 (3%)	2 (1%)	17	33
1	B	216/231 (94%)	207 (96%)	7 (3%)	2 (1%)	17	33
All	All	432/462 (94%)	415 (96%)	13 (3%)	4 (1%)	17	33

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	5	GLN
1	A	4	ASP
1	B	218	GLY
1	B	67	GLU

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	193/204 (95%)	190 (98%)	3 (2%)	62	81
1	B	193/204 (95%)	183 (95%)	10 (5%)	23	44
All	All	386/408 (95%)	373 (97%)	13 (3%)	37	61

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	112	ARG
1	A	191	LEU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	A	204	GLN
1	B	66	CYS
1	B	85	LEU
1	B	135	LEU
1	B	174	PHE
1	B	186	TYR
1	B	193	LYS
1	B	194	GLU
1	B	214	MET
1	B	215	LYS
1	B	217	PHE

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GSH	B	301	-	18,19,19	0.71	0	23,24,24	0.83	0
2	GSH	A	301	-	18,19,19	0.70	0	23,24,24	0.83	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GSH	B	301	-	-	7/24/24/24	-
2	GSH	A	301	-	-	12/24/24/24	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (19) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
2	A	301	GSH	N1-CA1-CB1-CG1
2	A	301	GSH	C1-CA1-CB1-CG1
2	A	301	GSH	N2-CA2-CB2-SG2
2	B	301	GSH	N1-CA1-CB1-CG1
2	B	301	GSH	C1-CA1-CB1-CG1
2	A	301	GSH	O2-C2-N3-CA3
2	A	301	GSH	CA2-C2-N3-CA3
2	A	301	GSH	OE1-CD1-N2-CA2
2	A	301	GSH	CG1-CD1-N2-CA2
2	B	301	GSH	OE1-CD1-N2-CA2
2	B	301	GSH	CG1-CD1-N2-CA2
2	A	301	GSH	C2-CA2-CB2-SG2
2	B	301	GSH	N2-CA2-CB2-SG2
2	A	301	GSH	O11-C1-CA1-N1
2	A	301	GSH	O12-C1-CA1-N1
2	B	301	GSH	C2-CA2-CB2-SG2
2	B	301	GSH	C3-CA3-N3-C2
2	A	301	GSH	O32-C3-CA3-N3
2	A	301	GSH	O31-C3-CA3-N3

There are no ring outliers.

1 monomer is involved in 4 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	A	301	GSH	4	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	218/231 (94%)	-0.35	4 (1%) 68 64	56, 68, 94, 114	0
1	B	218/231 (94%)	-0.36	3 (1%) 75 71	58, 70, 87, 109	0
All	All	436/462 (94%)	-0.35	7 (1%) 72 68	56, 69, 92, 114	0

All (7) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	3	SER	3.4
1	B	220	GLU	3.0
1	B	217	PHE	2.7
1	A	217	PHE	2.6
1	A	219	ILE	2.5
1	A	220	GLU	2.5
1	A	213	LEU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)


There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GSH	A	301	20/20	0.86	0.17	65,78,90,93	0
2	GSH	B	301	20/20	0.87	0.21	68,79,87,96	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







## 6.5 Other polymers [i](#)

There are no such residues in this entry.