



Full wwPDB X-ray Structure Validation Report ⓘ

May 22, 2020 – 01:28 pm BST

PDB ID : 2JBX
Title : Crystal Structure of the myxoma virus anti-apoptotic protein M11L
Authors : Kvensakul, M.; Van Delft, M.F.; Lee, E.F.; Gulbis, J.M.; Fairlie, W.D.; Huang, D.C.S.; Colman, P.M.
Deposited on : 2006-12-14
Resolution : 2.73 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.73 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	1271 (2.76-2.72)
Clashscore	141614	1322 (2.76-2.72)
Ramachandran outliers	138981	1297 (2.76-2.72)
Sidechain outliers	138945	1298 (2.76-2.72)
RSRZ outliers	127900	1243 (2.76-2.72)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	145	 44% 31% 14% • 9%
1	B	145	 49% 34% • • 11%

2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 2132 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called M11L PROTEIN.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	132	1049	666	168	205	5	5	12	1	1
1	B	129	1016	643	163	200	5	5	5	0	1

- Molecule 2 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	43	Total	O	0	0
			43	43		
2	B	24	Total	O	0	0
			24	24		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: M11L PROTEIN

Chain A: 



- Molecule 1: M11L PROTEIN

Chain B: 



4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, α , β , γ	39.88Å 39.40Å 49.22Å 71.79° 90.09° 81.66°	Depositor
Resolution (Å)	20.00 – 2.73 46.70 – 2.73	Depositor EDS
% Data completeness (in resolution range)	97.3 (20.00-2.73) 97.3 (46.70-2.73)	Depositor EDS
R_{merge}	0.11	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.08 (at 2.73Å)	Xtrriage
Refinement program	REFMAC 5.2.0005	Depositor
R, R_{free}	0.214 , 0.286 0.218 , 0.235	Depositor DCC
R_{free} test set	336 reflections (4.61%)	wwPDB-VP
Wilson B-factor (Å ²)	46.0	Xtrriage
Anisotropy	0.321	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.33 , 53.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	2132	wwPDB-VP
Average B, all atoms (Å ²)	43.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 11.83% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.85	1/1060 (0.1%)	1.16	8/1424 (0.6%)
1	B	3.41	3/1023 (0.3%)	1.77	5/1376 (0.4%)
All	All	2.46	4/2083 (0.2%)	1.49	13/2800 (0.5%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	4	ARG	CZ-NH2	78.95	2.35	1.33
1	B	4	ARG	CZ-NH1	68.09	2.21	1.33
1	B	92	LYS	CE-NZ	-18.09	1.03	1.49
1	A	126	LYS	CG-CD	9.53	1.84	1.52

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	4	ARG	NH1-CZ-NH2	-53.76	60.27	119.40
1	A	126	LYS	CB-CG-CD	-15.56	71.14	111.60
1	A	74	ASP	CB-CG-OD1	-14.77	105.01	118.30
1	B	4	ARG	NE-CZ-NH1	-11.91	114.35	120.30
1	B	4	ARG	NE-CZ-NH2	10.85	125.73	120.30
1	A	126	LYS	CG-CD-CE	-9.00	84.90	111.90
1	B	92	LYS	CD-CE-NZ	8.65	131.59	111.70
1	A	51	ILE	CG1-CB-CG2	-6.63	96.81	111.40
1	A	81	LEU	CA-CB-CG	-6.44	100.48	115.30
1	A	74	ASP	OD1-CG-OD2	6.31	135.29	123.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	54	ARG	NE-CZ-NH1	6.16	123.38	120.30
1	A	43	LYS	CB-CG-CD	-5.61	97.03	111.60
1	B	57	LEU	CB-CG-CD1	5.01	119.52	111.00

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	4	ARG	Sidechain

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1049	0	1066	79	0
1	B	1016	0	1031	47	0
2	A	43	0	0	12	0
2	B	24	0	0	9	0
All	All	2132	0	2097	120	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 29.

All (120) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:65:LYS:HE3	2:A:2035:HOH:O	1.32	1.28
1:A:33:CYS:HB2	2:A:2040:HOH:O	1.15	1.24
1:A:35:ASP:HA	2:A:2016:HOH:O	1.38	1.23
1:A:92[A]:LYS:HG2	2:A:2027:HOH:O	1.30	1.23
1:B:47:THR:O	1:B:51:ILE:HG12	1.58	1.04
1:A:49:TYR:HE2	1:A:93:LYS:CE	1.76	0.97
1:A:57:LEU:O	1:A:58:SER:HB3	1.65	0.95
1:A:61:ILE:HG21	1:A:104:MSE:CE	1.99	0.91
1:A:49:TYR:CE2	1:A:93:LYS:CE	2.59	0.85

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:49:TYR:HE2	1:A:93:LYS:NZ	1.75	0.84
1:A:49:TYR:CE2	1:A:93:LYS:NZ	2.45	0.83
1:A:49:TYR:HE2	1:A:93:LYS:HZ2	1.27	0.82
1:A:54:ARG:NH2	1:B:54:ARG:HG3	1.97	0.80
1:B:14:ASN:HD21	1:B:102:ALA:H	1.29	0.79
1:B:17:ASP:HB3	2:B:2007:HOH:O	1.83	0.78
1:B:35:ASP:O	1:B:39:GLU:HG2	1.83	0.77
1:A:48:LEU:CD2	1:A:86:LEU:HD11	2.15	0.77
1:A:61:ILE:HG21	1:A:104:MSE:HE2	1.67	0.76
1:A:87:LEU:O	1:A:91:ILE:HG22	1.84	0.76
1:A:96:LYS:HG3	2:A:2029:HOH:O	1.88	0.73
1:A:49:TYR:CD1	1:A:49:TYR:C	2.61	0.73
1:A:49:TYR:HE2	1:A:93:LYS:CD	2.02	0.72
1:A:91:ILE:HD13	1:A:91:ILE:O	1.91	0.71
1:A:51:ILE:HG13	1:B:51:ILE:CG2	2.20	0.71
1:B:75:ALA:HB1	1:B:79:VAL:HG13	1.73	0.69
1:B:116:GLU:H	1:B:116:GLU:CD	1.97	0.69
1:B:18:ILE:CG2	1:B:24:MSE:HG3	2.25	0.67
1:A:48:LEU:HD22	1:A:86:LEU:HD11	1.76	0.66
1:A:91:ILE:HD13	1:A:91:ILE:C	2.17	0.65
1:B:18:ILE:HG23	1:B:24:MSE:HG3	1.78	0.64
1:A:49:TYR:CE2	1:A:93:LYS:HE3	2.31	0.64
1:B:48:LEU:O	1:B:51:ILE:HB	1.97	0.63
1:B:14:ASN:ND2	1:B:102:ALA:H	1.96	0.63
1:A:123:ILE:HG23	2:A:2040:HOH:O	1.98	0.63
1:A:49:TYR:HD1	1:A:49:TYR:C	2.02	0.62
1:A:6:LYS:NZ	1:A:110:ASP:OD2	2.33	0.62
1:B:49:TYR:CE2	1:B:93:LYS:HE2	2.35	0.62
1:A:51:ILE:HG13	1:B:51:ILE:HG23	1.81	0.61
1:A:30:LEU:CD2	1:A:84:LEU:HD21	2.31	0.60
1:A:26:LEU:O	1:A:30:LEU:HB2	2.02	0.60
1:A:49:TYR:CE2	1:A:93:LYS:CD	2.85	0.59
1:A:60:ASN:ND2	1:A:63:ASN:ND2	2.52	0.58
1:B:71:ALA:CB	2:B:2014:HOH:O	2.51	0.57
1:A:116:GLU:OE2	2:A:2038:HOH:O	2.18	0.57
1:A:49:TYR:CD2	1:A:93:LYS:HE3	2.40	0.57
1:A:30:LEU:HD21	1:A:84:LEU:HD21	1.88	0.56
1:B:45:TYR:CE2	1:B:85:THR:CG2	2.89	0.56
1:A:54:ARG:HH11	1:A:54:ARG:HG2	1.71	0.55
1:A:49:TYR:O	1:A:49:TYR:CD1	2.60	0.55
1:A:57:LEU:O	1:A:58:SER:CB	2.42	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:47:THR:O	1:B:51:ILE:CG1	2.44	0.54
1:A:107:GLU:H	1:A:107:GLU:CD	2.10	0.54
1:A:54:ARG:O	1:A:57:LEU:O	2.25	0.54
1:B:17:ASP:CB	2:B:2007:HOH:O	2.50	0.54
1:A:2:MSE:SE	1:A:6:LYS:HG2	2.58	0.54
1:A:84:LEU:HG	1:A:84:LEU:O	2.07	0.54
1:B:35:ASP:HA	2:B:2010:HOH:O	2.06	0.53
1:A:66:ASN:HD22	1:B:47:THR:HG21	1.73	0.53
1:A:76:SER:O	1:A:78:SER:N	2.41	0.53
1:A:17:ASP:O	1:A:19:THR:O	2.27	0.53
1:A:127:CYS:SG	1:A:127:CYS:O	2.67	0.53
1:A:95:ASN:OD1	1:A:100:THR:N	2.38	0.53
1:B:45:TYR:CE2	1:B:85:THR:HG22	2.44	0.53
1:A:54:ARG:HH21	1:B:54:ARG:HD2	1.74	0.52
1:A:68:LEU:CD2	1:A:86:LEU:HD13	2.39	0.52
1:A:91:ILE:HD11	2:A:2030:HOH:O	2.09	0.52
1:A:45:TYR:OH	1:A:82:ALA:HA	2.10	0.52
1:B:3:SER:O	1:B:4:ARG:HB2	2.10	0.52
1:A:95:ASN:HA	1:A:98:GLN:O	2.10	0.51
1:A:61:ILE:CG2	2:A:2035:HOH:O	2.58	0.51
1:A:54:ARG:HH21	1:B:54:ARG:HG3	1.70	0.51
1:A:61:ILE:HG21	1:A:104:MSE:HE3	1.87	0.51
1:B:101:ASP:HA	2:B:2020:HOH:O	2.12	0.49
1:A:98:GLN:HG3	1:A:99:HIS:N	2.27	0.49
1:A:30:LEU:HD21	1:A:84:LEU:CD2	2.43	0.49
1:B:29:GLN:HB3	1:B:123:ILE:CG2	2.43	0.49
1:B:89:SER:O	1:B:92:LYS:N	2.46	0.49
1:A:72:LEU:HD23	1:A:72:LEU:HA	1.66	0.48
1:B:34:CYS:HB3	2:B:2006:HOH:O	2.12	0.48
1:A:76:SER:O	1:A:77:PRO:C	2.52	0.47
1:B:6:LYS:NZ	1:B:110:ASP:OD2	2.31	0.47
1:B:89:SER:O	1:B:90:VAL:C	2.52	0.47
1:B:18:ILE:HG22	1:B:24:MSE:HG3	1.96	0.47
1:A:54:ARG:HG2	2:A:2018:HOH:O	2.15	0.47
1:A:33:CYS:O	1:A:37:ILE:HG13	2.15	0.47
1:A:25:ASP:O	1:A:29:GLN:HG3	2.15	0.47
1:A:0:PRO:HA	2:A:2002:HOH:O	2.15	0.46
1:A:65:LYS:HE2	1:A:107:GLU:OE1	2.16	0.46
1:B:73:ARG:HA	2:B:2015:HOH:O	2.14	0.46
1:A:90:VAL:O	1:A:94:LEU:HG	2.15	0.46
1:B:25:ASP:C	1:B:25:ASP:OD1	2.54	0.46

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:69:THR:O	1:A:69:THR:CG2	2.64	0.46
1:B:88:ALA:O	1:B:92:LYS:HG2	2.16	0.46
1:A:2:MSE:CE	1:A:7:THR:HA	2.46	0.45
1:B:17:ASP:C	1:B:18:ILE:O	2.54	0.45
1:B:45:TYR:OH	1:B:82:ALA:HA	2.16	0.45
1:B:53:GLU:O	1:B:57:LEU:HG	2.17	0.45
1:B:71:ALA:HB2	2:B:2014:HOH:O	2.16	0.45
1:B:70:PHE:HA	1:B:73:ARG:NH1	2.32	0.44
1:B:108:VAL:HG12	1:B:112:ILE:CD1	2.48	0.44
1:A:69:THR:O	1:A:69:THR:HG22	2.18	0.43
1:A:4:ARG:HD3	2:A:2003:HOH:O	2.17	0.43
1:B:108:VAL:HG12	1:B:112:ILE:HD12	1.99	0.43
1:B:14:ASN:HD21	1:B:102:ALA:N	2.07	0.43
1:B:56:ILE:C	1:B:57:LEU:O	2.56	0.43
1:A:48:LEU:HD23	1:A:86:LEU:HD11	1.98	0.42
1:A:29:GLN:HB3	1:A:29:GLN:HE21	1.58	0.42
1:A:128:LYS:O	1:A:129:TYR:C	2.58	0.42
1:A:72:LEU:HD13	1:A:111:GLY:HA3	2.01	0.42
1:A:2:MSE:HE3	1:A:7:THR:HA	2.01	0.42
1:A:95:ASN:OD1	1:A:99:HIS:HA	2.20	0.41
1:B:116:GLU:OE1	2:B:2022:HOH:O	2.22	0.41
1:B:61:ILE:HD11	1:B:104:MSE:HE1	2.02	0.41
1:A:6:LYS:HD2	1:A:10:TYR:HE2	1.86	0.41
1:A:54:ARG:O	1:A:58:SER:HB3	2.21	0.41
1:A:128:LYS:HG3	1:A:129:TYR:N	2.36	0.40
1:B:52:MSE:CE	1:B:64:ILE:HG23	2.52	0.40
1:B:76:SER:OG	1:B:78:SER:OG	2.35	0.40
1:A:45:TYR:OH	1:A:82:ALA:CA	2.69	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	131/145 (90%)	107 (82%)	18 (14%)	6 (5%)	2	2
1	B	127/145 (88%)	99 (78%)	25 (20%)	3 (2%)	6	9
All	All	258/290 (89%)	206 (80%)	43 (17%)	9 (4%)	3	5

All (9) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	129	TYR
1	A	130	ASN
1	B	4	ARG
1	B	18	ILE
1	B	72	LEU
1	A	106	SER
1	A	98	GLN
1	A	77	PRO
1	A	56	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	121/127 (95%)	95 (78%)	26 (22%)	1	1
1	B	117/127 (92%)	101 (86%)	16 (14%)	3	5
All	All	238/254 (94%)	196 (82%)	42 (18%)	2	2

All (42) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	6	LYS
1	A	19	THR
1	A	20	GLU
1	A	27	LEU
1	A	29	GLN
1	A	30	LEU
1	A	47	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	48	LEU
1	A	49	TYR
1	A	51	ILE
1	A	54	ARG
1	A	57	LEU
1	A	58	SER
1	A	72	LEU
1	A	81	LEU
1	A	86	LEU
1	A	87	LEU
1	A	91	ILE
1	A	93	LYS
1	A	101	ASP
1	A	107	GLU
1	A	113	VAL
1	A	116	GLU
1	A	127	CYS
1	A	128	LYS
1	A	129	TYR
1	B	2	MSE
1	B	9	VAL
1	B	18	ILE
1	B	22	THR
1	B	25	ASP
1	B	27	LEU
1	B	40	THR
1	B	58	SER
1	B	73	ARG
1	B	81	LEU
1	B	83	THR
1	B	85	THR
1	B	87	LEU
1	B	89	SER
1	B	115	GLU
1	B	128	LYS

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (7) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	29	GLN
1	A	63	ASN
1	A	66	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	117	GLN
1	A	124	GLN
1	B	14	ASN
1	B	124	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	127/145 (87%)	-0.13	0 100 100	22, 40, 60, 69	6 (4%)
1	B	124/145 (85%)	-0.06	0 100 100	29, 42, 59, 68	3 (2%)
All	All	251/290 (86%)	-0.10	0 100 100	22, 42, 60, 69	9 (3%)

There are no RSRZ outliers to report.

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.