



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 16, 2019 – 05:55 PM EDT

PDB ID : 6JIJ  
Title : The Crystal Structure of Main Protease from Mouse Hepatitis Virus A59 in Complex with an inhibitor  
Authors : Cui, W.; Cui, S.S.  
Deposited on : 2019-02-21  
Resolution : 2.65 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.0 (224370), CSD as540be (2019)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031633  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031633


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.65 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	1112 (2.68-2.64)
Clashscore	122126	1151 (2.68-2.64)
Ramachandran outliers	120053	1133 (2.68-2.64)
Sidechain outliers	120020	1133 (2.68-2.64)
RSRZ outliers	108989	1098 (2.68-2.64)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	 3% 78% 20% ..
1	B	301	 2% 84% 14% ..
1	C	301	 4% 82% 17% .
2	D	6	 50% 33% 17%
2	E	6	 50% 33% 17%

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	6	 33% 67%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	02J	F	1	-	-	-	X

## 2 Entry composition

There are 3 unique types of molecules in this entry. The entry contains 7002 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Replicative polyprotein 1ab.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	B	297	2279	1445	377	435	22	0	0	0
1	C	298	2286	1450	378	436	22	0	0	0
1	A	297	2279	1445	377	435	22	0	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	?	-	SER	deletion	UNP Q66WN6
B	?	-	ALA	deletion	UNP Q66WN6
B	282	PHE	LEU	conflict	UNP Q66WN6
C	?	-	SER	deletion	UNP Q66WN6
C	?	-	ALA	deletion	UNP Q66WN6
C	282	PHE	LEU	conflict	UNP Q66WN6
A	?	-	SER	deletion	UNP Q66WN6
A	?	-	ALA	deletion	UNP Q66WN6
A	282	PHE	LEU	conflict	UNP Q66WN6

- Molecule 2 is a protein called 02J-ALA-VAL-LEU-PJE-010.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	D	6	49	35	6	8	0	0	0
2	E	6	49	35	6	8	0	0	0
2	F	6	49	35	6	8	0	0	0


- Molecule 3 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>	<b>ZeroOcc</b>	<b>AltConf</b>
3	B	4	Total O 4 4	0	0
3	C	2	Total O 2 2	0	0
3	A	5	Total O 5 5	0	0


### 3 Residue-property plots i

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Replicative polyprotein 1ab



- Molecule 1: Replicative polyprotein 1ab



- Molecule 1: Replicative polyprotein 1ab



- Molecule 2: 02J-ALA-VAL-LEU-PJE-010






- Molecule 2: 02J-ALA-VAL-LEU-PJE-010



- Molecule 2: 02J-ALA-VAL-LEU-PJE-010



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	167.79Å 64.03Å 117.95Å 90.00° 125.97° 90.00°	Depositor
Resolution (Å)	36.96 – 2.65 36.96 – 2.65	Depositor EDS
% Data completeness (in resolution range)	99.3 (36.96-2.65) 99.4 (36.96-2.65)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.31 (at 2.65Å)	Xtrriage
Refinement program	PHENIX (1.14_3260: ???)	Depositor
R, $R_{free}$	0.226 , 0.270 0.225 , 0.267	Depositor DCC
$R_{free}$ test set	1503 reflections (5.06%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	59.5	Xtrriage
Anisotropy	0.709	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 39.8	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	7002	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	65.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.91% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: 010, PJE, 02J

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.29	0/2331	0.50	0/3173
1	B	0.29	0/2331	0.51	1/3173 (0.0%)
1	C	0.35	0/2338	0.54	0/3183
2	D	2.09	2/19 (10.5%)	1.29	0/25
2	E	2.07	2/19 (10.5%)	1.22	0/25
2	F	2.11	2/19 (10.5%)	1.34	0/25
All	All	0.36	6/7057 (0.1%)	0.53	1/9604 (0.0%)

All (6) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	3	VAL	C-N	6.42	1.48	1.34
2	D	3	VAL	C-N	6.24	1.48	1.34
2	E	3	VAL	C-N	6.19	1.48	1.34
2	F	2	ALA	C-N	5.25	1.46	1.34
2	D	2	ALA	C-N	5.16	1.46	1.34
2	E	2	ALA	C-N	5.14	1.45	1.34

All (1) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	244	LEU	CA-CB-CG	7.56	132.68	115.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2279	0	2225	41	0
1	B	2279	0	2225	30	2
1	C	2286	0	2234	34	0
2	D	49	0	45	3	0
2	E	49	0	45	2	0
2	F	49	0	45	2	2
3	A	5	0	0	0	0
3	B	4	0	0	0	0
3	C	2	0	0	0	0
All	All	7002	0	6819	99	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 7.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:275:GLN:HG3	1:C:278:GLY:HA2	1.29	1.09
1:B:163:LEU:HD21	2:D:2:ALA:HB1	1.64	0.80
1:A:191:LEU:HD22	1:A:192:PRO:HD2	1.66	0.77
1:C:275:GLN:HG3	1:C:278:GLY:CA	2.14	0.76
1:B:1:SER:OG	1:A:138:PHE:O	2.04	0.75
1:A:253:THR:HG23	1:A:255:VAL:H	1.51	0.74
1:B:7:VAL:HG21	1:A:7:VAL:HG21	1.70	0.71
1:C:275:GLN:HB2	1:C:279:SER:N	2.06	0.70
1:A:104:VAL:HG23	1:A:128:LEU:HD22	1.75	0.68
1:C:104:VAL:HG13	1:C:128:LEU:HD22	1.75	0.67
1:A:44:CYS:SG	1:A:45:SER:N	2.68	0.66
1:B:163:LEU:HD22	1:B:165:LEU:HD23	1.78	0.66
1:B:226:GLU:HA	1:B:229:ASN:HD22	1.60	0.66
1:C:221:ASP:OD1	1:C:266:ARG:NH1	2.29	0.65
1:B:116:TYR:HD2	1:B:121:GLN:HG3	1.62	0.65
1:C:164:GLU:HB3	2:E:3:VAL:HG23	1.77	0.65
1:C:275:GLN:HB2	1:C:279:SER:H	1.66	0.60
1:C:200:VAL:HG12	1:C:240:ILE:HG13	1.82	0.60

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:40:ARG:HA	1:A:85:LEU:HG	1.83	0.59
1:C:219:GLN:HB2	1:C:266:ARG:NH1	2.18	0.58
1:A:222:SER:HB3	1:A:258:GLU:HB3	1.87	0.56
1:A:207:TYR:CE1	1:A:253:THR:HG21	2.41	0.55
1:A:217:PHE:O	1:A:266:ARG:NH2	2.37	0.55
1:C:271:PHE:CD1	1:C:271:PHE:N	2.73	0.55
1:B:217:PHE:O	1:B:266:ARG:NH2	2.40	0.55
1:C:293:GLN:O	1:C:298:VAL:HB	2.08	0.54
1:C:30:LEU:HD13	1:C:146:VAL:HG11	1.90	0.54
1:C:219:GLN:HB2	1:C:266:ARG:HH12	1.73	0.53
1:C:185:ASP:OD1	1:C:185:ASP:N	2.42	0.53
1:C:129:ARG:NH2	1:C:284:ASP:OD2	2.38	0.53
1:B:225:LEU:HD12	1:B:225:LEU:H	1.73	0.53
1:A:207:TYR:HE1	1:A:253:THR:HG21	1.75	0.52
1:B:14:GLU:OE1	1:A:11:SER:HB3	2.10	0.51
1:C:109:THR:HG23	1:C:287:THR:HG23	1.92	0.51
1:A:211:PHE:CD1	1:A:295:LEU:HD13	2.46	0.51
1:B:104:VAL:HG13	1:B:108:GLU:HB2	1.93	0.50
1:B:6:MET:HE3	1:A:122:GLY:HA3	1.94	0.50
1:C:151:THR:HG23	1:C:154:SER:HB3	1.94	0.49
1:C:161:HIS:HE1	1:C:170:HIS:HB3	1.78	0.49
1:A:200:VAL:HG22	1:A:240:ILE:HG13	1.94	0.49
1:C:203:VAL:HG13	1:C:260:VAL:HG21	1.95	0.49
1:A:102:GLY:HA3	1:A:156:ARG:HH11	1.77	0.49
1:B:129:ARG:NH2	1:B:284:ASP:OD2	2.42	0.49
1:B:40:ARG:HA	1:B:85:LEU:HG	1.95	0.49
1:B:41:HIS:CD2	2:D:4:LEU:HD13	2.48	0.48
1:C:28:ASN:O	1:C:144:GLY:HA3	2.13	0.48
1:A:166:SER:OG	1:A:167:THR:N	2.47	0.48
1:A:219:GLN:O	1:A:259:GLN:NE2	2.46	0.48
1:A:226:GLU:HA	1:A:229:ASN:HD22	1.78	0.47
1:A:157:PHE:HB3	1:A:175:PHE:CE1	2.49	0.47
1:C:131:SER:N	1:C:195:ASP:OD1	2.46	0.47
1:B:7:VAL:HG22	1:A:123:ALA:O	2.15	0.47
1:B:163:LEU:HD23	1:B:164:GLU:N	2.30	0.46
1:A:109:THR:HG23	1:A:287:THR:HG23	1.97	0.46
1:B:226:GLU:OE1	1:B:229:ASN:ND2	2.49	0.46
1:C:47:MET:HG3	1:C:49:ASP:H	1.79	0.46
1:B:44:CYS:O	1:B:45:SER:HB3	2.15	0.46
1:C:132:HIS:HB3	1:C:180:TYR:HB3	1.98	0.46
1:B:59:VAL:O	1:B:78:TYR:OH	2.26	0.46

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:161:HIS:HE1	1:A:170:HIS:HB3	1.82	0.45
1:B:56:LEU:HD22	1:B:78:TYR:HE2	1.82	0.45
1:A:22:TYR:CD2	1:A:43:ILE:HA	2.52	0.45
1:B:31:TRP:CE2	1:B:93:ASN:HB2	2.52	0.45
1:C:161:HIS:CE1	1:C:170:HIS:HB3	2.52	0.44
1:B:225:LEU:O	1:B:229:ASN:ND2	2.50	0.44
1:B:41:HIS:NE2	2:D:4:LEU:HD13	2.33	0.44
1:A:44:CYS:C	1:A:46:ASP:H	2.21	0.44
1:B:122:GLY:HA2	1:A:9:PRO:HD3	2.00	0.44
1:C:21:THR:OG1	1:C:65:CYS:HB3	2.18	0.43
1:A:206:LEU:O	1:A:210:ILE:HG13	2.18	0.43
1:B:208:ALA:HB2	1:B:291:VAL:HG13	2.01	0.43
1:B:161:HIS:HE1	1:B:170:HIS:HB3	1.83	0.43
1:C:263:ALA:HA	1:C:266:ARG:HE	1.84	0.43
1:B:219:GLN:HG3	1:B:266:ARG:NH2	2.33	0.43
1:C:268:HIS:ND1	1:C:268:HIS:C	2.72	0.43
1:A:187:GLN:HE21	1:A:187:GLN:HB3	1.58	0.43
1:A:19:SER:OG	1:A:67:MET:HB2	2.19	0.42
1:B:40:ARG:HG3	1:B:52:TYR:CE1	2.55	0.42
1:C:81:GLN:O	1:C:177:GLY:HA3	2.19	0.42
1:C:143:CYS:HB2	2:E:5:PJE:H4	1.47	0.42
1:C:268:HIS:HA	1:C:282:PHE:CZ	2.54	0.42
1:A:291:VAL:O	1:A:295:LEU:HG	2.19	0.42
1:B:40:ARG:HD2	1:B:80:MET:HE2	2.01	0.42
1:C:12:LYS:NZ	1:C:153:ASP:HA	2.34	0.41
1:A:213:ARG:HH11	1:A:213:ARG:HA	1.84	0.41
1:A:67:MET:SD	1:A:72:SER:OG	2.64	0.41
1:C:268:HIS:HA	1:C:282:PHE:CE2	2.56	0.41
1:A:225:LEU:O	1:A:229:ASN:ND2	2.53	0.41
1:A:164:GLU:OE2	2:F:5:PJE:N6	2.53	0.41
1:A:40:ARG:HD3	1:A:83:CYS:HA	2.03	0.41
1:A:164:GLU:HB2	2:F:5:PJE:C29	2.51	0.41
1:A:249:LEU:O	1:A:253:THR:HG22	2.21	0.40
1:B:58:ARG:HA	1:B:58:ARG:CZ	2.51	0.40
1:C:291:VAL:O	1:C:295:LEU:HG	2.20	0.40
1:A:265:LYS:H	1:A:265:LYS:HG2	1.72	0.40
1:A:58:ARG:HD2	1:A:58:ARG:HA	1.93	0.40
1:A:3:ILE:HD11	1:A:295:LEU:HD21	2.03	0.40
1:A:21:THR:HB	1:A:65:CYS:HB3	2.04	0.40
1:C:116:TYR:CZ	1:C:142:SER:HB3	2.57	0.40

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the sym-

metry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:244:LEU:CD1	2:F:6:010:C3[2_555]	1.50	0.70
1:B:244:LEU:CD1	2:F:6:010:C2[2_555]	1.76	0.44

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	295/301 (98%)	275 (93%)	20 (7%)	0	100	100
1	B	295/301 (98%)	276 (94%)	19 (6%)	0	100	100
1	C	296/301 (98%)	283 (96%)	13 (4%)	0	100	100
2	D	1/6 (17%)	1 (100%)	0	0	100	100
2	E	1/6 (17%)	1 (100%)	0	0	100	100
2	F	1/6 (17%)	1 (100%)	0	0	100	100
All	All	889/921 (96%)	837 (94%)	52 (6%)	0	100	100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	259/263 (98%)	255 (98%)	4 (2%)	67	83
1	B	259/263 (98%)	253 (98%)	6 (2%)	53	72

*Continued on next page...*

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	C	260/263 (99%)	257 (99%)	3 (1%)	74	86
2	D	2/2 (100%)	2 (100%)	0	100	100
2	E	2/2 (100%)	1 (50%)	1 (50%)	0	0
2	F	2/2 (100%)	2 (100%)	0	100	100
All	All	784/795 (99%)	770 (98%)	14 (2%)	62	79

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	58	ARG
1	B	67	MET
1	B	91	LEU
1	B	153	ASP
1	B	187	GLN
1	B	283	GLU
1	C	48	THR
1	C	100	SER
1	C	275	GLN
1	A	58	ARG
1	A	74	THR
1	A	151	THR
1	A	187	GLN
2	E	3	VAL

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	272	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

6 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	02J	D	1	2	6,8,9	3.07	2 (33%)	4,10,12	2.36	1 (25%)
2	PJE	D	5	2	12,13,14	3.34	4 (33%)	11,16,18	1.43	2 (18%)
2	02J	E	1	2	6,8,9	3.01	2 (33%)	4,10,12	2.95	1 (25%)
2	PJE	E	5	2	12,13,14	3.44	4 (33%)	11,16,18	1.45	3 (27%)
2	02J	F	1	2	6,8,9	3.07	2 (33%)	4,10,12	2.83	1 (25%)
2	PJE	F	5	2	12,13,14	3.40	4 (33%)	11,16,18	1.39	2 (18%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	02J	D	1	2	-	0/0/2/4	0/1/1/1
2	PJE	D	5	2	-	1/7/18/19	0/1/1/1
2	02J	E	1	2	-	0/0/2/4	0/1/1/1
2	PJE	E	5	2	-	0/7/18/19	0/1/1/1
2	02J	F	1	2	-	0/0/2/4	0/1/1/1
2	PJE	F	5	2	-	1/7/18/19	0/1/1/1

All (18) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	5	PJE	C27-C28	-8.86	1.39	1.53
2	E	5	PJE	C27-C28	-8.82	1.39	1.53
2	D	5	PJE	C27-C28	-8.72	1.39	1.53
2	E	5	PJE	C25-C26	-5.36	1.41	1.53
2	D	5	PJE	C25-C26	-5.33	1.41	1.53
2	F	5	PJE	C25-C26	-5.19	1.41	1.53
2	D	5	PJE	C21-C22	3.05	1.53	1.44
2	F	5	PJE	C21-C22	3.12	1.53	1.44
2	E	5	PJE	C21-C22	3.44	1.54	1.44
2	D	5	PJE	C28-N6	3.66	1.53	1.46
2	E	5	PJE	C28-N6	3.67	1.53	1.46
2	F	5	PJE	C28-N6	3.74	1.54	1.46

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	1	02J	C6-C5	3.91	1.53	1.48
2	E	1	02J	C6-C5	3.93	1.53	1.48
2	D	1	02J	C6-C5	3.97	1.53	1.48
2	E	1	02J	C3-C41	5.99	1.55	1.48
2	D	1	02J	C3-C41	6.15	1.55	1.48
2	F	1	02J	C3-C41	6.22	1.55	1.48

All (10) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	E	1	02J	O42-C41-C3	-5.51	119.88	124.39
2	F	1	02J	O42-C41-C3	-5.24	120.10	124.39
2	D	1	02J	O42-C41-C3	-4.17	120.98	124.39
2	E	5	PJE	C25-C26-C27	-2.24	109.08	117.41
2	F	5	PJE	C25-C26-C27	-2.24	109.08	117.41
2	E	5	PJE	C28-N6-C29	-2.20	109.46	113.83
2	E	5	PJE	O7-C22-C21	-2.19	118.18	125.71
2	D	5	PJE	C25-C26-C27	-2.13	109.48	117.41
2	D	5	PJE	C28-N6-C29	-2.09	109.67	113.83
2	F	5	PJE	C28-N6-C29	-2.08	109.69	113.83

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	F	5	PJE	O7-C22-C21-C20
2	D	5	PJE	O7-C22-C21-C20

There are no ring outliers.

2 monomers are involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	E	5	PJE	1	0
2	F	5	PJE	2	0

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.



## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	297/301 (98%)	0.17	9 (3%) 50 46	44, 64, 88, 102	0
1	B	297/301 (98%)	-0.02	5 (1%) 70 67	41, 62, 84, 101	0
1	C	298/301 (99%)	0.15	12 (4%) 38 34	43, 63, 94, 108	0
2	D	3/6 (50%)	-0.22	0 100 100	58, 58, 58, 64	0
2	E	3/6 (50%)	0.47	0 100 100	68, 68, 70, 77	0
2	F	3/6 (50%)	0.35	0 100 100	72, 72, 74, 78	0
All	All	901/921 (97%)	0.10	26 (2%) 51 48	41, 63, 90, 108	0

All (26) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	152	GLY	5.1
1	A	47	MET	4.0
1	A	191	LEU	3.8
1	A	58	ARG	3.7
1	B	58	ARG	3.4
1	C	236	GLY	3.4
1	C	230	VAL	3.4
1	A	297	GLY	3.3
1	A	230	VAL	3.2
1	B	191	LEU	3.1
1	C	282	PHE	2.9
1	A	45	SER	2.7
1	C	226	GLU	2.7
1	C	276	ILE	2.6
1	C	234	THR	2.6
1	C	267	LEU	2.6
1	C	264	ILE	2.4
1	B	297	GLY	2.4
1	A	98	LYS	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	272	GLN	2.2
1	C	45	SER	2.2
1	C	218	VAL	2.2
1	B	192	PRO	2.1
1	C	241	LYS	2.1
1	C	261	LEU	2.1
1	A	54	ASN	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q<0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	02J	F	1	8/9	0.78	0.42	90,94,101,102	0
2	02J	E	1	8/9	0.86	0.27	78,88,92,92	0
2	PJE	E	5	13/14	0.90	0.18	53,58,75,75	0
2	PJE	F	5	13/14	0.91	0.20	61,67,78,78	0
2	02J	D	1	8/9	0.91	0.16	69,75,79,81	0
2	PJE	D	5	13/14	0.94	0.15	47,56,72,73	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.