



wwPDB EM Validation Summary Report ⓘ

Feb 29, 2024 – 10:12 PM JST

PDB ID : 8JME
EMDB ID : EMD-36417
Title : The cryo-EM structure of insect gustatory receptor Gr64a from *Drosophila melanogaster*
Authors : Ma, D.; Guo, J.
Deposited on : 2023-06-04
Resolution : 2.50 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/EMValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

EMDB validation analysis : **FAILED**
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
MapQ : **FAILED**
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.36


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

ELECTRON MICROSCOPY

The reported resolution of this entry is 2.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	EM structures (#Entries)
Clashscore	158937	4297
Ramachandran outliers	154571	4023
Sidechain outliers	154315	3826

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Mol	Chain	Length	Quality of chain
1	A	473	77% 9% 14%
1	B	473	78% 8% 14%
1	C	473	77% 9% 14%
1	D	473	77% 9% 14%

2 Entry composition [i](#)

There is only 1 type of molecule in this entry. The entry contains 27164 atoms, of which 13812 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Gustatory receptor for sugar taste 64a.

Mol	Chain	Residues	Atoms						AltConf	Trace
			Total	C	H	N	O	S		
1	A	407	6791	2206	3453	553	555	24	0	0
1	B	407	6791	2206	3453	553	555	24	0	0
1	C	407	6791	2206	3453	553	555	24	0	0
1	D	407	6791	2206	3453	553	555	24	0	0

There are 68 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-16	MET	-	initiating methionine	UNP P83293
A	-15	TRP	-	expression tag	UNP P83293
A	-14	SER	-	expression tag	UNP P83293
A	-13	HIS	-	expression tag	UNP P83293
A	-12	PRO	-	expression tag	UNP P83293
A	-11	GLN	-	expression tag	UNP P83293
A	-10	PHE	-	expression tag	UNP P83293
A	-9	GLU	-	expression tag	UNP P83293
A	-8	LYS	-	expression tag	UNP P83293
A	-7	GLY	-	expression tag	UNP P83293
A	-6	GLY	-	expression tag	UNP P83293
A	-5	SER	-	expression tag	UNP P83293
A	-4	SER	-	expression tag	UNP P83293
A	-3	GLY	-	expression tag	UNP P83293
A	-2	GLY	-	expression tag	UNP P83293
A	-1	VAL	-	expression tag	UNP P83293
A	0	ASP	-	expression tag	UNP P83293
B	-16	MET	-	initiating methionine	UNP P83293
B	-15	TRP	-	expression tag	UNP P83293
B	-14	SER	-	expression tag	UNP P83293
B	-13	HIS	-	expression tag	UNP P83293
B	-12	PRO	-	expression tag	UNP P83293

Continued on next page...

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-11	GLN	-	expression tag	UNP P83293
B	-10	PHE	-	expression tag	UNP P83293
B	-9	GLU	-	expression tag	UNP P83293
B	-8	LYS	-	expression tag	UNP P83293
B	-7	GLY	-	expression tag	UNP P83293
B	-6	GLY	-	expression tag	UNP P83293
B	-5	SER	-	expression tag	UNP P83293
B	-4	SER	-	expression tag	UNP P83293
B	-3	GLY	-	expression tag	UNP P83293
B	-2	GLY	-	expression tag	UNP P83293
B	-1	VAL	-	expression tag	UNP P83293
B	0	ASP	-	expression tag	UNP P83293
C	-16	MET	-	initiating methionine	UNP P83293
C	-15	TRP	-	expression tag	UNP P83293
C	-14	SER	-	expression tag	UNP P83293
C	-13	HIS	-	expression tag	UNP P83293
C	-12	PRO	-	expression tag	UNP P83293
C	-11	GLN	-	expression tag	UNP P83293
C	-10	PHE	-	expression tag	UNP P83293
C	-9	GLU	-	expression tag	UNP P83293
C	-8	LYS	-	expression tag	UNP P83293
C	-7	GLY	-	expression tag	UNP P83293
C	-6	GLY	-	expression tag	UNP P83293
C	-5	SER	-	expression tag	UNP P83293
C	-4	SER	-	expression tag	UNP P83293
C	-3	GLY	-	expression tag	UNP P83293
C	-2	GLY	-	expression tag	UNP P83293
C	-1	VAL	-	expression tag	UNP P83293
C	0	ASP	-	expression tag	UNP P83293
D	-16	MET	-	initiating methionine	UNP P83293
D	-15	TRP	-	expression tag	UNP P83293
D	-14	SER	-	expression tag	UNP P83293
D	-13	HIS	-	expression tag	UNP P83293
D	-12	PRO	-	expression tag	UNP P83293
D	-11	GLN	-	expression tag	UNP P83293
D	-10	PHE	-	expression tag	UNP P83293
D	-9	GLU	-	expression tag	UNP P83293
D	-8	LYS	-	expression tag	UNP P83293
D	-7	GLY	-	expression tag	UNP P83293
D	-6	GLY	-	expression tag	UNP P83293
D	-5	SER	-	expression tag	UNP P83293
D	-4	SER	-	expression tag	UNP P83293

Continued on next page...


Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
D	-3	GLY	-	expression tag	UNP P83293
D	-2	GLY	-	expression tag	UNP P83293
D	-1	VAL	-	expression tag	UNP P83293
D	0	ASP	-	expression tag	UNP P83293


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Gustatory receptor for sugar taste 64a



- Molecule 1: Gustatory receptor for sugar taste 64a




- Molecule 1: Gustatory receptor for sugar taste 64a



- Molecule 1: Gustatory receptor for sugar taste 64a

Chain D: 77% 9% 14%



4 Experimental information

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	372952	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE CORRECTION	Depositor
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose ($e^-/\text{\AA}^2$)	52	Depositor
Minimum defocus (nm)	800	Depositor
Maximum defocus (nm)	1500	Depositor
Magnification	Not provided	
Image detector	FEI FALCON IV (4k x 4k)	Depositor

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.25	0/3422	0.48	0/4641
1	B	0.26	0/3422	0.48	0/4641
1	C	0.26	0/3422	0.49	0/4641
1	D	0.25	0/3422	0.48	0/4641
All	All	0.26	0/13688	0.48	0/18564

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3338	3453	3453	22	0
1	B	3338	3453	3453	21	0
1	C	3338	3453	3453	22	0
1	D	3338	3453	3453	22	0
All	All	13352	13812	13812	79	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

The worst 5 of 79 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:423:ARG:NH1	1:C:417:PHE:O	2.00	0.95
1:C:423:ARG:NH1	1:D:417:PHE:O	2.00	0.94
1:A:417:PHE:O	1:D:423:ARG:NH1	2.01	0.94
1:A:423:ARG:NH1	1:B:417:PHE:O	2.01	0.93
1:C:274:LEU:HD13	1:C:312:LEU:HD12	1.79	0.64

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	405/473 (86%)	392 (97%)	13 (3%)	0	100	100
1	B	405/473 (86%)	392 (97%)	13 (3%)	0	100	100
1	C	405/473 (86%)	392 (97%)	13 (3%)	0	100	100
1	D	405/473 (86%)	392 (97%)	13 (3%)	0	100	100
All	All	1620/1892 (86%)	1568 (97%)	52 (3%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	368/426 (86%)	360 (98%)	8 (2%)	52	77
1	B	368/426 (86%)	360 (98%)	8 (2%)	52	77

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	C	368/426 (86%)	360 (98%)	8 (2%)	52 77
1	D	368/426 (86%)	360 (98%)	8 (2%)	52 77
All	All	1472/1704 (86%)	1440 (98%)	32 (2%)	54 77

5 of 32 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	D	286	ARG
1	D	294	CYS
1	B	286	ARG
1	B	225	ASN
1	D	416	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.