



# Full wwPDB X-ray Structure Validation Report ⓘ

Sep 11, 2023 – 12:19 AM EDT

PDB ID : 4JN6  
Title : Crystal Structure of the Aldolase-Dehydrogenase Complex from Mycobacterium tuberculosis HRv37  
Authors : Carere, J.; McKenna, S.E.; Kimber, M.S.; Seah, S.Y.K.  
Deposited on : 2013-03-14  
Resolution : 1.93 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.93 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4310 (1.96-1.92)
Clashscore	141614	1023 (1.94-1.94)
Ramachandran outliers	138981	1007 (1.94-1.94)
Sidechain outliers	138945	1007 (1.94-1.94)
RSRZ outliers	127900	4250 (1.96-1.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	346	 3% 89% 8%
1	C	346	 92% 6%
2	B	306	 15% 85% 12%
2	D	306	 18% 87% 9%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard

residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
3	OXL	A	401	-	X	-	-

## 2 Entry composition

There are 6 unique types of molecules in this entry. The entry contains 9781 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 4-hydroxy-2-oxovalerate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	338	2499	1559	449	478	13	0	0	0
1	C	339	2504	1562	450	479	13	0	0	0


- Molecule 2 is a protein called Acetaldehyde dehydrogenase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	296	2205	1397	372	429	7	0	0	0
2	D	296	2205	1397	372	429	7	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	-2	GLY	-	expression tag	UNP P71866
B	-1	SER	-	expression tag	UNP P71866
B	0	HIS	-	expression tag	UNP P71866
D	-2	GLY	-	expression tag	UNP P71866
D	-1	SER	-	expression tag	UNP P71866
D	0	HIS	-	expression tag	UNP P71866

- Molecule 3 is OXALATE ION (three-letter code: OXL) (formula: C<sub>2</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C O 6 2 4	0	0
3	C	1	Total C O 6 2 4	0	0

- Molecule 4 is MANGANESE (II) ION (three-letter code: MN) (formula: Mn).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Mn 1 1	0	0
4	C	1	Total Mn 1 1	0	0

- Molecule 5 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	C	1	Total Na 1 1	0	0

- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	101	Total O 101 101	0	0
6	B	42	Total O 42 42	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
6	C	132	Total 132	O 132	0	0
6	D	78	Total 78	O 78	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 4-hydroxy-2-oxovalerate aldolase




- Molecule 1: 4-hydroxy-2-oxovalerate aldolase




- Molecule 2: Acetaldehyde dehydrogenase



- Molecule 2: Acetaldehyde dehydrogenase







## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.69Å 142.69Å 148.17Å 90.00° 95.08° 90.00°	Depositor
Resolution (Å)	45.89 – 1.93 45.89 – 1.93	Depositor EDS
% Data completeness (in resolution range)	98.1 (45.89-1.93) 98.1 (45.89-1.93)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.14 (at 1.92Å)	Xtrriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
R, $R_{free}$	0.174 , 0.206 0.177 , 0.208	Depositor DCC
$R_{free}$ test set	5296 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.4	Xtrriage
Anisotropy	0.192	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.38 , 51.0	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	9781	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	54.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.59% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: MN, NA, OXL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.31	0/2536	0.47	0/3425
1	C	0.38	0/2541	0.51	0/3432
2	B	0.28	0/2250	0.48	0/3078
2	D	0.31	0/2250	0.51	0/3078
All	All	0.32	0/9577	0.49	0/13013

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2499	0	2486	17	0
1	C	2504	0	2491	10	0
2	B	2205	0	2211	20	0
2	D	2205	0	2211	16	0
3	A	6	0	0	0	0
3	C	6	0	0	0	0
4	A	1	0	0	0	0
4	C	1	0	0	0	0
5	C	1	0	0	0	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
6	A	101	0	0	2	0
6	B	42	0	0	3	0
6	C	132	0	0	2	0
6	D	78	0	0	6	1
All	All	9781	0	9399	59	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:265:VAL:O	6:C:565:HOH:O	1.89	0.90
1:A:122:GLN:NE2	6:A:576:HOH:O	2.15	0.79
2:B:49:LYS:NZ	6:B:440:HOH:O	2.24	0.70
1:A:82:ARG:NH1	1:A:133:GLU:OE1	2.27	0.67
2:D:242:GLU:OE1	6:D:455:HOH:O	2.12	0.66
2:D:164:ARG:O	6:D:416:HOH:O	2.13	0.66
2:B:7:VAL:HG12	2:B:71:LEU:HB3	1.78	0.65
2:B:186:ARG:NH2	6:B:417:HOH:O	2.31	0.64
2:D:293:ALA:HA	2:D:297:LEU:HB2	1.78	0.64
2:D:163:THR:N	6:D:475:HOH:O	2.31	0.62
1:A:101:ARG:NH2	1:A:102:ASP:OD1	2.33	0.61
1:A:94:LYS:HB3	1:A:126:LEU:HD22	1.83	0.59
2:D:108:ILE:HG22	2:D:110:PRO:HD2	1.84	0.58
2:B:107:VAL:HG11	2:B:113:LEU:HD13	1.85	0.57
2:B:238:ARG:NH1	2:B:263:GLU:OE2	2.37	0.56
2:B:108:ILE:HG22	2:B:110:PRO:HD2	1.87	0.56
2:D:39:PRO:HG3	2:D:56:HIS:CD2	2.42	0.55
2:B:35:VAL:HG21	2:B:62:LEU:HD22	1.89	0.54
1:C:17:GLY:HA2	1:C:229:GLY:HA2	1.89	0.54
1:A:58:PHE:HE1	2:B:252:SER:HB2	1.73	0.54
2:D:109:PRO:HA	2:D:113:LEU:HB2	1.89	0.54
1:C:94:LYS:HB3	1:C:126:LEU:HD22	1.89	0.53
1:A:236:PRO:HG2	1:A:239:ALA:HB3	1.90	0.53
2:D:21:TYR:OH	2:D:46:ARG:HD2	2.12	0.50
2:D:71:LEU:HD12	2:D:94:ARG:HB2	1.94	0.50
2:D:271:LEU:HD13	2:D:278:LEU:HB2	1.94	0.50
2:D:246:ASP:OD2	6:D:446:HOH:O	2.20	0.49
2:B:17:THR:O	2:B:20:LEU:HB3	2.12	0.49
1:A:295:HIS:O	1:A:299:GLN:HG2	2.13	0.49

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:35:VAL:HG21	2:D:62:LEU:HD22	1.95	0.48
1:A:114:CYS:O	1:A:152:GLN:HG3	2.14	0.48
1:A:295:HIS:CD2	1:A:324:GLU:HG2	2.48	0.48
1:C:237:VAL:O	1:C:241:ILE:HG12	2.14	0.47
2:B:109:PRO:HA	2:B:113:LEU:HB2	1.97	0.46
1:A:237:VAL:O	1:A:241:ILE:HG12	2.16	0.46
1:A:142:HIS:HD2	6:A:513:HOH:O	1.99	0.46
2:D:160:GLY:O	6:D:475:HOH:O	2.20	0.45
2:B:15:ILE:HD11	2:B:278:LEU:HD22	1.98	0.45
1:A:254:ASP:HB3	1:A:257:ASP:HB2	1.99	0.45
1:A:281:LEU:CD1	1:C:238:GLU:HG3	2.47	0.44
2:B:57:GLU:HB2	2:B:61:TRP:HB2	1.98	0.44
2:D:103:VAL:N	6:D:438:HOH:O	2.51	0.44
2:B:125:ILE:HG23	2:B:126:THR:H	1.82	0.44
1:A:246:LYS:HE3	6:C:544:HOH:O	2.17	0.44
2:B:20:LEU:HD21	2:B:50:LEU:HD12	1.99	0.43
1:C:114:CYS:O	1:C:152:GLN:HG3	2.18	0.43
2:D:28:TRP:CE2	2:D:294:LYS:HD3	2.54	0.43
2:D:240:LEU:HD21	2:D:263:GLU:HB2	2.01	0.43
1:C:295:HIS:CD2	1:C:324:GLU:HG2	2.54	0.43
2:B:268:GLY:HA2	2:B:271:LEU:O	2.20	0.42
1:A:309:ALA:O	1:A:313:ARG:HG3	2.20	0.42
1:A:281:LEU:HD11	1:C:238:GLU:HG3	2.02	0.41
2:B:199:PRO:HB2	2:B:274:TYR:CG	2.55	0.41
1:C:116:GLU:O	1:C:119:VAL:HG22	2.21	0.41
2:B:223:ILE:O	2:B:227:VAL:HG23	2.21	0.41
1:A:238:GLU:HG3	1:C:281:LEU:CD1	2.51	0.41
2:B:144:GLU:O	2:B:211:PRO:HD3	2.21	0.41
2:B:201:ILE:HD13	2:B:274:TYR:HA	2.03	0.41
2:B:46:ARG:NH1	6:B:435:HOH:O	2.21	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:D:448:HOH:O	6:D:448:HOH:O[2_557]	2.11	0.09

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	336/346 (97%)	331 (98%)	3 (1%)	2 (1%)	25	13
1	C	337/346 (97%)	333 (99%)	2 (1%)	2 (1%)	25	13
2	B	294/306 (96%)	283 (96%)	10 (3%)	1 (0%)	41	32
2	D	294/306 (96%)	285 (97%)	8 (3%)	1 (0%)	41	32
All	All	1261/1304 (97%)	1232 (98%)	23 (2%)	6 (0%)	29	17

All (6) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	228	PHE
1	C	228	PHE
1	A	175	LEU
1	C	175	LEU
2	B	125	ILE
2	D	125	ILE

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	251/257 (98%)	248 (99%)	3 (1%)	71	64
1	C	251/257 (98%)	246 (98%)	5 (2%)	55	42
2	B	231/237 (98%)	227 (98%)	4 (2%)	60	49
2	D	231/237 (98%)	227 (98%)	4 (2%)	60	49

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
All	All	964/988 (98%)	948 (98%)	16 (2%)	60 49

All (16) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	48	THR
1	A	228	PHE
1	A	301	GLU
2	B	56	HIS
2	B	63	LEU
2	B	79	TYR
2	B	297	LEU
1	C	4	MET
1	C	48	THR
1	C	101	ARG
1	C	228	PHE
1	C	301	GLU
2	D	114	ARG
2	D	226	VAL
2	D	257	LEU
2	D	297	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry

Of 5 ligands modelled in this entry, 3 are monoatomic - leaving 2 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
3	OXL	C	401	4	5,5,5	1.55	0	6,6,6	1.51	1 (16%)
3	OXL	A	401	4	5,5,5	1.41	0	6,6,6	1.75	2 (33%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	OXL	C	401	4	-	0/4/4/4	-
3	OXL	A	401	4	-	4/4/4/4	-

There are no bond length outliers.

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	A	401	OXL	O3-C1-C2	2.68	121.13	113.16
3	C	401	OXL	O4-C2-C1	2.56	120.78	113.16
3	A	401	OXL	O4-C2-C1	2.38	120.23	113.16

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	401	OXL	O1-C1-C2-O2
3	A	401	OXL	O1-C1-C2-O4
3	A	401	OXL	O3-C1-C2-O2
3	A	401	OXL	O3-C1-C2-O4

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	338/346 (97%)	0.35	11 (3%) 46 54	36, 49, 68, 95	0
1	C	339/346 (97%)	0.12	1 (0%) 94 96	28, 39, 54, 85	0
2	B	296/306 (96%)	1.00	47 (15%) 1 2	48, 69, 104, 120	0
2	D	296/306 (96%)	0.80	56 (18%) 1 1	27, 49, 100, 121	0
All	All	1269/1304 (97%)	0.55	115 (9%) 9 14	27, 49, 92, 121	0

All (115) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	63	LEU	10.4
2	B	61	TRP	10.0
2	B	64	ALA	7.8
2	D	62	LEU	7.5
2	B	37	ILE	6.7
2	B	57	GLU	6.6
2	D	297	LEU	6.5
2	B	66	PRO	6.2
2	D	52	LEU	6.0
2	B	79	TYR	5.9
2	B	60	ASP	5.6
2	D	64	ALA	5.2
1	A	341	GLY	5.2
2	B	88	TYR	5.0
2	B	39	PRO	4.7
2	B	91	ALA	4.6
2	D	39	PRO	4.6
2	D	61	TRP	4.6
2	D	55	THR	4.5
2	B	38	ASP	4.5
2	D	35	VAL	4.4

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	D	59	VAL	4.4
2	D	37	ILE	4.2
2	B	67	ASP	4.2
2	D	93	ILE	4.2
2	D	296	THR	4.2
2	B	58	GLY	4.1
2	B	298	VAL	4.0
2	D	41	SER	4.0
2	B	59	VAL	4.0
2	B	68	LYS	3.9
2	D	68	LYS	3.8
2	B	86	PRO	3.8
2	B	297	LEU	3.8
2	B	35	VAL	3.7
2	D	20	LEU	3.6
2	D	298	VAL	3.5
2	D	33	TRP	3.5
2	D	66	PRO	3.5
2	D	90	GLU	3.5
2	D	91	ALA	3.5
2	D	69	PRO	3.5
1	A	340	SER	3.4
2	D	67	ASP	3.4
2	B	296	THR	3.4
2	D	48	ALA	3.4
2	B	56	HIS	3.4
2	B	293	ALA	3.4
2	D	103	VAL	3.3
2	B	90	GLU	3.3
2	D	50	LEU	3.3
1	A	4	MET	3.3
2	B	33	TRP	3.3
2	D	292	ILE	3.2
2	B	40	GLU	3.2
1	A	338	LEU	3.2
2	D	47	ALA	3.1
2	D	63	LEU	3.1
2	D	88	TYR	3.1
2	B	93	ILE	3.1
2	D	34	MET	3.0
2	B	55	THR	3.0
2	B	216	ARG	3.0

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	B	295	GLU	3.0
2	B	114	ARG	3.0
2	D	46	ARG	2.9
2	D	5	ALA	2.9
2	B	62	LEU	2.9
1	A	191	GLU	2.8
2	D	6	LYS	2.8
1	A	88	LEU	2.8
2	D	32	ARG	2.8
2	B	89	ALA	2.8
2	D	44	LEU	2.8
2	D	57	GLU	2.8
2	D	94	ARG	2.7
2	D	45	ALA	2.7
2	D	86	PRO	2.7
2	D	56	HIS	2.6
2	B	214	ALA	2.6
2	D	38	ASP	2.6
2	D	40	GLU	2.5
2	B	213	ASP	2.5
2	D	43	GLY	2.4
2	D	53	GLU	2.4
2	B	65	GLN	2.4
2	D	92	GLY	2.4
2	D	114	ARG	2.4
1	A	192	ASP	2.4
2	B	83	ASP	2.4
2	B	246	ASP	2.4
2	D	42	ASP	2.4
2	B	142	ILE	2.4
2	D	71	LEU	2.4
2	D	36	GLY	2.4
2	D	54	THR	2.3
2	B	220	ALA	2.3
2	B	82	ARG	2.3
2	D	28	TRP	2.3
1	A	170	ALA	2.2
2	D	95	ALA	2.2
1	A	339	ASP	2.2
2	D	58	GLY	2.2
2	D	60	ASP	2.2
2	B	71	LEU	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	240	LEU	2.1
1	A	129	GLU	2.1
2	B	190	ILE	2.1
1	C	205	LEU	2.1
2	D	118	ASP	2.1
2	B	245	PHE	2.1
2	D	102	ALA	2.1
2	B	94	ARG	2.0
2	B	84	ALA	2.0
1	A	336	ARG	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	OXL	A	401	6/6	0.94	0.21	27,38,43,52	6
3	OXL	C	401	6/6	0.96	0.18	22,31,33,42	6
5	NA	C	402	1/1	0.97	0.08	37,37,37,37	0
4	MN	C	403	1/1	0.98	0.14	34,34,34,34	0
4	MN	A	402	1/1	0.98	0.17	41,41,41,41	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.