



# wwPDB X-ray Structure Validation Summary Report ⓘ

Oct 3, 2021 – 02:19 PM EDT

PDB ID : 3JR6  
Title : Sequential reorganization of beta-sheet topology by insertion of a single strand  
Authors : Sagermann, M.; Baas, W.A.; Matthews, B.W.  
Deposited on : 2009-09-08  
Resolution : 3.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.23.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.23.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 3.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	2416 (3.00-3.00)
Ramachandran outliers	138981	2333 (3.00-3.00)
Sidechain outliers	138945	2336 (3.00-3.00)
RSRZ outliers	127900	1990 (3.00-3.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	170	
1	B	170	
1	C	170	
1	D	170	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	SO4	A	806	-	-	X	-
2	SO4	D	804	-	-	X	-

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 5252 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Lysozyme.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	170	1351	852	246	248	5	0	0	0
1	B	156	1249	787	228	229	5	0	0	0
1	C	159	1270	801	231	233	5	0	0	0
1	D	169	1342	846	245	246	5	0	0	0

There are 32 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	28	GLY	-	insertion	UNP P00720
A	29	ILE	-	insertion	UNP P00720
A	30	GLY	-	insertion	UNP P00720
A	31	HIS	-	insertion	UNP P00720
A	32	LEU	-	insertion	UNP P00720
A	33	LEU	-	insertion	UNP P00720
A	60	THR	CYS	engineered mutation	UNP P00720
A	103	ALA	CYS	engineered mutation	UNP P00720
B	28	GLY	-	insertion	UNP P00720
B	29	ILE	-	insertion	UNP P00720
B	30	GLY	-	insertion	UNP P00720
B	31	HIS	-	insertion	UNP P00720
B	32	LEU	-	insertion	UNP P00720
B	33	LEU	-	insertion	UNP P00720
B	60	THR	CYS	engineered mutation	UNP P00720
B	103	ALA	CYS	engineered mutation	UNP P00720
C	28	GLY	-	insertion	UNP P00720
C	29	ILE	-	insertion	UNP P00720
C	30	GLY	-	insertion	UNP P00720
C	31	HIS	-	insertion	UNP P00720
C	32	LEU	-	insertion	UNP P00720

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
C	33	LEU	-	insertion	UNP P00720
C	60	THR	CYS	engineered mutation	UNP P00720
C	103	ALA	CYS	engineered mutation	UNP P00720
D	28	GLY	-	insertion	UNP P00720
D	29	ILE	-	insertion	UNP P00720
D	30	GLY	-	insertion	UNP P00720
D	31	HIS	-	insertion	UNP P00720
D	32	LEU	-	insertion	UNP P00720
D	33	LEU	-	insertion	UNP P00720
D	60	THR	CYS	engineered mutation	UNP P00720
D	103	ALA	CYS	engineered mutation	UNP P00720

- Molecule 2 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total O S 5 4 1	0	0
2	A	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	B	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0
2	C	1	Total O S 5 4 1	0	0

Continued on next page...


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>			<b>ZeroOcc</b>	<b>AltConf</b>
2	D	1	Total	O	S	0	0
			5	4	1		
2	D	1	Total	O	S	0	0
			5	4	1		


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Lysozyme




- Molecule 1: Lysozyme





- Molecule 1: Lysozyme





- Molecule 1: Lysozyme





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	59.23Å 75.92Å 81.55Å 90.00° 94.06° 90.00°	Depositor
Resolution (Å)	32.44 – 3.00 32.44 – 3.00	Depositor EDS
% Data completeness (in resolution range)	96.5 (32.44-3.00) 96.6 (32.44-3.00)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.07	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	12.49 (at 3.00Å)	Xtrriage
Refinement program	CCP4 (refmac), CNS	Depositor
R, $R_{free}$	0.250 , 0.320 0.260 , (Not available)	Depositor DCC
$R_{free}$ test set	No test flags present.	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.4	Xtrriage
Anisotropy	0.359	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.30 , 22.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.88	EDS
Total number of atoms	5252	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	2.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 36.40 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 5.0257e-04. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	2.50	8/1372 (0.6%)	1.68	14/1847 (0.8%)
1	B	0.86	0/1267	0.98	0/1704
1	C	0.86	1/1288 (0.1%)	1.03	3/1731 (0.2%)
1	D	0.97	3/1363 (0.2%)	1.08	5/1836 (0.3%)
All	All	1.49	12/5290 (0.2%)	1.23	22/7118 (0.3%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1
1	D	0	1
All	All	0	2

The worst 5 of 12 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	44	SER	CB-OG	78.26	2.44	1.42
1	A	45	LEU	CB-CG	20.12	2.10	1.52
1	A	42	SER	CB-OG	17.58	1.65	1.42
1	A	43	PRO	N-CD	15.37	1.69	1.47
1	D	46	ASN	CB-CG	12.96	1.80	1.51

The worst 5 of 22 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	42	SER	C-N-CD	-29.57	55.54	120.60
1	A	43	PRO	N-CA-CB	26.86	135.53	103.30

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	43	PRO	CA-N-CD	-16.39	88.55	111.50
1	A	44	SER	CA-CB-OG	-16.32	67.15	111.20
1	A	45	LEU	CB-CG-CD2	13.94	134.70	111.00

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	61	ASN	Mainchain
1	D	24	TYR	Sidechain

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1351	0	1382	151	0
1	B	1249	0	1270	143	0
1	C	1270	0	1295	108	0
1	D	1342	0	1371	99	0
2	A	10	0	0	3	0
2	B	10	0	0	1	0
2	C	10	0	0	0	0
2	D	10	0	0	3	0
All	All	5252	0	5318	488	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 46.

The worst 5 of 488 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:46:ASN:CB	1:D:46:ASN:CG	1.80	1.46
1:A:43:PRO:CG	1:A:43:PRO:CB	1.90	1.46
1:A:42:SER:OG	1:A:42:SER:CB	1.65	1.43
1:A:24:TYR:CE2	1:A:32:LEU:HD22	1.61	1.35
1:B:64:ILE:CG2	1:B:68:GLU:HB3	1.60	1.31

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	168/170 (99%)	135 (80%)	25 (15%)	8 (5%)	2	13
1	B	152/170 (89%)	129 (85%)	14 (9%)	9 (6%)	1	9
1	C	155/170 (91%)	132 (85%)	16 (10%)	7 (4%)	2	14
1	D	167/170 (98%)	143 (86%)	21 (13%)	3 (2%)	8	37
All	All	642/680 (94%)	539 (84%)	76 (12%)	27 (4%)	3	16

5 of 27 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	43	PRO
1	A	44	SER
1	B	17	ILE
1	B	82	ARG
1	C	168	LYS

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	141/141 (100%)	121 (86%)	20 (14%)	3	16
1	B	129/141 (92%)	120 (93%)	9 (7%)	15	47
1	C	131/141 (93%)	122 (93%)	9 (7%)	15	48

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	D	140/141 (99%)	127 (91%)	13 (9%)	9	33
All	All	541/564 (96%)	490 (91%)	51 (9%)	8	32

5 of 51 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	B	169	ASN
1	C	138	ASN
1	D	129	GLN
1	C	10	ASP
1	C	75	GLN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 21 such sidechains are listed below:

Mol	Chain	Res	Type
1	C	129	GLN
1	D	129	GLN
1	D	150	ASN
1	D	138	ASN
1	D	75	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

8 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and

the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	SO4	D	801	-	4,4,4	0.25	0	6,6,6	0.32	0
2	SO4	D	804	-	4,4,4	0.37	0	6,6,6	0.55	0
2	SO4	C	802	-	4,4,4	0.43	0	6,6,6	0.37	0
2	SO4	B	808	-	4,4,4	0.36	0	6,6,6	0.31	0
2	SO4	A	806	-	4,4,4	0.27	0	6,6,6	0.33	0
2	SO4	B	807	-	4,4,4	0.43	0	6,6,6	0.42	0
2	SO4	A	805	-	4,4,4	0.39	0	6,6,6	0.23	0
2	SO4	C	803	-	4,4,4	0.39	0	6,6,6	0.45	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

5 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	D	801	SO4	1	0
2	D	804	SO4	2	0
2	A	806	SO4	2	0
2	B	807	SO4	1	0
2	A	805	SO4	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	170/170 (100%)	-0.37	2 (1%) 79 54	1, 2, 8, 19	0
1	B	156/170 (91%)	-0.34	2 (1%) 77 51	1, 2, 6, 8	0
1	C	159/170 (93%)	-0.35	2 (1%) 77 51	1, 2, 5, 19	0
1	D	169/170 (99%)	-0.41	2 (1%) 79 54	1, 2, 6, 11	0
All	All	654/680 (96%)	-0.37	8 (1%) 79 54	1, 2, 6, 19	0

The worst 5 of 8 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	61	ASN	4.2
1	A	24	TYR	3.2
1	C	34	GLY	3.2
1	D	40	THR	2.9
1	D	42	SER	2.7

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled ‘Q< 0.9’ lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	SO4	D	804	5/5	0.93	0.18	1,4,13,16	0
2	SO4	C	802	5/5	0.95	0.17	1,1,12,12	0
2	SO4	A	805	5/5	0.95	0.19	1,6,11,16	0
2	SO4	C	803	5/5	0.96	0.17	1,1,3,3	0
2	SO4	B	807	5/5	0.96	0.13	1,3,11,13	0
2	SO4	D	801	5/5	0.97	0.14	1,1,1,4	0
2	SO4	B	808	5/5	0.97	0.12	1,1,6,6	0
2	SO4	A	806	5/5	0.99	0.07	1,1,1,2	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.