



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 3, 2024 – 04:34 PM EST

PDB ID : 1JSH
Title : CRYSTAL STRUCTURE OF H9 HAEMAGGLUTININ COMPLEXED WITH LSTA RECEPTOR ANALOG
Authors : Ha, Y.; Stevens, D.J.; Skehel, J.J.; Wiley, D.C.
Deposited on : 2001-08-17
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	319	
2	B	176	
3	C	2	
3	E	2	
3	G	2	
4	D	3	
5	F	2	

2 Entry composition [i](#)

There are 7 unique types of molecules in this entry. The entry contains 4053 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called HAEMAGGLUTININ (HA1 CHAIN).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	317	2472	1551	435	473	13	0	0	0

- Molecule 2 is a protein called HAEMAGGLUTININ (HA2 CHAIN).


Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	160	1251	779	214	250	8	0	0	0

- Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
3	C	2	28	16	2	10	0	0	0
3	E	2	28	16	2	10	0	0	0
3	G	2	28	16	2	10	0	0	0

- Molecule 4 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose.




Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
4	D	3	42	24	3	15	0	0	0

- Molecule 5 is an oligosaccharide called N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	F	2	32	17	1	14	0	0	0

- Molecule 6 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
6	A	1	14	8	1	5	0	0

- Molecule 7 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	A	86	Total	O	0	0
			86	86		

Continued on next page...

Continued from previous page...


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
7	B	72	Total	O	0	0
			72	72		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.

- Molecule 1: HAEMAGGLUTININ (HA1 CHAIN)



- Molecule 2: HAEMAGGLUTININ (HA2 CHAIN)



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose



- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose




- Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  100%

MAG1
MAG2

- Molecule 4: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-[2-acetamido-2-deoxy-beta-D-glucopyranose-(1-6)]2-acetamido-2-deoxy-beta-D-glucopyranose

Chain D:  67% 33%

MAG1
MAG2
MAG3

- Molecule 5: N-acetyl-alpha-neuraminic acid-(2-3)-beta-D-galactopyranose

Chain F:  50% 50%

GAL1
SIA2

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 63	Depositor
Cell constants a, b, c, α , β , γ	108.56Å 108.56Å 149.07Å 90.00° 90.00° 120.00°	Depositor
Resolution (Å)	40.00 – 2.40	Depositor
% Data completeness (in resolution range)	(Not available) (40.00-2.40)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.200 , 0.227	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	4053	wwPDB-VP
Average B, all atoms (Å ²)	34.0	wwPDB-VP

5 Model quality

5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, GAL, SIA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.34	0/2531	0.64	0/3452
2	B	0.36	0/1272	0.60	0/1716
All	All	0.35	0/3803	0.63	0/5168

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2472	0	2406	61	0
2	B	1251	0	1181	11	0
3	C	28	0	25	2	0
3	E	28	0	25	2	0
3	G	28	0	25	0	0
4	D	42	0	37	1	0
5	F	32	0	28	2	0
6	A	14	0	13	0	0
7	A	86	0	0	4	0
7	B	72	0	0	0	0
All	All	4053	0	3740	73	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (73) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:303:LYS:HD2	3:E:1:NAG:H4	1.66	0.78
1:A:300:VAL:HG12	1:A:302:VAL:HG22	1.64	0.78
1:A:298:LYS:HE2	2:B:92:TRP:CE3	2.27	0.70
1:A:162:ARG:HB3	1:A:162:ARG:NH1	2.08	0.68
1:A:175:PRO:HG2	1:A:181:GLN:NE2	2.08	0.68
1:A:162:ARG:HB3	1:A:162:ARG:HH11	1.61	0.66
1:A:153:ILE:HD12	1:A:153:ILE:N	2.10	0.65
1:A:187:ARG:HD2	1:A:238:ASN:ND2	2.11	0.65
1:A:77:SER:O	1:A:254:SER:HB2	1.97	0.65
1:A:187:ARG:HD2	1:A:238:ASN:HD22	1.62	0.64
3:C:1:NAG:H61	3:C:2:NAG:H82	1.81	0.63
1:A:153:ILE:HD12	1:A:153:ILE:H	1.64	0.63
1:A:271:GLN:HE22	1:A:281:THR:H	1.47	0.62
1:A:142:ARG:HB3	7:A:381:HOH:O	1.99	0.62
1:A:192:THR:HG23	1:A:237:SER:HB2	1.83	0.61
3:C:1:NAG:C6	3:C:2:NAG:H82	2.31	0.60
1:A:175:PRO:HG2	1:A:181:GLN:HE21	1.65	0.60
1:A:144:LEU:HD23	7:A:381:HOH:O	2.01	0.59
2:B:125:ALA:O	2:B:160:ARG:HA	2.05	0.57
1:A:207:ILE:HD12	1:A:207:ILE:N	2.20	0.56
1:A:185:TYR:O	1:A:186:THR:HB	2.04	0.56
2:B:158:ASN:O	2:B:159:ARG:CB	2.54	0.56
1:A:60:LEU:C	1:A:60:LEU:HD23	2.27	0.56
1:A:126:TYR:HA	7:A:381:HOH:O	2.06	0.55
1:A:162:ARG:HH11	1:A:162:ARG:CB	2.21	0.53
1:A:186:THR:O	1:A:186:THR:HG22	2.08	0.53
2:B:60:ASN:HD22	2:B:60:ASN:C	2.12	0.53
1:A:181:GLN:HG2	1:A:188:THR:HA	1.91	0.53
1:A:96:GLU:OE2	1:A:201:ARG:NH2	2.41	0.52
1:A:75:GLU:HA	1:A:109:SER:O	2.10	0.51
1:A:271:GLN:NE2	1:A:281:THR:H	2.09	0.50
1:A:153:ILE:H	1:A:153:ILE:CD1	2.22	0.50
1:A:41:LEU:HD13	1:A:80:VAL:HG21	1.95	0.49
1:A:271:GLN:NE2	1:A:280:ASN:HA	2.27	0.49
1:A:154:GLN:O	1:A:236:ARG:HA	2.13	0.48
1:A:73:GLY:HA2	1:A:111:TYR:CE2	2.48	0.48
2:B:30:GLN:NE2	2:B:145:ASP:HB2	2.29	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:278:GLY:HA3	7:A:361:HOH:O	2.13	0.47
1:A:93:GLY:HA3	1:A:220:ILE:O	2.14	0.47
1:A:233:LEU:HD12	1:A:234:ARG:N	2.30	0.47
1:A:173:ASN:ND2	1:A:185:TYR:OH	2.47	0.47
1:A:233:LEU:HD12	1:A:234:ARG:H	1.80	0.46
1:A:153:ILE:N	1:A:153:ILE:CD1	2.77	0.46
4:D:2:NAG:H82	4:D:3:NAG:H2	1.96	0.46
1:A:115:GLN:NE2	1:A:118:PRO:HB3	2.30	0.46
1:A:52:LEU:HB3	1:A:55:CYS:O	2.17	0.45
1:A:53:ASP:HA	1:A:82:ARG:HB2	1.99	0.44
1:A:190:THR:OG1	1:A:240:ASN:ND2	2.50	0.44
1:A:151:TYR:HB3	1:A:187:ARG:HH12	1.83	0.44
1:A:272:CYS:SG	1:A:292:ALA:HB1	2.58	0.43
2:B:131:MET:HB3	2:B:141:TYR:OH	2.17	0.43
1:A:186:THR:O	1:A:186:THR:CG2	2.66	0.43
1:A:128:GLY:CA	5:F:2:SIA:H113	2.49	0.43
2:B:130:ALA:HB2	2:B:140:LEU:HD23	2.00	0.43
2:B:131:MET:HG3	2:B:139:GLU:HB2	2.01	0.43
1:A:34:HIS:O	1:A:286:HIS:HA	2.19	0.42
2:B:143:LYS:HD3	2:B:143:LYS:HA	1.87	0.42
1:A:207:ILE:N	1:A:207:ILE:CD1	2.82	0.42
1:A:271:GLN:HE21	1:A:280:ASN:HA	1.85	0.42
1:A:192:THR:HG21	1:A:241:LEU:HB2	2.02	0.41
1:A:177:THR:OG1	1:A:179:THR:HG23	2.20	0.41
1:A:196:THR:HA	1:A:232:THR:O	2.20	0.41
1:A:192:THR:HG21	1:A:241:LEU:HD13	2.01	0.41
1:A:190:THR:HA	1:A:238:ASN:HD21	1.85	0.41
1:A:128:GLY:HA3	5:F:2:SIA:H113	2.01	0.41
1:A:198:ASP:OD2	1:A:231:GLN:NE2	2.54	0.41
1:A:19:LEU:HD22	2:B:102:LEU:HD11	2.02	0.41
1:A:168:PHE:C	1:A:169:MET:SD	2.99	0.41
1:A:296:CYS:HA	1:A:297:PRO:HD3	1.88	0.41
1:A:300:VAL:CG1	1:A:302:VAL:HG22	2.43	0.41
2:B:60:ASN:HD22	2:B:61:LYS:N	2.18	0.41
1:A:276:ARG:HH12	3:E:1:NAG:H82	1.86	0.40
1:A:91:TYR:CE1	1:A:216:LEU:HD13	2.56	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	315/319 (99%)	302 (96%)	13 (4%)	0	100	100
2	B	158/176 (90%)	152 (96%)	5 (3%)	1 (1%)	22	33
All	All	473/495 (96%)	454 (96%)	18 (4%)	1 (0%)	44	59

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
2	B	159	ARG

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	279/282 (99%)	274 (98%)	5 (2%)	54	73
2	B	132/150 (88%)	128 (97%)	4 (3%)	36	57
All	All	411/432 (95%)	402 (98%)	9 (2%)	47	67

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	22	THR
1	A	53	ASP
1	A	146	GLN
1	A	162	ARG
1	A	179	THR

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	47	LYS
2	B	59	MET
2	B	60	ASN
2	B	131	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	97	ASN
1	A	115	GLN
1	A	146	GLN
1	A	157	GLN
1	A	173	ASN
1	A	181	GLN
1	A	240	ASN
1	A	271	GLN
2	B	30	GLN
2	B	60	ASN
2	B	68	HIS
2	B	78	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
3	NAG	C	1	3,1	14,14,15	0.53	0	17,19,21	0.66	0
3	NAG	C	2	3	14,14,15	0.70	1 (7%)	17,19,21	0.60	0
4	NAG	D	1	1,4	14,14,15	0.56	0	17,19,21	0.82	1 (5%)
4	NAG	D	2	4	14,14,15	0.55	0	17,19,21	0.74	1 (5%)
4	NAG	D	3	4	14,14,15	0.73	0	17,19,21	0.64	0
3	NAG	E	1	3,1	14,14,15	0.97	1 (7%)	17,19,21	1.33	3 (17%)
3	NAG	E	2	3	14,14,15	0.82	1 (7%)	17,19,21	0.74	0
5	GAL	F	1	5	12,12,12	0.51	0	17,17,17	0.37	0
5	SIA	F	2	5	20,20,21	0.65	0	21,28,31	0.91	1 (4%)
3	NAG	G	1	2,3	14,14,15	0.62	0	17,19,21	0.66	0
3	NAG	G	2	3	14,14,15	0.65	0	17,19,21	0.65	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	C	1	3,1	-	4/6/23/26	0/1/1/1
3	NAG	C	2	3	-	5/6/23/26	0/1/1/1
4	NAG	D	1	1,4	-	4/6/23/26	0/1/1/1
4	NAG	D	2	4	-	4/6/23/26	0/1/1/1
4	NAG	D	3	4	-	4/6/23/26	0/1/1/1
3	NAG	E	1	3,1	-	0/6/23/26	0/1/1/1
3	NAG	E	2	3	-	4/6/23/26	0/1/1/1
5	GAL	F	1	5	-	0/2/22/22	0/1/1/1
5	SIA	F	2	5	-	0/18/34/38	0/1/1/1
3	NAG	G	1	2,3	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	-	4/6/23/26	0/1/1/1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	E	1	NAG	C1-C2	2.74	1.56	1.52
3	E	2	NAG	C1-C2	2.61	1.55	1.52
3	C	2	NAG	C1-C2	2.02	1.55	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	F	2	SIA	O1B-C1-C2	2.43	119.03	112.71
3	E	1	NAG	C4-C3-C2	-2.35	107.58	111.02
3	E	1	NAG	C2-N2-C7	-2.31	119.80	122.90
4	D	2	NAG	C2-N2-C7	-2.25	119.89	122.90
4	D	1	NAG	C2-N2-C7	-2.17	120.00	122.90
3	E	1	NAG	C3-C4-C5	-2.17	106.31	110.23

There are no chirality outliers.

All (29) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
3	C	1	NAG	C8-C7-N2-C2
3	C	1	NAG	O7-C7-N2-C2
3	C	2	NAG	C1-C2-N2-C7
3	C	2	NAG	C8-C7-N2-C2
3	C	2	NAG	O7-C7-N2-C2
3	E	2	NAG	C8-C7-N2-C2
3	E	2	NAG	O7-C7-N2-C2
4	D	3	NAG	C8-C7-N2-C2
4	D	3	NAG	O7-C7-N2-C2
4	D	2	NAG	C4-C5-C6-O6
3	G	2	NAG	C8-C7-N2-C2
3	G	2	NAG	O7-C7-N2-C2
4	D	1	NAG	C8-C7-N2-C2
4	D	1	NAG	O7-C7-N2-C2
3	G	2	NAG	O5-C5-C6-O6
4	D	2	NAG	O5-C5-C6-O6
4	D	3	NAG	O5-C5-C6-O6
4	D	3	NAG	C4-C5-C6-O6
4	D	2	NAG	C8-C7-N2-C2
3	G	2	NAG	C4-C5-C6-O6
4	D	1	NAG	O5-C5-C6-O6
3	C	1	NAG	O5-C5-C6-O6
4	D	2	NAG	O7-C7-N2-C2
3	C	1	NAG	C4-C5-C6-O6
4	D	1	NAG	C4-C5-C6-O6
3	C	2	NAG	O5-C5-C6-O6
3	C	2	NAG	C4-C5-C6-O6
3	E	2	NAG	C4-C5-C6-O6
3	E	2	NAG	O5-C5-C6-O6


There are no ring outliers.


6 monomers are involved in 7 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	D	2	NAG	1	0
3	E	1	NAG	2	0
3	C	2	NAG	2	0
4	D	3	NAG	1	0
3	C	1	NAG	2	0
5	F	2	SIA	2	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.











5.6 Ligand geometry [i](#)

1 ligand is modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
6	NAG	A	320	1	14,14,15	0.47	0	17,19,21	0.64	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	NAG	A	320	1	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (2) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	A	320	NAG	C8-C7-N2-C2
6	A	320	NAG	O7-C7-N2-C2

There are no ring outliers.

No monomer is involved in short contacts.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.