



# Full wwPDB X-ray Structure Validation Report ⓘ

May 29, 2020 – 05:14 am BST

PDB ID : 4JTE  
Title : Crystal structure of F114A mutant of 3-deoxy-D-manno-octulosonate 8-phosphate synthase (KDO8PS) from *Neisseria meningitidis*  
Authors : Allison, T.M.; Cochrane, F.C.; Jameson, G.B.; Parker, E.J.  
Deposited on : 2013-03-23  
Resolution : 1.90 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.90 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	6207 (1.90-1.90)
Clashscore	141614	6847 (1.90-1.90)
Ramachandran outliers	138981	6760 (1.90-1.90)
Sidechain outliers	138945	6760 (1.90-1.90)
RSRZ outliers	127900	6082 (1.90-1.90)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	280	 9% 84% 5% 10%
1	B	280	 5% 78% 11% • 9%
1	C	280	 3% 76% 13% • 10%
1	D	280	 15% 80% 10% • 8%

## 2 Entry composition [i](#)

There are 4 unique types of molecules in this entry. The entry contains 8085 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2-dehydro-3-deoxyphosphooctonate aldolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	251	1909	1225	325	348	11	0	1	0
1	B	256	1956	1250	335	360	11	0	0	0
1	C	253	1945	1249	331	354	11	0	1	0
1	D	257	1948	1247	335	355	11	0	1	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	114	ALA	PHE	ENGINEERED MUTATION	UNP Q9JZ55
B	114	ALA	PHE	ENGINEERED MUTATION	UNP Q9JZ55
C	114	ALA	PHE	ENGINEERED MUTATION	UNP Q9JZ55
D	114	ALA	PHE	ENGINEERED MUTATION	UNP Q9JZ55

- Molecule 2 is SODIUM ION (three-letter code: NA) (formula: Na).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	C	1	Total	Na	0	0
			1	1		

- Molecule 3 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	D	1	Total	Cl	0	0
			1	1		
3	C	1	Total	Cl	0	0
			1	1		


- Molecule 4 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
4	A	80	Total 80	O 80	0	0
4	B	95	Total 95	O 95	0	0
4	C	89	Total 89	O 89	0	0
4	D	60	Total 60	O 60	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase




- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase




- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase



- Molecule 1: 2-dehydro-3-deoxyphosphooctonate aldolase





## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.36Å 85.47Å 163.06Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.95 – 1.90 39.95 – 1.90	Depositor EDS
% Data completeness (in resolution range)	97.2 (39.95-1.90) 97.3 (39.95-1.90)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.22 (at 1.91Å)	Xtrriage
Refinement program	REFMAC	Depositor
R, $R_{free}$	0.229 , 0.267 0.232 , 0.266	Depositor DCC
$R_{free}$ test set	4369 reflections (5.02%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	39.2	Xtrriage
Anisotropy	0.241	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 37.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.36$	Xtrriage
Estimated twinning fraction	0.010 for k,h,-l	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	8085	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.75% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup> Intensities estimated from amplitudes.

<sup>2</sup> Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NA, CL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.78	0/1941	0.84	2/2625 (0.1%)
1	B	0.87	1/1987 (0.1%)	0.89	4/2683 (0.1%)
1	C	0.87	0/1978	0.87	2/2674 (0.1%)
1	D	0.80	0/1979	0.88	5/2680 (0.2%)
All	All	0.83	1/7885 (0.0%)	0.87	13/10662 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	95	GLU	CG-CD	5.14	1.59	1.51

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed( $^{\circ}$ )	Ideal( $^{\circ}$ )
1	B	67	ARG	NE-CZ-NH1	5.95	123.28	120.30
1	D	201	LEU	CA-CB-CG	5.79	128.62	115.30
1	D	165	ARG	NE-CZ-NH2	5.66	123.13	120.30
1	D	201	LEU	CB-CA-C	-5.64	99.49	110.20
1	C	252	LEU	CA-CB-CG	5.54	128.03	115.30
1	D	67	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	C	92	ASP	CB-CG-OD1	5.53	123.28	118.30
1	D	235	LEU	CB-CG-CD2	-5.47	101.69	111.00
1	B	177	ASP	CB-CG-OD2	-5.46	113.38	118.30
1	A	201	LEU	CB-CA-C	-5.44	99.87	110.20
1	B	117	ARG	NE-CZ-NH1	-5.37	117.61	120.30
1	A	201	LEU	CA-CB-CG	5.21	127.29	115.30
1	B	201	LEU	CB-CA-C	-5.08	100.56	110.20

There are no chirality outliers.

There are no planarity outliers.



## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1909	0	1939	10	0
1	B	1956	0	2001	30	0
1	C	1945	0	1988	29	0
1	D	1948	0	1973	24	0
2	C	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	80	0	0	2	0
4	B	95	0	0	4	0
4	C	89	0	0	2	0
4	D	60	0	0	2	0
All	All	8085	0	7901	88	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (88) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:263:ARG:NH2	1:D:263:ARG:NH1	2.09	1.01
1:D:196:ASP:OD2	1:D:199:HIS:HD2	1.61	0.83
1:C:196:ASP:OD2	1:C:199[A]:HIS:HD2	1.63	0.82
1:D:263:ARG:NH2	1:D:263:ARG:NE	2.27	0.81
1:D:165:ARG:HH21	1:D:165:ARG:HG3	1.45	0.80
1:D:263:ARG:NE	1:D:263:ARG:NH1	2.29	0.79
1:A:165:ARG:HG3	1:A:165:ARG:HH11	1.49	0.78
1:D:144:GLN:HG3	4:D:448:HOH:O	1.89	0.73
1:C:140:LEU:HD23	1:C:145:MET:CE	2.20	0.71
1:C:141:SER:OG	1:C:144:GLN:HG3	1.90	0.70
1:C:82:LYS:HE2	4:C:442:HOH:O	1.89	0.70
1:A:165:ARG:CG	1:A:165:ARG:HH11	2.04	0.70
1:B:140:LEU:HD23	1:B:145:MET:CE	2.22	0.69
1:D:165:ARG:NH2	1:D:165:ARG:HG3	2.08	0.68
1:B:199:HIS:HD2	4:B:386:HOH:O	1.78	0.64
1:C:196:ASP:OD2	1:C:199[A]:HIS:CD2	2.49	0.64

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:255:HIS:CE1	1:C:256:LEU:HD23	2.33	0.64
1:C:173:ASN:OD1	1:C:202:GLN:NE2	2.31	0.64
1:D:73:GLU:O	1:D:76:LYS:HG3	1.98	0.63
1:B:23:ASN:HD21	1:B:57:LYS:HE3	1.65	0.62
1:B:45:LEU:O	1:B:265:LYS:HE3	2.01	0.61
1:D:196:ASP:OD2	1:D:199:HIS:CD2	2.49	0.61
1:B:144:GLN:HG3	4:B:373:HOH:O	2.01	0.60
1:B:71:LEU:C	1:B:71:LEU:HD13	2.22	0.60
1:C:255:HIS:CE1	1:C:256:LEU:CD2	2.85	0.59
1:C:255:HIS:CD2	1:C:256:LEU:CD2	2.85	0.59
1:C:255:HIS:NE2	1:C:256:LEU:HD21	2.17	0.59
1:B:277:LEU:O	1:B:277:LEU:HD13	2.02	0.59
1:C:26:GLU:OE1	1:C:66:TYR:HD2	1.86	0.59
1:C:23:ASN:HD21	1:C:57:LYS:HE3	1.69	0.58
1:D:165:ARG:CG	1:D:165:ARG:HH21	2.13	0.58
1:B:199:HIS:HB2	1:B:236:GLU:OE2	2.03	0.58
1:D:145:MET:HA	1:D:145:MET:HE2	1.85	0.58
1:A:165:ARG:HG3	1:A:165:ARG:NH1	2.14	0.58
1:A:60:ARG:NH2	1:C:120:ASP:OD1	2.29	0.57
1:B:198:THR:HG21	1:B:236:GLU:HB2	1.86	0.57
4:A:373:HOH:O	1:C:94:HIS:HD2	1.86	0.57
1:B:140:LEU:HD23	1:B:145:MET:HE2	1.88	0.55
1:C:45:LEU:O	1:C:265:LYS:HE3	2.07	0.55
1:C:140:LEU:HD23	1:C:145:MET:HE2	1.89	0.54
1:D:116:ALA:HB2	4:D:460:HOH:O	2.07	0.54
1:C:256:LEU:CD2	1:C:256:LEU:N	2.73	0.52
1:C:255:HIS:NE2	1:C:256:LEU:CD2	2.74	0.51
1:B:22:ILE:HG23	1:B:34:THR:HG21	1.92	0.51
1:B:23:ASN:ND2	1:B:57:LYS:HE3	2.24	0.51
1:D:169:PHE:HB2	1:D:175[B]:VAL:HG22	1.93	0.50
1:D:69:VAL:HG13	1:D:73:GLU:CB	2.42	0.50
1:B:22:ILE:HD11	1:B:25:LEU:HD23	1.94	0.49
1:B:276:ILE:O	1:B:277:LEU:HB2	2.13	0.49
1:C:256:LEU:HB3	4:C:482:HOH:O	2.13	0.48
1:C:4:LYS:HE3	1:C:9:THR:OG1	2.13	0.48
1:B:63:ILE:HG22	1:D:119:THR:HG23	1.96	0.47
1:C:22:ILE:HD11	1:C:25:LEU:HD23	1.96	0.47
1:B:52:LYS:HE2	1:B:234:PHE:CZ	2.50	0.47
1:B:56:ASP:OD2	1:B:94:HIS:HE1	1.97	0.47
1:A:260:PHE:HD2	4:A:316:HOH:O	1.97	0.47
1:A:22:ILE:HG23	1:A:34:THR:HG21	1.97	0.47

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:201:LEU:HD11	1:B:218:ALA:HA	1.96	0.47
1:D:63:ILE:HG13	1:D:64:HIS:CD2	2.49	0.46
1:C:277:LEU:HD23	1:C:277:LEU:C	2.35	0.46
1:A:22:ILE:HD11	1:A:25:LEU:HD23	1.98	0.46
1:B:202:GLN:HG3	1:B:202:GLN:H	1.49	0.46
1:B:198:THR:CG2	1:B:236:GLU:HB2	2.46	0.45
1:B:196:ASP:OD2	1:B:236:GLU:OE2	2.34	0.45
1:C:255:HIS:ND1	1:C:256:LEU:HD23	2.32	0.45
1:B:63:ILE:HA	1:D:119:THR:CG2	2.47	0.45
1:B:71:LEU:CD1	1:B:71:LEU:C	2.85	0.45
1:B:63:ILE:HA	1:D:119:THR:HG21	1.99	0.44
1:D:237:SER:HA	1:D:252:LEU:O	2.17	0.44
1:D:141:SER:OG	1:D:144:GLN:HG2	2.19	0.43
1:D:238:HIS:HB2	1:D:239:PRO:HD2	1.99	0.43
1:A:254:LEU:CB	1:A:256:LEU:CD1	2.96	0.43
1:C:141:SER:HG	1:C:144:GLN:HG3	1.84	0.43
1:D:22:ILE:HD11	1:D:25:LEU:HD23	1.99	0.43
1:B:202:GLN:NE2	4:B:386:HOH:O	2.52	0.43
1:A:60:ARG:HB3	1:A:65:SER:HB2	2.01	0.43
1:C:52:LYS:HE2	1:C:234:PHE:CZ	2.53	0.43
1:B:117:ARG:O	1:D:59:ASN:HA	2.19	0.42
1:B:58:ALA:HB2	1:B:94:HIS:CE1	2.55	0.42
1:B:180:GLY:HA3	4:B:371:HOH:O	2.18	0.42
1:D:22:ILE:HG23	1:D:34:THR:HG21	2.00	0.42
1:C:62:SER:O	1:C:63:ILE:C	2.57	0.42
1:C:135:LYS:HB3	1:C:165:ARG:HD2	2.02	0.42
1:C:201:LEU:O	1:C:202:GLN:C	2.58	0.42
1:C:255:HIS:CG	1:C:256:LEU:CD2	3.03	0.42
1:B:135:LYS:HB3	1:B:165:ARG:HD2	2.03	0.41
1:B:23:ASN:HB2	1:B:237:SER:HB2	2.03	0.40
1:A:59:ASN:C	1:A:59:ASN:OD1	2.59	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	246/280 (88%)	243 (99%)	2 (1%)	1 (0%)	34	24
1	B	250/280 (89%)	243 (97%)	7 (3%)	0	100	100
1	C	248/280 (89%)	242 (98%)	5 (2%)	1 (0%)	34	24
1	D	252/280 (90%)	249 (99%)	2 (1%)	1 (0%)	34	24
All	All	996/1120 (89%)	977 (98%)	16 (2%)	3 (0%)	41	31

All (3) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	229	ARG
1	C	229	ARG
1	A	229	ARG

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	206/235 (88%)	202 (98%)	4 (2%)	57	53
1	B	214/235 (91%)	208 (97%)	6 (3%)	43	36
1	C	212/235 (90%)	204 (96%)	8 (4%)	33	24
1	D	210/235 (89%)	202 (96%)	8 (4%)	33	24
All	All	842/940 (90%)	816 (97%)	26 (3%)	40	32

All (26) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	62	SER
1	A	135	LYS
1	A	154	GLU
1	A	165	ARG
1	B	40	GLU

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	71	LEU
1	B	202	GLN
1	B	214	ARG
1	B	237	SER
1	B	255	HIS
1	C	57	LYS
1	C	72	GLU
1	C	215	ARG
1	C	235	LEU
1	C	236	GLU
1	C	252	LEU
1	C	256	LEU
1	C	258	GLU
1	D	7	ASP
1	D	27	SER
1	D	52	LYS
1	D	69	VAL
1	D	76	LYS
1	D	145	MET
1	D	165	ARG
1	D	202	GLN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (12) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	6	ASN
1	A	217	GLN
1	B	23	ASN
1	B	94	HIS
1	B	110	GLN
1	C	23	ASN
1	C	100	GLN
1	C	173	ASN
1	C	202	GLN
1	D	23	ASN
1	D	64	HIS
1	D	199	HIS

### 5.3.3 RNA

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

Of 3 ligands modelled in this entry, 3 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	251/280 (89%)	0.72	25 (9%) <b>7</b> <b>8</b>	29, 44, 78, 106	2 (0%)
1	B	256/280 (91%)	0.42	15 (5%) <b>22</b> <b>25</b>	28, 39, 67, 101	0
1	C	253/280 (90%)	0.40	9 (3%) <b>42</b> <b>45</b>	28, 39, 61, 106	0
1	D	257/280 (91%)	0.91	41 (15%) <b>1</b> <b>2</b>	30, 51, 87, 108	2 (0%)
All	All	1017/1120 (90%)	0.61	90 (8%) <b>10</b> <b>11</b>	28, 41, 81, 108	4 (0%)

All (90) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	255	HIS	13.1
1	A	66	TYR	9.6
1	B	211	SER	6.6
1	D	239	PRO	6.4
1	A	202	GLN	6.1
1	A	253	PRO	6.0
1	C	254	LEU	5.8
1	B	206	ALA	5.1
1	B	212	GLY	5.0
1	A	254	LEU	4.6
1	D	33	GLN	4.5
1	A	64	HIS	4.3
1	D	66	TYR	4.3
1	D	252	LEU	4.3
1	A	69	VAL	4.3
1	C	213	GLY	4.1
1	B	277	LEU	4.1
1	B	256	LEU	4.1
1	B	255	HIS	4.1
1	C	253	PRO	3.9
1	A	27	SER	3.9

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	D	203	THR	3.9
1	D	36	ALA	3.8
1	C	277	LEU	3.8
1	A	67	ARG	3.7
1	A	216	ALA	3.7
1	C	252	LEU	3.6
1	D	71	LEU	3.6
1	D	30	SER	3.6
1	A	31	THR	3.6
1	A	32	LEU	3.5
1	D	75	LEU	3.5
1	D	81	VAL	3.5
1	A	68	GLY	3.4
1	D	254	LEU	3.4
1	D	76	LYS	3.3
1	A	22	ILE	3.3
1	A	24	VAL	3.3
1	D	79	GLU	3.3
1	D	63	ILE	3.2
1	A	71	LEU	3.1
1	D	253	PRO	3.1
1	A	29	ASP	3.1
1	D	32	LEU	3.1
1	A	30	SER	3.1
1	D	80	LYS	3.0
1	B	205	ASP	3.0
1	A	61	SER	2.9
1	D	267	LEU	2.9
1	D	29	ASP	2.8
1	D	82	LYS	2.8
1	C	251	ALA	2.7
1	D	213	GLY	2.7
1	B	270	LEU	2.7
1	D	62	SER	2.7
1	D	128	THR	2.7
1	D	78	PHE	2.6
1	D	105	VAL	2.6
1	B	87	ILE	2.6
1	D	276	ILE	2.6
1	A	76	LYS	2.5
1	C	199[A]	HIS	2.5
1	D	104	GLU	2.5

*Continued on next page...*



*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	D	83	ALA	2.5
1	A	201	LEU	2.5
1	B	36	ALA	2.5
1	D	28	LEU	2.5
1	D	72	GLU	2.4
1	D	214	ARG	2.4
1	D	215	ARG	2.4
1	D	100	GLN	2.4
1	D	202	GLN	2.3
1	A	70	GLY	2.3
1	B	7	ASP	2.3
1	B	43	ARG	2.3
1	D	224	ALA	2.3
1	A	72	GLU	2.3
1	A	65	SER	2.2
1	D	61	SER	2.2
1	B	202	GLN	2.2
1	D	27	SER	2.1
1	B	86	GLY	2.1
1	D	8	ILE	2.1
1	A	75	LEU	2.1
1	C	202	GLN	2.1
1	D	60	ARG	2.1
1	A	33	GLN	2.1
1	D	201	LEU	2.0
1	D	233	LEU	2.0
1	B	121	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,

median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	CL	C	302	1/1	0.92	0.15	70,70,70,70	0
2	NA	C	301	1/1	0.95	0.27	42,42,42,42	0
3	CL	D	301	1/1	0.95	0.09	65,65,65,65	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.