



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 15, 2023 – 07:50 PM JST

PDB ID : 6JVV  
Title : Crystal structure of maleylpyruvate hydrolase from *Sphingobium.sp* SYK-6  
Authors : Hong, H.; Kim, K.-J.  
Deposited on : 2019-04-17  
Resolution : 1.51 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.51 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	4009 (1.54-1.50)
Clashscore	141614	4249 (1.54-1.50)
Ramachandran outliers	138981	4148 (1.54-1.50)
Sidechain outliers	138945	4146 (1.54-1.50)
RSRZ outliers	127900	3943 (1.54-1.50)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	303	 2% 77% 9% 15%
1	B	303	 5% 78% 9% 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	GOL	A	404	-	-	X	-

## 2 Entry composition i

There are 5 unique types of molecules in this entry. The entry contains 4458 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called maleylpyruvate hydrolase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	259	1977	1259	342	365	11	0	2	0
1	B	266	2085	1322	369	383	11	0	8	0

There are 16 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	296	LEU	-	expression tag	UNP G2IPX5
A	297	GLU	-	expression tag	UNP G2IPX5
A	298	HIS	-	expression tag	UNP G2IPX5
A	299	HIS	-	expression tag	UNP G2IPX5
A	300	HIS	-	expression tag	UNP G2IPX5
A	301	HIS	-	expression tag	UNP G2IPX5
A	302	HIS	-	expression tag	UNP G2IPX5
A	303	HIS	-	expression tag	UNP G2IPX5
B	296	LEU	-	expression tag	UNP G2IPX5
B	297	GLU	-	expression tag	UNP G2IPX5
B	298	HIS	-	expression tag	UNP G2IPX5
B	299	HIS	-	expression tag	UNP G2IPX5
B	300	HIS	-	expression tag	UNP G2IPX5
B	301	HIS	-	expression tag	UNP G2IPX5
B	302	HIS	-	expression tag	UNP G2IPX5
B	303	HIS	-	expression tag	UNP G2IPX5

- Molecule 2 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	A	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0
2	B	1	Total C O 6 3 3	0	0

- Molecule 3 is DI(HYDROXYETHYL)ETHER (three-letter code: PEG) (formula: C<sub>4</sub>H<sub>10</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
3	B	1	Total	C	O	0	0
			7	4	3		

- Molecule 4 is SULFATE ION (three-letter code: SO<sub>4</sub>) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	B	1	Total	O	S	0	0
			5	4	1		

- Molecule 5 is water.

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
5	A	158	Total 158	O 158	0	0
5	B	172	Total 172	O 172	0	0





## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	67.09Å 74.52Å 67.97Å 90.00° 118.56° 90.00°	Depositor
Resolution (Å)	31.63 – 1.51 31.61 – 1.51	Depositor EDS
% Data completeness (in resolution range)	98.9 (31.63-1.51) 98.9 (31.61-1.51)	Depositor EDS
$R_{merge}$	0.05	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	3.00 (at 1.51Å)	Xtrriage
Refinement program	REFMAC 5.8.0238	Depositor
R, $R_{free}$	0.151 , 0.174 0.163 , 0.186	Depositor DCC
$R_{free}$ test set	4511 reflections (4.94%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.8	Xtrriage
Anisotropy	0.118	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.44 , 51.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.000 for -h-l,k,h 0.000 for l,k,-h-l 0.007 for h,-k,-h-l 0.006 for -h-l,-k,l 0.013 for l,-k,h	Xtrriage
$F_o, F_c$ correlation	0.97	EDS
Total number of atoms	4458	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	20.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 49.87 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 7.0130e-05. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4, PEG, GOL

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.80	1/2018 (0.0%)	1.00	4/2754 (0.1%)
1	B	0.80	0/2128	1.01	4/2900 (0.1%)
All	All	0.80	1/4146 (0.0%)	1.00	8/5654 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	A	113	SER	CB-OG	-6.43	1.33	1.42

All (8) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	B	5	ARG	NE-CZ-NH1	10.66	125.63	120.30
1	A	178	TYR	CB-CG-CD1	-8.25	116.05	121.00
1	B	10	ARG	NE-CZ-NH1	5.99	123.30	120.30
1	A	10	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	A	258	ARG	NE-CZ-NH1	5.52	123.06	120.30
1	B	258	ARG	NE-CZ-NH1	5.42	123.01	120.30
1	B	40	ARG	CG-CD-NE	-5.38	100.50	111.80
1	A	258	ARG	NE-CZ-NH2	-5.15	117.73	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1977	0	1994	19	1
1	B	2085	0	2087	27	1
2	A	24	0	32	7	0
2	B	30	0	40	3	0
3	B	7	0	10	2	0
4	B	5	0	0	0	0
5	A	158	0	0	7	0
5	B	172	0	0	5	0
All	All	4458	0	4163	49	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (49) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:A:404:GOL:H12	5:A:504:HOH:O	1.65	0.93
1:A:140:GLY:HA3	2:A:404:GOL:O1	1.74	0.88
2:A:404:GOL:C1	5:A:504:HOH:O	2.20	0.88
1:B:201[B]:ASP:HB3	1:B:203:LYS:NZ	1.93	0.83
2:A:404:GOL:H11	5:A:503:HOH:O	1.79	0.81
1:B:201[B]:ASP:HB3	1:B:203:LYS:HZ2	1.46	0.79
5:A:540:HOH:O	2:B:401:GOL:H32	1.86	0.76
1:B:49[A]:ARG:NH2	5:B:501:HOH:O	1.94	0.72
1:B:32:GLN:HE22	1:B:40:ARG:NH1	1.94	0.64
1:B:195:ASN:ND2	5:B:502:HOH:O	2.30	0.64
1:A:248:GLU:HA	2:A:402:GOL:H12	1.80	0.63
1:B:260:ARG:HH11	3:B:406:PEG:H31	1.64	0.61
1:A:250:VAL:HB	2:A:402:GOL:H11	1.84	0.59
1:A:124:VAL:CG1	1:A:127:ARG:HD2	2.32	0.59
1:B:84[A]:VAL:HG13	5:B:602:HOH:O	2.04	0.56
1:B:260:ARG:HH11	3:B:406:PEG:C3	2.18	0.56
1:B:130:HIS:CE1	1:B:167:ARG:HG3	2.40	0.56
1:A:124:VAL:HG12	1:A:127:ARG:HD2	1.89	0.55
1:A:156:PHE:HB2	2:A:404:GOL:O1	2.07	0.55
1:A:195:ASN:HB3	5:A:502:HOH:O	2.07	0.53
1:B:131:HIS:HE1	5:B:634:HOH:O	1.91	0.53
1:B:201[B]:ASP:CB	1:B:203:LYS:NZ	2.70	0.52
1:B:201[B]:ASP:HB3	1:B:203:LYS:HZ1	1.73	0.51
1:A:32:GLN:HE22	1:A:40:ARG:NH1	2.09	0.51

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:3:LEU:HD12	1:A:156:PHE:HZ	1.76	0.50
1:B:201[A]:ASP:HB3	1:B:203:LYS:NZ	2.27	0.49
1:B:124:VAL:HG11	1:B:127:ARG:HD2	1.94	0.49
1:B:201[B]:ASP:CB	1:B:203:LYS:HZ1	2.25	0.49
1:A:15:ILE:HD11	1:A:61:ARG:CD	2.43	0.49
1:B:186:PRO:HG2	1:B:187:TRP:CD2	2.49	0.48
1:B:107[A]:LYS:NZ	1:B:182:CYS:SG	2.82	0.48
1:B:176:LYS:NZ	5:B:509:HOH:O	2.47	0.48
1:A:56:LEU:HD23	1:A:59:LEU:HD12	1.97	0.46
1:A:2:ARG:HD3	5:A:593:HOH:O	2.14	0.46
1:A:131:HIS:HE1	5:A:609:HOH:O	1.98	0.45
1:B:201[A]:ASP:HB3	1:B:203:LYS:HZ2	1.81	0.45
1:A:186:PRO:HG2	1:A:187:TRP:CD2	2.52	0.44
1:A:15:ILE:HD11	1:A:61:ARG:HD3	2.00	0.44
1:B:151:TRP:CE2	1:B:152:LYS:HG3	2.53	0.44
1:B:78:LYS:HA	2:B:401:GOL:H2	2.00	0.44
1:A:56:LEU:HD23	1:A:56:LEU:HA	1.90	0.44
1:B:124:VAL:CG1	1:B:127:ARG:HD2	2.48	0.43
1:A:3:LEU:HD12	1:A:156:PHE:CZ	2.54	0.42
1:A:167:ARG:HD2	1:A:167:ARG:HA	1.81	0.41
1:B:259:ILE:O	1:B:270:ALA:HA	2.20	0.41
1:B:2:ARG:HD2	1:B:187:TRP:CE2	2.57	0.40
1:B:224:PRO:HG2	2:B:405:GOL:H32	2.03	0.40
1:A:172:ARG:HH11	1:B:102:GLN:HA	1.86	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:53:GLU:OE1	1:B:172:ARG:NH1[2_747]	2.18	0.02

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was

analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	255/303 (84%)	251 (98%)	3 (1%)	1 (0%)	34	13
1	B	268/303 (88%)	264 (98%)	1 (0%)	3 (1%)	14	2
All	All	523/606 (86%)	515 (98%)	4 (1%)	4 (1%)	25	4

All (4) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	91	ARG
1	A	84	VAL
1	B	84[A]	VAL
1	B	84[B]	VAL

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	210/247 (85%)	208 (99%)	2 (1%)	76	56
1	B	220/247 (89%)	219 (100%)	1 (0%)	88	78
All	All	430/494 (87%)	427 (99%)	3 (1%)	84	69

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	56	LEU
1	A	203	LYS
1	B	195	ASN

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (6) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	37	ASN
1	A	131	HIS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	B	37	ASN
1	B	66	GLN
1	B	131	HIS
1	B	195	ASN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

11 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z  > 2$	Counts	RMSZ	$\# Z  > 2$
2	GOL	B	403	-	5,5,5	0.12	0	5,5,5	0.35	0
2	GOL	A	401	-	5,5,5	0.15	0	5,5,5	0.34	0
3	PEG	B	406	-	6,6,6	0.40	0	5,5,5	0.38	0
2	GOL	B	404	-	5,5,5	0.16	0	5,5,5	0.55	0
4	SO4	B	407	-	4,4,4	0.14	0	6,6,6	0.34	0
2	GOL	A	403	-	5,5,5	0.17	0	5,5,5	0.26	0
2	GOL	A	402	-	5,5,5	0.19	0	5,5,5	0.22	0
2	GOL	B	401	-	5,5,5	0.09	0	5,5,5	0.78	0
2	GOL	B	402	-	5,5,5	0.11	0	5,5,5	0.19	0
2	GOL	A	404	-	5,5,5	0.20	0	5,5,5	0.84	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GOL	B	405	-	5,5,5	0.33	0	5,5,5	0.57	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GOL	B	403	-	-	4/4/4/4	-
2	GOL	A	401	-	-	0/4/4/4	-
3	PEG	B	406	-	-	4/4/4/4	-
2	GOL	B	404	-	-	2/4/4/4	-
2	GOL	A	403	-	-	0/4/4/4	-
2	GOL	A	402	-	-	4/4/4/4	-
2	GOL	B	401	-	-	4/4/4/4	-
2	GOL	B	402	-	-	4/4/4/4	-
2	GOL	A	404	-	-	4/4/4/4	-
2	GOL	B	405	-	-	0/4/4/4	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (26) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	402	GOL	O1-C1-C2-C3
2	A	404	GOL	O1-C1-C2-C3
2	B	401	GOL	O1-C1-C2-C3
2	B	402	GOL	C1-C2-C3-O3
2	B	403	GOL	C1-C2-C3-O3
2	B	404	GOL	O1-C1-C2-C3
2	B	401	GOL	O1-C1-C2-O2
3	B	406	PEG	O2-C3-C4-O4
2	A	402	GOL	C1-C2-C3-O3
2	B	401	GOL	C1-C2-C3-O3
2	B	402	GOL	O1-C1-C2-C3
2	B	403	GOL	O1-C1-C2-C3
2	A	402	GOL	O1-C1-C2-O2
2	B	402	GOL	O1-C1-C2-O2

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms
2	B	402	GOL	O2-C2-C3-O3
2	B	403	GOL	O2-C2-C3-O3
2	B	404	GOL	O1-C1-C2-O2
3	B	406	PEG	O1-C1-C2-O2
2	A	404	GOL	O2-C2-C3-O3
2	A	402	GOL	O2-C2-C3-O3
3	B	406	PEG	C4-C3-O2-C2
3	B	406	PEG	C1-C2-O2-C3
2	A	404	GOL	O1-C1-C2-O2
2	B	401	GOL	O2-C2-C3-O3
2	B	403	GOL	O1-C1-C2-O2
2	A	404	GOL	C1-C2-C3-O3

There are no ring outliers.

5 monomers are involved in 12 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	406	PEG	2	0
2	A	402	GOL	2	0
2	B	401	GOL	2	0
2	A	404	GOL	5	0
2	B	405	GOL	1	0

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	259/303 (85%)	-0.13	5 (1%) 66 71	11, 17, 34, 52	0
1	B	266/303 (87%)	0.01	14 (5%) 26 29	11, 17, 41, 82	0
All	All	525/606 (86%)	-0.06	19 (3%) 42 47	11, 17, 36, 82	0

All (19) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	169	ARG	8.0
1	B	91	ARG	6.2
1	A	168	GLY	4.7
1	B	89	HIS	4.4
1	B	92	GLU	4.0
1	B	124	VAL	4.0
1	B	195	ASN	3.9
1	B	90	GLY	3.9
1	B	168	GLY	3.8
1	A	125	PRO	3.5
1	B	126	GLY	3.4
1	B	102	GLN	3.2
1	B	125	PRO	3.1
1	B	127	ARG	2.9
1	A	195	ASN	2.9
1	B	65	ALA	2.3
1	A	167	ARG	2.3
1	B	172	ARG	2.2
1	A	127	ARG	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	GOL	B	405	6/6	0.66	0.22	39,44,51,52	0
2	GOL	A	402	6/6	0.69	0.32	34,42,48,57	0
3	PEG	B	406	7/7	0.77	0.15	31,37,44,45	0
2	GOL	B	404	6/6	0.86	0.14	29,44,45,52	0
2	GOL	A	404	6/6	0.87	0.30	42,45,50,68	0
2	GOL	B	402	6/6	0.89	0.15	38,48,53,55	0
2	GOL	B	401	6/6	0.91	0.12	17,22,26,27	0
2	GOL	B	403	6/6	0.92	0.26	25,45,51,62	0
2	GOL	A	401	6/6	0.92	0.08	20,27,27,29	0
2	GOL	A	403	6/6	0.93	0.15	30,40,45,45	0
4	SO4	B	407	5/5	0.94	0.20	34,37,40,41	0

### 6.5 Other polymers [i](#)

There are no such residues in this entry.