



# Full wwPDB X-ray Structure Validation Report ⓘ

Apr 30, 2024 – 07:38 PM JST

PDB ID : 8JYX  
Title : Crystal structure of the gasdermin-like protein RCD-1-1 from *Neurospora crassa*  
Authors : Li, Y.; Hou, Y.J.; Ding, J.  
Deposited on : 2023-07-04  
Resolution : 2.35 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36.2  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36.2


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.35 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1164 (2.36-2.36)
Clashscore	141614	1232 (2.36-2.36)
Ramachandran outliers	138981	1211 (2.36-2.36)
Sidechain outliers	138945	1212 (2.36-2.36)
RSRZ outliers	127900	1150 (2.36-2.36)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	635	 4% 81% 8% 11%
1	B	635	 4% 77% 11% 12%
2	C	2	 100%
2	D	2	 100%

## 2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 8949 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Maltodextrin-binding protein, Gasdermin-like protein rcd-1-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	565	4396	2829	736	812	19	0	0	0
1	B	558	4345	2796	727	803	19	0	0	0

There are 44 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-377	SER	-	expression tag	UNP A0A376KDN7
A	-376	GLY	-	expression tag	UNP A0A376KDN7
A	-375	ARG	-	expression tag	UNP A0A376KDN7
A	-374	PRO	-	expression tag	UNP A0A376KDN7
A	-373	MET	-	expression tag	UNP A0A376KDN7
A	-291	ALA	ASP	engineered mutation	UNP A0A376KDN7
A	-290	ALA	LYS	engineered mutation	UNP A0A376KDN7
A	-201	ALA	GLU	engineered mutation	UNP A0A376KDN7
A	-200	ALA	ASN	engineered mutation	UNP A0A376KDN7
A	-134	ALA	LYS	engineered mutation	UNP A0A376KDN7
A	-11	ALA	LYS	engineered mutation	UNP A0A376KDN7
A	-10	ALA	ASP	engineered mutation	UNP A0A376KDN7
A	-6	ASN	-	linker	UNP A0A376KDN7
A	-5	ALA	-	linker	UNP A0A376KDN7
A	-4	ALA	-	linker	UNP A0A376KDN7
A	-3	ARG	-	linker	UNP A0A376KDN7
A	-2	ALA	-	linker	UNP A0A376KDN7
A	-1	ALA	-	linker	UNP A0A376KDN7
A	0	ALA	-	linker	UNP A0A376KDN7
A	174	ALA	LYS	engineered mutation	UNP Q7SBA0
A	175	ALA	LYS	engineered mutation	UNP Q7SBA0
A	176	ALA	LYS	engineered mutation	UNP Q7SBA0
B	-377	SER	-	expression tag	UNP A0A376KDN7
B	-376	GLY	-	expression tag	UNP A0A376KDN7
B	-375	ARG	-	expression tag	UNP A0A376KDN7

*Continued on next page...*

Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
B	-374	PRO	-	expression tag	UNP A0A376KDN7
B	-373	MET	-	expression tag	UNP A0A376KDN7
B	-291	ALA	ASP	engineered mutation	UNP A0A376KDN7
B	-290	ALA	LYS	engineered mutation	UNP A0A376KDN7
B	-201	ALA	GLU	engineered mutation	UNP A0A376KDN7
B	-200	ALA	ASN	engineered mutation	UNP A0A376KDN7
B	-134	ALA	LYS	engineered mutation	UNP A0A376KDN7
B	-11	ALA	LYS	engineered mutation	UNP A0A376KDN7
B	-10	ALA	ASP	engineered mutation	UNP A0A376KDN7
B	-6	ASN	-	linker	UNP A0A376KDN7
B	-5	ALA	-	linker	UNP A0A376KDN7
B	-4	ALA	-	linker	UNP A0A376KDN7
B	-3	ARG	-	linker	UNP A0A376KDN7
B	-2	ALA	-	linker	UNP A0A376KDN7
B	-1	ALA	-	linker	UNP A0A376KDN7
B	0	ALA	-	linker	UNP A0A376KDN7
B	174	ALA	LYS	engineered mutation	UNP Q7SBA0
B	175	ALA	LYS	engineered mutation	UNP Q7SBA0
B	176	ALA	LYS	engineered mutation	UNP Q7SBA0

- Molecule 2 is an oligosaccharide called alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose.



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf	Trace
2	C	2	Total	C	O	0	0	0
			23	12	11			
2	D	2	Total	C	O	0	0	0
			23	12	11			

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	115	Total	O	0	0
			115	115		
3	B	47	Total	O	0	0
			47	47		



Chain C:  100%

GLC1  
GLC2

- Molecule 2: alpha-D-glucopyranose-(1-4)-alpha-D-glucopyranose

Chain D:  100%

GLC1  
GLC2

## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	81.14Å 91.91Å 105.87Å 90.00° 101.38° 90.00°	Depositor
Resolution (Å)	45.96 – 2.35 79.55 – 2.35	Depositor EDS
% Data completeness (in resolution range)	99.8 (45.96-2.35) 93.7 (79.55-2.35)	Depositor EDS
$R_{merge}$	0.09	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.20 (at 2.34Å)	Xtrriage
Refinement program	PHENIX 1.20.1_4487	Depositor
R, $R_{free}$	0.216 , 0.248 0.216 , 0.246	Depositor DCC
$R_{free}$ test set	2000 reflections (3.15%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	42.0	Xtrriage
Anisotropy	0.731	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.35 , 44.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.50$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	8949	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	58.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.37% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.25	0/4509	0.43	0/6116
1	B	0.24	0/4455	0.43	0/6040
All	All	0.25	0/8964	0.43	0/12156

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4396	0	4358	26	0
1	B	4345	0	4310	44	0
2	C	23	0	21	0	0
2	D	23	0	21	0	0
3	A	115	0	0	0	0
3	B	47	0	0	2	0
All	All	8949	0	8710	69	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (69) close contacts within the same asymmetric unit are listed below, sorted by their clash



magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:121:GLU:OE1	1:B:169:ARG:NH1	2.24	0.71
1:B:3:LYS:HD2	1:B:257:PHE:HA	1.73	0.70
1:A:231:LYS:NZ	1:A:233:GLY:O	2.29	0.66
1:A:22:ARG:NH2	1:A:58:MET:O	2.29	0.66
1:A:121:GLU:OE1	1:A:169:ARG:NH1	2.29	0.65
1:A:154:MET:SD	1:B:-48:GLN:NE2	2.70	0.64
1:B:30:ALA:HB3	1:B:221:VAL:HG21	1.83	0.60
1:B:36:ILE:HG21	1:B:42:LEU:HD12	1.83	0.60
1:A:-373:MET:HG3	1:A:-337:THR:HG21	1.84	0.59
1:B:52:GLU:OE1	1:B:138:ARG:NH1	2.36	0.59
1:B:-251:LEU:HD11	1:B:-238:LEU:HD21	1.85	0.58
1:B:254:LYS:NZ	3:B:302:HOH:O	2.33	0.58
1:B:69:PHE:HB2	1:B:118:ASP:HA	1.86	0.57
1:A:-251:LEU:HD21	1:A:-247:PRO:HD3	1.88	0.55
1:A:36:ILE:HG21	1:A:42:LEU:HD12	1.87	0.55
1:B:67:GLU:HG3	1:B:103:LEU:HB2	1.89	0.55
1:B:-141:TRP:HB2	1:B:-75:PRO:HG2	1.90	0.53
1:B:-80:VAL:HG12	1:B:-74:LEU:HD11	1.90	0.53
1:B:22:ARG:NH2	1:B:58:MET:O	2.43	0.52
1:A:129:ARG:NH2	1:A:238:TRP:O	2.43	0.51
1:A:139:ASP:HB3	1:A:143:ARG:NH1	2.26	0.51
1:B:-103:SER:O	1:B:-100:LYS:NZ	2.40	0.50
1:A:-214:PRO:HG3	1:A:-116:PRO:HA	1.93	0.50
1:B:119:ARG:HB3	1:B:169:ARG:HB2	1.93	0.50
1:B:-373:MET:HG3	1:B:-337:THR:HG21	1.94	0.49
1:A:-141:TRP:HB2	1:A:-75:PRO:HG2	1.95	0.48
1:B:-284:LEU:HB2	1:B:-279:TRP:NE1	2.27	0.48
1:B:9:ASP:N	1:B:9:ASP:OD1	2.46	0.48
1:A:113:ASN:HB3	1:A:212:TRP:CE2	2.49	0.48
1:B:-305:GLY:HA3	1:B:-41:ASN:O	2.14	0.48
1:B:-144:PRO:HA	1:B:-141:TRP:CE2	2.48	0.48
1:B:58:MET:HE1	1:B:128:THR:H	1.78	0.48
1:A:-280:THR:HB	1:A:-266:PRO:HB2	1.95	0.47
1:A:-27:ALA:HB2	1:A:-9:ALA:HB2	1.96	0.47
1:A:-284:LEU:HB2	1:A:-279:TRP:NE1	2.29	0.47
1:B:-102:PRO:HB3	1:B:57:THR:HB	1.97	0.47
1:B:-105:ALA:HB1	1:B:61:HIS:NE2	2.29	0.47
1:B:-195:ILE:HD11	1:B:101:VAL:HG11	1.96	0.47
1:A:-305:GLY:HA3	1:A:-41:ASN:O	2.14	0.47
1:B:-263:VAL:HG22	1:B:-112:VAL:HG22	1.96	0.46
1:B:22:ARG:NH2	1:B:59:ASP:HA	2.30	0.46

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:-364:ILE:HG21	1:B:-353:LEU:HD21	1.98	0.46
1:A:214:THR:O	1:A:217:THR:HG22	2.16	0.45
1:A:-144:PRO:HA	1:A:-141:TRP:CE2	2.52	0.45
1:B:159:VAL:HG22	1:B:229:ILE:HB	2.00	0.44
1:B:65:ILE:HD13	1:B:121:GLU:HG2	1.98	0.44
1:B:-363:TRP:CD2	1:B:-316:PRO:HG3	2.53	0.44
1:A:-311:TRP:CD1	1:A:-307:ARG:HG3	2.52	0.43
1:B:-361:ASN:ND2	3:B:305:HOH:O	2.51	0.43
1:B:-140:SER:HB2	1:B:-75:PRO:HD3	2.00	0.43
1:B:34:HIS:CD2	1:B:51:ILE:HD11	2.54	0.43
1:B:-216:THR:HG21	1:B:-26:VAL:HG11	2.01	0.43
1:A:251:ARG:HD2	1:A:251:ARG:HA	1.89	0.42
1:B:122:ARG:HG2	1:B:124:ILE:HG12	2.01	0.42
1:A:71:TRP:CE3	1:A:115:TRP:HA	2.54	0.42
1:B:4:CYS:HB3	1:B:244:PHE:CD1	2.54	0.42
1:B:-140:SER:HB3	1:B:-76:LYS:HD3	2.01	0.42
1:B:254:LYS:HD2	1:B:254:LYS:H	1.85	0.42
1:B:144:TRP:CZ3	1:B:159:VAL:HG12	2.55	0.42
1:A:158:GLU:HB2	1:A:230:THR:HG22	2.02	0.41
1:A:-102:PRO:HB3	1:A:57:THR:HB	2.02	0.41
1:A:228:LYS:HG2	1:A:230:THR:HG23	2.03	0.41
1:B:-280:THR:HB	1:B:-266:PRO:HB2	2.01	0.41
1:B:-276:VAL:HG21	1:B:-266:PRO:HD3	2.01	0.41
1:A:122:ARG:HG2	1:A:124:ILE:HG12	2.03	0.41
1:B:-219:PRO:HA	1:B:-216:THR:HG22	2.03	0.41
1:A:34:HIS:CD2	1:A:51:ILE:HD11	2.55	0.40
1:B:25:HIS:HA	1:B:49:LYS:HA	2.03	0.40
1:B:61:HIS:O	1:B:123:TYR:HA	2.21	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	559/635 (88%)	540 (97%)	19 (3%)	0	100	100
1	B	550/635 (87%)	536 (98%)	14 (2%)	0	100	100
All	All	1109/1270 (87%)	1076 (97%)	33 (3%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	451/505 (89%)	445 (99%)	6 (1%)	69	80
1	B	447/505 (88%)	445 (100%)	2 (0%)	91	95
All	All	898/1010 (89%)	890 (99%)	8 (1%)	78	87

All (8) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	-203	LYS
1	A	-127	VAL
1	A	-115	PHE
1	A	156	ARG
1	A	157	TRP
1	A	217	THR
1	B	-115	PHE
1	B	9	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

4 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	GLC	C	1	2	12,12,12	0.19	0	17,17,17	0.28	0
2	GLC	C	2	2	11,11,12	0.40	0	15,15,17	0.52	0
2	GLC	D	1	2	12,12,12	0.20	0	17,17,17	0.33	0
2	GLC	D	2	2	11,11,12	0.36	0	15,15,17	0.50	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	GLC	C	1	2	-	0/2/22/22	0/1/1/1
2	GLC	C	2	2	-	0/2/19/22	0/1/1/1
2	GLC	D	1	2	-	0/2/22/22	0/1/1/1
2	GLC	D	2	2	-	0/2/19/22	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.


There are no chirality outliers.


There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.





## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	565/635 (88%)	0.63	25 (4%) 34 46	38, 51, 78, 123	0
1	B	558/635 (87%)	0.74	28 (5%) 28 41	43, 58, 88, 122	0
All	All	1123/1270 (88%)	0.68	53 (4%) 31 44	38, 55, 85, 123	0

All (53) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	0	ALA	7.5
1	B	101	VAL	7.0
1	B	-371	ILE	6.6
1	A	101	VAL	6.3
1	B	250	PHE	5.4
1	A	1	MET	4.8
1	A	250	PHE	4.3
1	B	251	ARG	4.3
1	B	10	ASN	3.9
1	B	70	LYS	3.8
1	B	215	SER	3.7
1	B	1	MET	3.7
1	B	-201	ALA	3.5
1	B	217	THR	3.5
1	B	-372	LYS	3.3
1	A	-201	ALA	3.2
1	A	10	ASN	3.1
1	B	-134	ALA	3.0
1	B	-200	ALA	2.9
1	A	251	ARG	2.9
1	B	-4	ALA	2.8
1	A	-373	MET	2.8
1	B	100	ASN	2.8
1	B	-202	TYR	2.7

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	157	TRP	2.7
1	A	124	ILE	2.7
1	A	-371	ILE	2.7
1	A	71	TRP	2.6
1	A	216	GLN	2.6
1	A	42	LEU	2.6
1	B	60	ILE	2.4
1	A	-232	ALA	2.4
1	B	216	GLN	2.4
1	A	-198	LYS	2.4
1	B	12	HIS	2.4
1	A	214	THR	2.3
1	B	-306	PHE	2.3
1	A	213	GLY	2.3
1	A	-372	LYS	2.2
1	A	-306	PHE	2.2
1	A	-72	ALA	2.2
1	B	218	GLY	2.2
1	A	-2	ALA	2.1
1	B	-195	ILE	2.1
1	B	40	ALA	2.1
1	B	-367	LYS	2.1
1	A	8	LEU	2.1
1	B	42	LEU	2.1
1	B	-373	MET	2.1
1	A	-56	ILE	2.1
1	A	159	VAL	2.1
1	B	69	PHE	2.1
1	B	-252	LEU	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.


## 6.3 Carbohydrates [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.



Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors( $\text{\AA}^2$ )	Q<0.9
2	GLC	D	1	12/12	0.93	0.15	45,52,56,61	0
2	GLC	C	1	12/12	0.94	0.17	42,47,49,56	0
2	GLC	D	2	11/12	0.95	0.16	41,46,53,54	0
2	GLC	C	2	11/12	0.97	0.16	40,42,42,47	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.





## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.