



# Full wwPDB X-ray Structure Validation Report i

Sep 13, 2021 – 03:17 PM EDT

PDB ID : 7JZI  
Title : Crystal structure of LAIR1 ectodomain (from MGD21) in complex with Plasmodium RIFIN (PF3D7\_1040300) V2 domain  
Authors : Xu, K.; Kwong, P.D.  
Deposited on : 2020-09-02  
Resolution : 2.71 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)  
A user guide is available at  
<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>  
with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) ①) were used in the production of this report:

MolProbity	:	4.02b-467
Xtriage (Phenix)	:	1.13
EDS	:	2.23.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*


The reported resolution of this entry is 2.71 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 4 unique types of molecules in this entry. The entry contains 3430 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called LAIR1 ectodomain from antibody MGD21.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	98	780	488	137	153	2	0	0	0
1	C	98	780	488	137	153	2	0	0	0

- Molecule 2 is a protein called Rifin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	122	891	561	151	174	5	0	0	0
2	D	132	966	608	162	191	5	0	0	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
B	232	GLN	ASN	conflict	UNP A0A2I0BRG0
D	232	GLN	ASN	conflict	UNP A0A2I0BRG0

- Molecule 3 is PLATINUM (II) ION (three-letter code: PT) (formula: Pt).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	B	1	Total	Pt	0	0
3	D	1	Total	Pt	0	0

- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	1	Total	O	0	0

*Continued on next page...*


*Continued from previous page...*

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	B	1	Total O 1 1	0	0
4	C	8	Total O 8 8	0	0
4	D	1	Total O 1 1	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: LAIR1 ectodomain from antibody MGD21




- Molecule 1: LAIR1 ectodomain from antibody MGD21



- Molecule 2: Rifin



- Molecule 2: Rifin



## 4 Data and refinement statistics i

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	144.41Å 72.18Å 57.92Å 90.00° 91.08° 90.00°	Depositor
Resolution (Å)	44.76 – 2.71 44.76 – 2.71	Depositor EDS
% Data completeness (in resolution range)	97.6 (44.76-2.71) 90.8 (44.76-2.71)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	2.39 (at 2.73Å)	Xtriage
Refinement program	PHENIX 1.12_2829	Depositor
$R$ , $R_{free}$	0.240 , 0.281 0.247 , 0.270	Depositor DCC
$R_{free}$ test set	1613 reflections (10.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	86.3	Xtriage
Anisotropy	0.573	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.29 , 64.3	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtriage
Estimated twinning fraction	0.298 for -h,-k,l	Xtriage
$F_o, F_c$ correlation	0.94	EDS
Total number of atoms	3430	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	111.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.02% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [\(i\)](#)

### 5.1 Standard geometry [\(i\)](#)

Bond lengths and bond angles in the following residue types are not validated in this section: PT

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.38	0/797	0.70	0/1081
1	C	0.48	1/797 (0.1%)	0.72	0/1081
2	B	0.42	0/899	0.81	2/1212 (0.2%)
2	D	0.42	1/974 (0.1%)	0.73	1/1314 (0.1%)
All	All	0.43	2/3467 (0.1%)	0.74	3/4688 (0.1%)

All (2) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	C	148	ARG	CB-CG	-5.63	1.37	1.52
2	D	297	GLU	CB-CG	5.52	1.62	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	297	GLU	CA-CB-CG	7.50	129.89	113.40
2	B	214	LEU	CA-CB-CG	6.36	129.94	115.30
2	B	220	ASP	CB-CG-OD2	-5.57	113.29	118.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	780	0	755	24	0
1	C	780	0	755	26	0
2	B	891	0	928	35	0
2	D	966	0	1005	37	0
3	B	1	0	0	0	0
3	D	1	0	0	0	0
4	A	1	0	0	0	0
4	B	1	0	0	0	0
4	C	8	0	0	1	0
4	D	1	0	0	0	0
All	All	3430	0	3443	115	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (115) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:210:VAL:HG11	2:D:239:VAL:HG11	1.24	1.12
2:D:209:ILE:HD11	2:D:299:ALA:HA	1.34	1.03
1:A:129:VAL:HG12	1:A:172:ILE:HD11	1.50	0.92
2:B:209:ILE:HD11	2:B:299:ALA:HA	1.54	0.90
2:D:221:THR:HG21	2:D:263:PRO:HG2	1.57	0.85
2:B:219:VAL:HG12	2:B:227:PHE:HB2	1.62	0.81
1:C:129:VAL:HG12	1:C:172:ILE:HD11	1.62	0.79
1:C:161:THR:OG1	1:C:165:GLU:HB3	1.84	0.76
2:B:205:HIS:HD2	2:B:306:LEU:HD12	1.50	0.76
2:B:209:ILE:HD12	2:B:298:LYS:HG3	1.68	0.75
1:A:143:ARG:HD2	1:A:186:ILE:HD11	1.69	0.74
1:C:197:SER:N	4:C:301:HOH:O	2.18	0.73
2:B:301:GLN:HG2	2:B:301:GLN:O	1.89	0.72
1:A:139:VAL:HA	1:A:189:LYS:HB2	1.70	0.71
2:D:188:GLU:HB3	2:D:189:LEU:HD22	1.73	0.71
2:B:285:LYS:NZ	2:B:289:THR:HG21	2.07	0.70
2:D:209:ILE:HD12	2:D:298:LYS:HG3	1.75	0.67
2:B:210:VAL:O	2:B:214:LEU:HD22	1.95	0.67
2:B:208:LYS:O	2:B:211:ILE:HG13	1.95	0.66
1:C:109:ASP:O	1:C:110:LEU:HD23	1.97	0.64
1:C:176:ASN:CB	2:D:259:ILE:HD12	2.29	0.63
2:D:294:GLN:HA	2:D:297:GLU:HG3	1.81	0.62
1:A:157:ASP:OD2	1:A:157:ASP:O	2.17	0.62
2:D:217:LEU:HD21	2:D:291:VAL:HG11	1.82	0.61

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:222:LEU:HD22	2:B:249:GLU:HG2	1.83	0.61
2:D:222:LEU:HD23	2:D:249:GLU:HG2	1.81	0.61
2:B:288:ILE:O	2:B:292:LEU:HD12	2.01	0.60
2:D:209:ILE:HD11	2:D:299:ALA:CA	2.21	0.59
2:B:249:GLU:O	2:B:253:VAL:HG22	2.05	0.57
1:C:176:ASN:HB2	2:D:259:ILE:HD12	1.86	0.57
2:D:268:ARG:HA	2:D:271:LEU:HD12	1.86	0.57
2:D:294:GLN:O	2:D:297:GLU:HB2	2.04	0.57
1:C:180:ALA:HB2	1:C:204:VAL:HG13	1.86	0.56
1:C:161:THR:OG1	1:C:165:GLU:HG2	2.04	0.56
2:D:264:ILE:O	2:D:267:VAL:HG22	2.04	0.56
2:D:236:TYR:O	2:D:239:VAL:HG13	2.06	0.55
1:A:172:ILE:HD12	1:A:172:ILE:O	2.07	0.55
2:D:244:ILE:O	2:D:248:THR:HG23	2.06	0.55
2:D:207:MET:O	2:D:211:ILE:HG13	2.07	0.55
1:C:172:ILE:HD12	1:C:172:ILE:O	2.08	0.54
2:B:296:LEU:O	2:B:300:THR:HG23	2.08	0.53
1:A:186:ILE:CG2	1:A:196:GLN:HG3	2.39	0.53
1:C:161:THR:OG1	1:C:165:GLU:CB	2.56	0.52
1:A:130:THR:CG2	1:A:171:ARG:HB2	2.40	0.52
1:C:121:THR:HG22	1:C:200:LEU:HD11	1.92	0.52
2:B:295:ALA:O	2:B:298:LYS:HB3	2.10	0.52
2:B:240:THR:O	2:B:243:ALA:HB3	2.10	0.52
2:D:214:LEU:HD21	2:D:242:LEU:HD13	1.90	0.52
2:D:201:ALA:O	2:D:204:ILE:HG13	2.10	0.52
1:C:190:SER:O	1:C:191:ARG:HB2	2.10	0.51
2:D:296:LEU:O	2:D:300:THR:HG23	2.11	0.51
2:B:285:LYS:HZ2	2:B:289:THR:HG21	1.76	0.51
2:D:205:HIS:CE1	2:D:302:SER:HB3	2.45	0.51
2:D:224:SER:OG	2:D:226:ILE:HG13	2.11	0.51
1:A:172:ILE:CG2	2:B:257:THR:HG23	2.41	0.51
2:B:205:HIS:CD2	2:B:306:LEU:HD12	2.39	0.50
2:D:281:MET:SD	2:D:283:LYS:NZ	2.81	0.50
2:B:217:LEU:HD11	2:B:291:VAL:HG11	1.94	0.50
1:C:127:SER:O	1:C:174:SER:HA	2.11	0.50
1:C:128:HIS:HD1	1:C:128:HIS:C	2.15	0.50
2:B:205:HIS:HD2	2:B:306:LEU:CD1	2.22	0.49
2:D:267:VAL:HA	2:D:270:LYS:HD2	1.93	0.49
2:B:221:THR:HG23	2:B:262:HIS:HE1	1.78	0.49
2:B:212:GLU:HA	2:B:215:LYS:HE2	1.95	0.49
2:B:230:PHE:CZ	2:B:236:TYR:HA	2.48	0.49

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:D:239:VAL:HA	2:D:242:LEU:HD12	1.96	0.48
2:B:242:LEU:O	2:B:246:ILE:HG12	2.13	0.48
2:B:236:TYR:O	2:B:239:VAL:HG13	2.13	0.48
1:A:123:ILE:HG13	1:A:129:VAL:HB	1.96	0.47
1:A:127:SER:O	1:A:174:SER:HA	2.14	0.47
1:A:177:ALA:HA	1:A:204:VAL:HG21	1.95	0.47
1:A:172:ILE:HG22	2:B:257:THR:HG23	1.97	0.47
2:B:217:LEU:HA	2:B:267:VAL:HG21	1.95	0.47
1:A:126:GLY:O	1:A:174:SER:HB2	2.14	0.47
1:A:139:VAL:HG21	1:A:142:PHE:CE2	2.49	0.47
2:B:230:PHE:HZ	2:B:236:TYR:HA	1.79	0.47
2:D:294:GLN:HA	2:D:297:GLU:CG	2.43	0.46
1:C:130:THR:HG22	1:C:171:ARG:CB	2.45	0.46
1:C:145:GLU:OE2	1:C:186:ILE:HD11	2.16	0.46
2:B:244:ILE:O	2:B:248:THR:HG23	2.16	0.46
2:D:253:VAL:HG11	2:D:262:HIS:CD2	2.51	0.46
2:B:247:ASP:OD2	2:B:285:LYS:HB2	2.16	0.45
2:B:218:LYS:HB3	2:B:221:THR:HG22	1.97	0.45
1:C:129:VAL:HG12	1:C:172:ILE:CD1	2.39	0.45
1:C:136:PRO:O	1:C:166:SER:HB3	2.17	0.45
1:A:147:GLU:OE2	1:C:199:TYR:N	2.40	0.45
1:A:130:THR:HG22	1:A:171:ARG:HB2	1.98	0.45
1:A:115:ILE:HD11	1:A:131:PHE:HB3	1.99	0.45
1:A:172:ILE:HD13	1:A:175:VAL:HG12	2.00	0.44
2:D:303:ALA:O	2:D:307:SER:HB2	2.16	0.44
2:D:226:ILE:HG22	2:D:242:LEU:HD23	1.99	0.44
2:D:219:VAL:O	2:D:222:LEU:HD12	2.18	0.43
2:B:273:VAL:HA	2:B:282:VAL:HG13	2.01	0.43
1:A:129:VAL:HG12	1:A:172:ILE:CD1	2.36	0.43
1:A:186:ILE:HG22	1:A:196:GLN:HA	1.99	0.43
1:A:134:ARG:NH1	1:A:167:GLU:OE1	2.51	0.43
1:A:139:VAL:HG23	1:A:158:VAL:HG11	2.01	0.43
1:C:130:THR:HG22	1:C:171:ARG:HB2	2.01	0.42
2:B:300:THR:O	2:B:304:GLU:HG3	2.20	0.42
2:D:227:PHE:O	2:D:231:VAL:HG13	2.18	0.42
2:D:247:ASP:OD2	2:D:284:GLN:NE2	2.52	0.42
2:D:284:GLN:O	2:D:288:ILE:HG13	2.19	0.42
1:A:147:GLU:HG3	1:C:199:TYR:HD1	1.84	0.42
2:D:236:TYR:CE2	2:D:237:THR:HG22	2.55	0.42
2:B:208:LYS:O	2:B:212:GLU:HG2	2.20	0.41
1:C:118:GLU:HA	1:C:119:PRO:HA	1.89	0.41

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:131:PHE:HZ	1:C:202:LEU:HD22	1.85	0.41
1:C:128:HIS:C	1:C:128:HIS:ND1	2.74	0.41
2:B:235:HIS:O	2:B:235:HIS:CD2	2.73	0.41
1:C:184:ARG:HD3	1:C:199:TYR:CE2	2.56	0.40
1:C:152:TYR:HA	2:D:256:ALA:HB2	2.03	0.40
2:B:285:LYS:HZ3	2:B:289:THR:HG21	1.82	0.40
1:A:117:ALA:HB2	1:A:200:LEU:HD21	2.04	0.40
2:D:219:VAL:HG12	2:D:227:PHE:HB2	2.04	0.40
2:D:269:GLU:HA	2:D:274:ILE:CD1	2.52	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles [\(i\)](#)

#### 5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	96/98 (98%)	94 (98%)	2 (2%)	0	100 100
1	C	96/98 (98%)	92 (96%)	4 (4%)	0	100 100
2	B	120/145 (83%)	118 (98%)	2 (2%)	0	100 100
2	D	130/145 (90%)	128 (98%)	2 (2%)	0	100 100
All	All	442/486 (91%)	432 (98%)	10 (2%)	0	100 100

There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	88/88 (100%)	85 (97%)	3 (3%)	37 66
1	C	88/88 (100%)	85 (97%)	3 (3%)	37 66
2	B	94/111 (85%)	93 (99%)	1 (1%)	73 90
2	D	102/111 (92%)	100 (98%)	2 (2%)	55 81
All	All	372/398 (94%)	363 (98%)	9 (2%)	49 77

All (9) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	130	THR
1	A	142	PHE
1	A	162	SER
2	B	267	VAL
1	C	128	HIS
1	C	132	VAL
1	C	148	ARG
2	D	188	GLU
2	D	285	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
2	B	235	HIS
2	B	251	ASN
2	B	284	GLN
2	D	205	HIS
2	D	251	ASN

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [\(i\)](#)

Of 2 ligands modelled in this entry, 2 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	98/98 (100%)	0.32	6 (6%) 21 20	30, 90, 118, 127	0
1	C	98/98 (100%)	0.46	4 (4%) 37 36	71, 94, 126, 155	0
2	B	122/145 (84%)	1.13	24 (19%) 1 0	81, 124, 184, 201	0
2	D	132/145 (91%)	0.41	6 (4%) 33 31	80, 122, 153, 167	0
All	All	450/486 (92%)	0.60	40 (8%) 9 7	30, 105, 167, 201	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
2	B	196	ALA	10.8
2	B	206	GLY	10.6
2	B	310	THR	7.3
2	B	202	GLY	7.0
2	B	296	LEU	5.5
2	B	195	ALA	5.1
1	C	108	THR	5.0
2	B	193	ALA	4.6
2	B	205	HIS	4.6
1	A	108	THR	3.9
2	B	311	ALA	3.8
1	A	175	VAL	3.7
1	C	204	VAL	3.7
1	A	125	LEU	3.6
2	B	192	LYS	3.4
2	B	194	GLY	3.1
2	B	303	ALA	3.1
2	B	199	LEU	3.1
2	D	188	GLU	3.0
1	A	139	VAL	3.0
2	B	200	LYS	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	292	LEU	2.9
2	B	232	GLN	2.9
2	D	306	LEU	2.9
2	B	231	VAL	2.8
1	C	125	LEU	2.7
2	D	229	SER	2.6
2	D	299	ALA	2.5
1	C	191	ARG	2.5
1	A	120	GLY	2.4
2	B	203	ASP	2.4
2	B	207	MET	2.4
2	D	232	GLN	2.3
2	B	308	GLU	2.3
2	B	219	VAL	2.3
2	B	309	THR	2.2
2	D	318	LEU	2.2
1	A	109	ASP	2.2
2	B	297	GLU	2.1
2	B	210	VAL	2.1

## 6.2 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [\(i\)](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	PT	D	401	1/1	0.71	0.20	272,272,272,272	0
3	PT	B	401	1/1	0.93	0.32	280,280,280,280	0

## 6.5 Other polymers [\(i\)](#)

There are no such residues in this entry.