



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 8, 2023 – 11:08 PM EDT

PDB ID : 7K2B  
Title : Kelch domain of human KEAP1 bound to Nrf2 peptide, ADEETGEFA  
Authors : Muellers, S.N.; Allen, K.N.  
Deposited on : 2020-09-08  
Resolution : 2.31 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtrriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.31 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5974 (2.34-2.30)
Clashscore	141614	6604 (2.34-2.30)
Ramachandran outliers	138981	6523 (2.34-2.30)
Sidechain outliers	138945	6523 (2.34-2.30)
RSRZ outliers	127900	5855 (2.34-2.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	301	
1	B	301	
2	P	11	

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 4597 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Kelch-like ECH-associated protein 1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	286	2192	1365	395	417	15	0	0	0
1	B	283	2159	1346	390	408	15	0	0	0

There are 4 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	540	ALA	GLU	engineered mutation	UNP Q14145
A	542	ALA	GLU	engineered mutation	UNP Q14145
B	540	ALA	GLU	engineered mutation	UNP Q14145
B	542	ALA	GLU	engineered mutation	UNP Q14145

- Molecule 2 is a protein called ACE-ALA-ASP-GLU-GLU-THR-GLY-GLU-PHE-ALA-NH2.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
2	P	11	71	42	10	19	0	0	1


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	129	Total	O	0	0
			129	129		
3	B	42	Total	O	0	0
			42	42		
3	P	4	Total	O	0	0
			4	4		

### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Kelch-like ECH-associated protein 1

Chain A:  78% 16% • 5%



- Molecule 1: Kelch-like ECH-associated protein 1

Chain B:  24% 67% 26% • 6%



- Molecule 2: ACE-ALA-ASP-GLU-GLU-THR-GLY-GLU-PHE-ALA-NH2

Chain P:  64% 27% 9%



## 4 Data and refinement statistics

Property	Value	Source
Space group	C 1 2 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	161.72Å 68.75Å 77.62Å 90.00° 117.60° 90.00°	Depositor
Resolution (Å)	39.23 – 2.31 39.23 – 2.31	Depositor EDS
% Data completeness (in resolution range)	98.0 (39.23-2.31) 98.0 (39.23-2.31)	Depositor EDS
$R_{merge}$	0.07	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.75 (at 2.31Å)	Xtrriage
Refinement program	PHENIX 1.17.1_3660	Depositor
R, $R_{free}$	0.215 , 0.258 0.215 , 0.258	Depositor DCC
$R_{free}$ test set	1994 reflections (6.12%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	33.6	Xtrriage
Anisotropy	0.353	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 43.4	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.49$ , $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	0.020 for -h-2*1,-k,l	Xtrriage
$F_o, F_c$ correlation	0.93	EDS
Total number of atoms	4597	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	46.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.18% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACE, NH2

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# $ Z  > 5$	RMSZ	# $ Z  > 5$
1	A	0.48	0/2246	0.65	0/3061
1	B	0.39	0/2212	0.64	1/3015 (0.0%)
2	P	1.76	3/68 (4.4%)	1.57	2/91 (2.2%)
All	All	0.49	3/4526 (0.1%)	0.67	3/6167 (0.0%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

All (3) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	P	85	ALA	C-O	9.72	1.41	1.23
2	P	84	PHE	C-N	-6.83	1.18	1.34
2	P	85	ALA	CA-C	5.84	1.68	1.52

All (3) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	P	85	ALA	CB-CA-C	-10.09	94.97	110.10
2	P	85	ALA	N-CA-C	6.67	129.01	111.00
1	B	353	LEU	CA-CB-CG	5.05	126.91	115.30

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	609	THR	Peptide

## 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2192	0	2083	37	0
1	B	2159	0	2052	62	0
2	P	71	0	53	3	0
3	A	129	0	0	3	0
3	B	42	0	0	1	0
3	P	4	0	0	0	0
All	All	4597	0	4188	99	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 12.

All (99) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:483:ARG:NH1	3:A:701:HOH:O	1.74	1.14
1:B:380:ARG:NH1	2:P:83:GLU:OE1	1.91	1.03
1:B:328:ILE:HD11	1:B:562:HIS:HB2	1.57	0.86
1:B:349:ASP:HB2	1:B:351:THR:HG22	1.62	0.82
1:B:581:VAL:HB	1:B:595:THR:HB	1.63	0.81
1:B:421:ILE:HG12	1:B:467:VAL:HG11	1.64	0.79
1:B:466:ALA:HB1	1:B:514:VAL:HG23	1.65	0.78
1:A:483:ARG:HD2	3:A:701:HOH:O	1.82	0.78
1:B:415:ARG:NH2	2:P:80:GLU:OE1	2.18	0.71
1:B:562:HIS:HB3	1:B:567:TYR:HE1	1.55	0.70
1:B:441:GLU:HB3	1:B:452:LEU:HD23	1.74	0.68
1:B:447:ARG:HD2	1:B:449:GLU:HB2	1.76	0.66
1:A:550:MET:CE	1:A:568:VAL:HG11	2.25	0.66
1:A:550:MET:HE2	1:A:568:VAL:HG11	1.79	0.64
1:B:416:ILE:HD11	1:B:427:ALA:HB1	1.80	0.63
1:B:562:HIS:HB3	1:B:567:TYR:CE1	2.35	0.62
1:A:430:GLY:C	1:A:461:ILE:HG22	2.20	0.62

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:388:THR:HG22	1:A:389:ASP:O	2.00	0.61
1:B:421:ILE:HG12	1:B:467:VAL:CG1	2.29	0.61
1:A:515:LEU:HD22	1:A:566:ILE:HG13	1.83	0.61
1:B:554:ARG:HG3	1:B:557:LEU:HD22	1.84	0.60
1:B:410:SER:OG	1:B:441:GLU:OE2	2.19	0.60
1:A:430:GLY:O	1:A:461:ILE:HG22	2.03	0.58
1:A:522:ALA:HB1	1:A:550:MET:HE3	1.86	0.58
1:B:561:VAL:HG12	1:B:566:ILE:HG12	1.85	0.58
1:B:468:LEU:HD12	1:B:469:ASN:H	1.69	0.58
1:B:355:LEU:HD13	1:B:396:TYR:OH	2.04	0.58
1:B:559:ILE:HG13	1:B:568:VAL:HG12	1.88	0.56
1:A:431:SER:HB3	1:A:461:ILE:HG21	1.87	0.55
1:B:424:HIS:CD2	1:B:444:GLU:HG3	2.42	0.55
1:B:515:LEU:HD22	1:B:566:ILE:HG13	1.88	0.55
1:B:443:TYR:HB2	1:B:450:TRP:CE2	2.42	0.53
1:B:380:ARG:NH1	1:B:382:ASN:HD22	2.06	0.53
1:B:347:PRO:HG2	1:B:562:HIS:ND1	2.24	0.53
1:A:490:TYR:CE2	1:A:492:PRO:HA	2.44	0.53
1:A:554:ARG:HH22	1:A:580:SER:HG	1.54	0.52
1:B:483:ARG:HG2	1:B:508:SER:HB2	1.91	0.52
1:B:400:THR:OG1	1:B:402:GLN:HG2	2.10	0.52
1:A:337:GLN:HG3	1:A:384:PRO:HD3	1.91	0.52
1:B:397:ASN:HB3	1:B:400:THR:OG1	2.10	0.52
1:B:532:ASN:HB2	1:B:550:MET:O	2.10	0.52
1:B:342:LEU:HD22	1:B:403:TRP:CZ2	2.45	0.52
1:B:366:ALA:HB1	1:B:418:VAL:HG22	1.92	0.51
1:B:494:ARG:NH1	1:B:496:GLU:OE2	2.36	0.51
1:B:328:ILE:CD1	1:B:562:HIS:HB2	2.37	0.51
1:B:594:VAL:HG12	1:B:595:THR:OG1	2.11	0.51
1:A:522:ALA:HB1	1:A:550:MET:CE	2.40	0.50
1:B:470:ARG:HE	1:B:470:ARG:H	1.58	0.50
1:A:467:VAL:O	1:A:514:VAL:HG21	2.11	0.50
1:B:365:LEU:H	1:B:365:LEU:HD23	1.77	0.50
1:A:466:ALA:HB1	1:A:514:VAL:HG23	1.93	0.48
1:B:460:ARG:HB3	1:B:463:VAL:HB	1.96	0.48
1:B:443:TYR:HA	1:B:449:GLU:O	2.14	0.48
1:B:470:ARG:H	1:B:470:ARG:NE	2.12	0.48
1:A:550:MET:HE1	1:A:568:VAL:HG11	1.95	0.48
1:B:345:TYR:HB2	1:B:352:TRP:CH2	2.49	0.47
1:B:369:VAL:HG13	1:B:607:ALA:HB1	1.95	0.47
1:B:443:TYR:HB2	1:B:450:TRP:CD2	2.49	0.47

*Continued on next page...*



*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:468:LEU:HD12	1:B:469:ASN:N	2.30	0.47
1:B:425:ILE:HB	1:B:443:TYR:HB3	1.97	0.47
1:A:362:ARG:NH1	1:A:394:ASP:OD1	2.48	0.47
1:B:604:VAL:HG23	1:B:606:VAL:HG23	1.97	0.47
1:A:443:TYR:HB2	1:A:450:TRP:CE2	2.49	0.46
1:A:460:ARG:HB3	1:A:463:VAL:HB	1.97	0.46
1:B:418:VAL:HA	1:B:426:TYR:O	2.15	0.46
1:B:342:LEU:HD22	1:B:403:TRP:HZ2	1.81	0.46
1:B:354:ARG:NH2	1:B:357:ASP:OD1	2.42	0.45
1:A:494:ARG:HD2	1:A:496:GLU:HG2	1.98	0.45
1:B:380:ARG:CZ	2:P:83:GLU:OE1	2.62	0.45
1:A:365:LEU:HD12	1:A:376:ALA:HB1	1.99	0.45
1:A:557:LEU:HD23	1:A:557:LEU:H	1.82	0.45
1:A:443:TYR:HB2	1:A:450:TRP:CD2	2.52	0.45
1:B:347:PRO:HG2	1:B:562:HIS:CE1	2.52	0.45
1:A:490:TYR:HB2	1:A:497:TRP:CE2	2.52	0.44
1:B:584:TYR:HB2	1:B:591:TRP:CZ2	2.52	0.44
1:B:345:TYR:HB2	1:B:352:TRP:CZ3	2.53	0.44
1:A:328:ILE:O	1:A:344:ALA:HA	2.17	0.43
1:B:425:ILE:O	1:B:442:ARG:HA	2.18	0.43
1:A:393:LEU:HD22	1:A:450:TRP:CZ2	2.54	0.43
1:A:512:VAL:HA	1:A:520:TYR:O	2.19	0.43
1:A:519:ILE:O	1:A:536:ARG:HA	2.18	0.43
1:A:337:GLN:HG2	3:A:778:HOH:O	2.19	0.42
1:A:346:ASN:HB3	1:A:349:ASP:HB2	2.01	0.42
1:B:512:VAL:HA	1:B:520:TYR:O	2.20	0.42
1:A:347:PRO:HG2	1:A:562:HIS:CE1	2.55	0.42
1:A:416:ILE:HD13	1:A:416:ILE:HG21	1.81	0.42
1:B:352:TRP:HE1	1:B:595:THR:CG2	2.33	0.42
1:B:443:TYR:CE1	1:B:448:ASP:HA	2.55	0.42
1:A:566:ILE:HB	1:A:584:TYR:HB3	2.01	0.42
1:B:375:TYR:OH	1:B:445:PRO:HB3	2.19	0.42
1:B:469:ASN:O	3:B:701:HOH:O	2.22	0.41
1:B:532:ASN:O	1:B:550:MET:N	2.42	0.41
1:A:446:GLU:H	1:A:446:GLU:CD	2.24	0.41
1:B:328:ILE:HG21	1:B:560:THR:HG21	2.02	0.41
1:B:470:ARG:HD3	1:B:470:ARG:HA	1.87	0.41
1:A:562:HIS:HB3	1:A:567:TYR:CE2	2.55	0.41
1:B:519:ILE:O	1:B:536:ARG:HA	2.21	0.41
1:A:349:ASP:HB3	1:A:351:THR:HG23	2.03	0.40
1:B:473:TYR:CD1	1:B:519:ILE:HD11	2.56	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	284/301 (94%)	274 (96%)	9 (3%)	1 (0%)	34	41
1	B	281/301 (93%)	269 (96%)	12 (4%)	0	100	100
2	P	9/11 (82%)	8 (89%)	0	1 (11%)	0	0
All	All	574/613 (94%)	551 (96%)	21 (4%)	2 (0%)	41	50

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	611	GLU
2	P	85	ALA

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	230/245 (94%)	227 (99%)	3 (1%)	69	81
1	B	225/245 (92%)	217 (96%)	8 (4%)	35	48
2	P	6/6 (100%)	6 (100%)	0	100	100
All	All	461/496 (93%)	450 (98%)	11 (2%)	49	65

All (11) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	536	ARG
1	A	557	LEU
1	A	580	SER
1	B	349	ASP
1	B	354	ARG
1	B	434	CYS
1	B	447	ARG
1	B	498	ARG
1	B	551	LYS
1	B	557	LEU
1	B	596	ARG

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	424	HIS

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

### 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues

The following chains have linkage breaks:

Mol	Chain	Number of breaks
2	P	1

All chain breaks are listed below:

Model	Chain	Residue-1	Atom-1	Residue-2	Atom-2	Distance (Å)
1	P	84:PHE	C	85:ALA	N	1.18

## 6 Fit of model and data [i](#)

### 6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	286/301 (95%)	-0.26	3 (1%) 82 86	21, 28, 48, 74	0
1	B	283/301 (94%)	1.27	72 (25%) 0 1	25, 56, 105, 125	0
2	P	9/11 (81%)	0.15	0 100 100	35, 39, 55, 55	0
All	All	578/613 (94%)	0.49	75 (12%) 3 5	21, 37, 96, 125	0

All (75) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	351	THR	8.9
1	B	329	TYR	7.3
1	B	562	HIS	6.8
1	B	590	THR	6.5
1	B	328	ILE	5.8
1	B	327	LEU	5.4
1	B	352	TRP	5.2
1	B	595	THR	5.1
1	B	402	GLN	5.0
1	B	347	PRO	5.0
1	B	447	ARG	4.8
1	B	373	LEU	4.8
1	B	567	TYR	4.7
1	B	353	LEU	4.7
1	B	449	GLU	4.2
1	B	491	TYR	4.2
1	B	423	GLY	4.0
1	B	421	ILE	4.0
1	B	395	CYS	4.0
1	B	445	PRO	4.0
1	B	398	PRO	3.9
1	B	348	SER	3.8
1	B	346	ASN	3.8

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
1	B	592	SER	3.7
1	B	345	TYR	3.7
1	B	591	TRP	3.6
1	B	399	MET	3.6
1	B	349	ASP	3.5
1	B	422	ASP	3.5
1	B	594	VAL	3.5
1	B	581	VAL	3.4
1	B	561	VAL	3.4
1	B	608	VAL	3.4
1	B	424	HIS	3.4
1	B	446	GLU	3.3
1	B	596	ARG	3.2
1	B	494	ARG	3.1
1	B	375	TYR	3.1
1	B	400	THR	3.1
1	B	443	TYR	3.0
1	B	408	PRO	2.9
1	B	350	GLY	2.9
1	B	397	ASN	2.9
1	B	401	ASN	2.9
1	B	344	ALA	2.8
1	B	588	THR	2.8
1	B	575	HIS	2.7
1	B	425	ILE	2.7
1	B	444	GLU	2.7
1	B	393	LEU	2.7
1	B	597	MET	2.7
1	A	385	ASP	2.6
1	B	396	TYR	2.6
1	B	593	GLU	2.6
1	A	610	MET	2.6
1	B	589	ASP	2.5
1	B	442	ARG	2.5
1	B	371	GLY	2.5
1	B	565	ARG	2.5
1	B	370	VAL	2.4
1	B	517	ASN	2.4
1	B	374	LEU	2.4
1	B	493	GLU	2.3
1	B	452	LEU	2.3
1	B	454	ALA	2.3

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	420	VAL	2.2
1	B	470	ARG	2.2
1	B	516	HIS	2.1
1	B	552	HIS	2.1
1	A	386	GLY	2.1
1	B	448	ASP	2.1
1	B	355	LEU	2.1
1	B	495	ASN	2.1
1	B	451	HIS	2.0
1	B	539	VAL	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 6.4 Ligands [i](#)

There are no ligands in this entry.

## 6.5 Other polymers [i](#)

There are no such residues in this entry.