

wwPDB EM Validation Summary Report (i)

Dec 17, 2024 – 02:08 PM JST

PDB ID	:	8K4S
EMDB ID	:	EMD-36890
Title	:	CryoEM structure of Gq coupled MRGPRX4 with agonist DCA-3P
Authors	:	Yang, J.; Fan, J.P.; Lei, X.G.
Deposited on	:	2023-07-20
Resolution	:	2.90 Å(reported)

This is a wwPDB EM Validation Summary Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/EMValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

EMDB validation analysis	:	FAILED
Mogul	:	1.8.5 (274361), CSD as541be (2020)
MolProbity	:	4.02b-467
buster-report	:	1.1.7(2018)
Percentile statistics	:	20231227.v01 (using entries in the PDB archive December 27th 2023)
MapQ	:	FAILED
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $ELECTRON\ MICROSCOPY$

The reported resolution of this entry is 2.90 Å.

Sidechain outliers

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



206894

The table below summarises the geometric issues observed across the polymeric chains and their fit to the map. The red, orange, yellow and green segments of the bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

16415

Mol	Chain	Length	Quality of chain							
1	Е	710	27% 9%	64%						
2	А	246	71%	19% 10%						
3	В	340	69%	27% ••						
4	С	80	46% 12%	41%						
5	D	254	69%	20% • 9%						



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 8196 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Soluble cytochrome b562,Mas-related G-protein coupled receptor member X4,Green fluorescent protein.

Mol	Chain	Residues	Atoms					AltConf	Trace
1	Е	255	Total 1956	C 1304	N 314	O 322	S 16	0	0

Chain	Residue	Modelled	Actual	Comment	Reference
Е	-115	ASP	-	expression tag	UNP P0ABE7
Е	-114	TYR	-	expression tag	UNP P0ABE7
Е	-113	LYS	-	expression tag	UNP P0ABE7
Е	-112	ASP	-	expression tag	UNP P0ABE7
Е	-111	ASP	-	expression tag	UNP P0ABE7
Е	-110	ASP	-	expression tag	UNP P0ABE7
Е	-109	ASP	-	expression tag	UNP P0ABE7
Е	-108	LYS	-	expression tag	UNP P0ABE7
Е	-107	GLU	-	expression tag	UNP P0ABE7
Е	-106	PHE	-	expression tag	UNP P0ABE7
Е	-99	TRP	MET	conflict	UNP P0ABE7
Е	-4	ILE	HIS	conflict	UNP P0ABE7
Е	0	LEU	ARG	conflict	UNP P0ABE7
Е	323	LEU	-	linker	UNP Q96LA9
Е	324	GLU	-	linker	UNP Q96LA9
Е	325	LEU	-	linker	UNP Q96LA9
Е	326	GLU	-	linker	UNP Q96LA9
Е	327	VAL	-	linker	UNP Q96LA9
Е	328	LEU	-	linker	UNP Q96LA9
Е	329	PHE	-	linker	UNP Q96LA9
Е	330	GLN	-	linker	UNP Q96LA9
Е	331	GLY	-	linker	UNP Q96LA9
Е	332	PRO	-	linker	UNP Q96LA9
Е	361	ARG	SER	conflict	UNP P42212
Е	370	ASN	TYR	conflict	UNP P42212
Е	395	LEU	PHE	conflict	UNP P42212
Е	396	THR	SER	conflict	UNP P42212

There are 63 discrepancies between the modelled and reference sequences:

Continued on next page...



Continu	ied from pre	vious page			
Chain	Residue	Modelled	Actual	Comment	Reference
Е	411	ARG	GLN	conflict	UNP P42212
Е	430	SER	PHE	conflict	UNP P42212
Е	436	THR	ASN	conflict	UNP P42212
Е	476	PHE	TYR	conflict	UNP P42212
Е	484	THR	MET	conflict	UNP P42212
Е	494	ALA	VAL	conflict	UNP P42212
Е	502	VAL	ILE	conflict	UNP P42212
Е	537	VAL	ALA	conflict	UNP P42212
Е	567	TRP	-	expression tag	UNP P42212
Е	568	SER	-	expression tag	UNP P42212
Е	569	HIS	-	expression tag	UNP P42212
Е	570	PRO	-	expression tag	UNP P42212
Е	571	GLN	-	expression tag	UNP P42212
Е	572	PHE	-	expression tag	UNP P42212
Е	573	GLU	-	expression tag	UNP P42212
Е	574	LYS	-	expression tag	UNP P42212
Е	575	GLY	-	expression tag	UNP P42212
Е	576	GLY	-	expression tag	UNP P42212
Е	577	GLY	-	expression tag	UNP P42212
Е	578	SER	-	expression tag	UNP P42212
Е	579	GLY	-	expression tag	UNP P42212
Е	580	GLY	-	expression tag	UNP P42212
Е	581	GLY	-	expression tag	UNP P42212
Е	582	SER	-	expression tag	UNP P42212
Е	583	GLY	-	expression tag	UNP P42212
Е	584	GLY	-	expression tag	UNP P42212
Е	585	SER	-	expression tag	UNP P42212
Е	586	ALA	-	expression tag	UNP P42212
Е	587	TRP	-	expression tag	UNP P42212
Е	588	SER	-	expression tag	UNP P42212
Е	589	HIS	-	expression tag	UNP P42212
Е	590	PRO	-	expression tag	UNP P42212
Е	591	GLN	-	expression tag	UNP P42212
Е	592	PHE	-	expression tag	UNP P42212
Е	593	GLU	-	expression tag	UNP P42212
Е	594	LYS	-	expression tag	UNP P42212

 α 1: 1 0

• Molecule 2 is a protein called Gs-mini-Gq chimera.

Mol	Chain	Residues	Atoms					AltConf	Trace
2	А	222	Total 1673	C 1069	N 296	0 301	S 7	0	0



• Molecule 3 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(T) subunit beta-1.

Mol	Chain	Residues	Atoms					AltConf	Trace
3	В	338	Total 2481	C 1544	N 440	0 477	S 20	0	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	1	GLY	-	expression tag	UNP P62873

- Molecule 4 is a protein called Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2.

Mol	Chain	Residues	Atoms				AltConf	Trace	
4	С	47	Total 329	C 210	N 56	O 61	$\begin{array}{c} \mathrm{S} \\ \mathrm{2} \end{array}$	0	0

There are 9 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
С	72	GLY	-	expression tag	UNP P59768
С	73	SER	-	expression tag	UNP P59768
С	74	ALA	-	expression tag	UNP P59768
С	75	GLY	-	expression tag	UNP P59768
С	76	SER	-	expression tag	UNP P59768
С	77	ALA	-	expression tag	UNP P59768
С	78	GLY	-	expression tag	UNP P59768
С	79	SER	-	expression tag	UNP P59768
С	80	ALA	-	expression tag	UNP P59768

• Molecule 5 is a protein called scFv16.

Mol	Chain	Residues	Atoms					AltConf	Trace
5	D	231	Total 1725	C 1098	N 288	O 329	S 10	0	0

• Molecule 6 is $(4 \{R\})-4-[(3 \{R\},5 \{R\},8 \{R\},9 \{S\},10 \{S\},12 \{S\},13 \{R\},14 \{S\},17 \{R\})-1 0,13-dimethyl-12-oxidanyl-3-phosphonooxy-2,3,4,5,6,7,8,9,11,12,14,15,16,17-tetradecahydr o-1 {H}-cyclopenta[a]phenanthren-17-yl]pentanoic acid (three-letter code: JW0) (formula: C₂₄H₄₁O₇P) (labeled as "Ligand of Interest" by depositor).$





Mol	Chain	Residues	Atoms			AltConf	
6	Ε	1	Total 32	C 24	O 7	Р 1	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Soluble cytochrome b562,Mas-related G-protein coupled receptor member X4,Green fluorescent protein



Chain A: 71% 19% 10%



• Molecule 3: Guanine nucleotide-binding protein $\rm G(I)/\rm G(S)/\rm G(T)$ subunit beta-1



• Molecule 4: Guanine nucleotide-binding protein G(I)/G(S)/G(O) subunit gamma-2



• Molecule 5: scFv16







4 Experimental information (i)

Property	Value	Source
EM reconstruction method	SINGLE PARTICLE	Depositor
Imposed symmetry	POINT, Not provided	
Number of particles used	451859	Depositor
Resolution determination method	FSC 0.143 CUT-OFF	Depositor
CTF correction method	PHASE FLIPPING AND AMPLITUDE	Depositor
	CORRECTION	
Microscope	FEI TITAN KRIOS	Depositor
Voltage (kV)	300	Depositor
Electron dose $(e^-/\text{\AA}^2)$	60	Depositor
Minimum defocus (nm)	1100	Depositor
Maximum defocus (nm)	2200	Depositor
Magnification	Not provided	
Image detector	GATAN K2 SUMMIT (4k x 4k)	Depositor



5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $\mathrm{JW0}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
	Unain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	Е	0.25	0/2003	0.52	0/2744	
2	А	0.26	0/1702	0.49	0/2309	
3	В	0.25	0/2528	0.52	0/3443	
4	С	0.23	0/335	0.44	0/459	
5	D	0.27	0/1769	0.53	0/2406	
All	All	0.26	0/8337	0.51	0/11361	

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	Е	1956	0	1986	40	0
2	А	1673	0	1571	34	0
3	В	2481	0	2305	69	0
4	С	329	0	309	8	0
5	D	1725	0	1614	35	0
6	Е	32	0	0	3	0
All	All	8196	0	7785	167	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including



hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 167 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:E:67:ASN:ND2	1:E:143:CYS:SG	2.44	0.88
1:E:100:SER:HG	1:E:158:TRP:HE1	1.20	0.87
3:B:104:ALA:HB3	3:B:113:ALA:HB3	1.69	0.74
3:B:283:ARG:HB3	4:C:51:LEU:HD21	1.72	0.72
2:A:98:GLN:OE1	3:B:314:ARG:NH1	2.22	0.72

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	entiles
1	Е	251/710~(35%)	231 (92%)	20 (8%)	0	100	100
2	А	216/246~(88%)	206 (95%)	10 (5%)	0	100	100
3	В	336/340~(99%)	325 (97%)	11 (3%)	0	100	100
4	С	45/80~(56%)	43 (96%)	2 (4%)	0	100	100
5	D	227/254~(89%)	214 (94%)	13 (6%)	0	100	100
All	All	1075/1630~(66%)	1019 (95%)	56 (5%)	0	100	100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all EM entries.

The Analysed column shows the number of residues for which the sidechain conformation was



Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	Ε	212/621~(34%)	199~(94%)	13~(6%)	15	43
2	А	160/213~(75%)	157 (98%)	3(2%)	52	81
3	В	248/282~(88%)	226~(91%)	22 (9%)	8	26
4	С	30/61~(49%)	30 (100%)	0	100	100
5	D	180/203~(89%)	167~(93%)	13~(7%)	12	35
All	All	830/1380~(60%)	779~(94%)	51~(6%)	18	43

analysed, and the total number of residues.

5 of 51 residues with a non-rotameric side chain are listed below:

Mol	Chain	\mathbf{Res}	Type
3	В	163	ASP
3	В	294	CYS
5	D	201	THR
3	В	198	LEU
3	В	258	ASP

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	Е	268	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.

5.6 Ligand geometry (i)

1 ligand is modelled in this entry.



In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol Type Chain	Chain	Ros	Ros	Bos	Link	B	ond leng	gths	B	ond ang	gles
WIOI	viol Type Chain Res	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z >2		
6	JW0	Е	601	-	$35,\!35,\!35$	2.16	11 (31%)	56,56,56	2.40	19 (33%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
6	JW0	Е	601	-	-	10/14/76/76	0/4/4/4

The worst 5 of 11 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
6	Е	601	JW0	P1-04	5.84	1.70	1.59
6	Е	601	JW0	O1-C13	-4.70	1.35	1.43
6	Е	601	JW0	C12-C17	-4.64	1.47	1.55
6	Е	601	JW0	C14-C10	-4.46	1.46	1.53
6	Е	601	JW0	C6-C5	-3.69	1.47	1.54

The worst 5 of 19 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
6	Е	601	JW0	C12-C17-C18	-8.68	109.14	119.50
6	Е	601	JW0	C12-C11-C9	-5.54	108.59	114.71
6	Е	601	JW0	C14-C10-C5	-4.81	108.77	113.73
6	Е	601	JW0	C11-C9-C10	-4.20	103.47	109.09
6	Е	601	JW0	C11-C12-C13	3.75	110.89	107.40

There are no chirality outliers.

5 of 10 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
6	Е	601	JW0	C1-C2-O4-P1

Continued on next page...



Mol	Chain	Res	Type	Atoms
6	Ε	601	JW0	C12-C17-C18-C21
6	Е	601	JW0	C16-C17-C18-C21
6	Ε	601	JW0	C12-C17-C18-C19
6	Е	601	JW0	C16-C17-C18-C19

Continued from previous page...

There are no ring outliers.

1 monomer is involved in 3 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
6	Е	601	JW0	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.





5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

