



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 04:27 AM EDT

PDB ID : 1K6C
Title : LACK OF SYNERGY FOR INHIBITORS TARGETING A MULTI-DRUG RESISTANT HIV-1 PROTEASE
Authors : Schiffer, C.A.
Deposited on : 2001-10-15
Resolution : 2.20 Å (reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtrriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
buster-report : 1.1.7 (2018)
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.20 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	5594 (2.20-2.20)
Ramachandran outliers	138981	5503 (2.20-2.20)
Sidechain outliers	138945	5504 (2.20-2.20)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	99	77% 21% .
1	B	99	81% 17% .

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	ACT	B	502	-	-	X	-
2	ACT	B	506	-	-	X	-

2 Entry composition

There are 4 unique types of molecules in this entry. The entry contains 1786 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called POL polyprotein.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	99	798	512	138	144	4	0	5	0
1	B	99	804	516	139	145	4	5	6	0

There are 8 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	7	LYS	GLN	engineered mutation	UNP P35963
A	14	ARG	LYS	engineered mutation	UNP P35963
A	82	THR	VAL	engineered mutation	UNP P35963
A	84	VAL	ILE	engineered mutation	UNP P35963
B	7	LYS	GLN	engineered mutation	UNP P35963
B	14	ARG	LYS	engineered mutation	UNP P35963
B	82	THR	VAL	engineered mutation	UNP P35963
B	84	VAL	ILE	engineered mutation	UNP P35963

- Molecule 2 is ACETATE ION (three-letter code: ACT) (formula: C₂H₃O₂).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	A	1	Total C O 4 2 2	0	0
2	A	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0
2	B	1	Total C O 4 2 2	0	0

- Molecule 3 is N-[2(R)-HYDROXY-1(S)-INDANYL]-5-[(2(S)-TERTIARY BUTYLAMINOCARBONYL)-4(3-PYRIDYLMETHYL)PIPERAZINO]-4(S)-HYDROXY-2(R)-PHENYLME THYLPENTANAMIDE (three-letter code: MK1) (formula: C₃₆H₄₇N₅O₄).



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
			Total	C	N	O		
3	B	1	45	36	5	4	0	0

- Molecule 4 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	50	Total	O	0	0
			50	50		
4	B	69	Total	O	0	0
			69	69		

3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


Note EDS was not executed.

- Molecule 1: POL polyprotein

Chain A:  77% 21%



- Molecule 1: POL polyprotein

Chain B:  81% 17%



4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α , β , γ	51.45Å 59.40Å 61.74Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	27.00 – 2.20	Depositor
% Data completeness (in resolution range)	99.2 (27.00-2.20)	Depositor
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
Refinement program	CNS	Depositor
R, R_{free}	0.194 , 0.231	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	1786	wwPDB-VP
Average B, all atoms (Å ²)	38.0	wwPDB-VP

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: ACT, MK1

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.35	0/733	0.65	0/988
1	B	0.36	0/726	0.64	0/976
All	All	0.36	0/1459	0.65	0/1964

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	798	0	838	27	0
1	B	804	0	844	34	0
2	A	8	0	6	0	0
2	B	12	0	9	7	0
3	B	45	0	47	4	0
4	A	50	0	0	13	0
4	B	69	0	0	8	0
All	All	1786	0	1744	59	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 17.

All (59) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:50[1]:ILE:HB	4:A:546:HOH:O	1.63	0.96
2:B:506:ACT:H2	4:B:929:HOH:O	1.66	0.96
2:B:506:ACT:CH3	4:B:929:HOH:O	2.16	0.92
4:A:546:HOH:O	1:B:49:GLY:HA3	1.74	0.86
1:B:61:GLN:HE21	1:B:61:GLN:HA	1.41	0.84
1:A:50[2]:ILE:HB	4:A:546:HOH:O	1.79	0.83
1:B:37:ASN:HD22	2:B:502:ACT:H3	1.44	0.83
1:B:37:ASN:ND2	2:B:502:ACT:H3	1.95	0.82
1:A:18:GLN:HB3	4:A:548:HOH:O	1.83	0.77
1:A:14:ARG:HH11	1:A:14:ARG:HB3	1.51	0.76
1:B:67:CYS:SG	4:B:970:HOH:O	2.45	0.74
1:B:20:LYS:HE3	1:B:36:MET:SD	2.26	0.74
2:B:506:ACT:O	4:B:940:HOH:O	2.14	0.66
1:B:61:GLN:HA	1:B:61:GLN:NE2	2.10	0.64
1:B:32[1]:VAL:HG22	1:B:76:LEU:HB2	1.78	0.63
1:B:55:LYS:HD2	4:B:958:HOH:O	2.00	0.61
1:A:14:ARG:HH11	1:A:14:ARG:CB	2.14	0.61
4:A:550:HOH:O	1:B:81:PRO:HG3	2.01	0.60
1:A:14:ARG:HB3	1:A:14:ARG:NH1	2.16	0.59
1:A:95:CYS:HA	1:B:99:PHE:HB3	1.86	0.58
1:A:35[2]:GLU:OE2	1:A:78:GLY:HA2	2.04	0.57
4:A:550:HOH:O	1:B:81:PRO:CG	2.53	0.56
1:A:63:PRO:HG2	4:A:535:HOH:O	2.06	0.55
1:B:37:ASN:ND2	2:B:502:ACT:CH3	2.68	0.55
1:A:47[1]:ILE:HD12	3:B:902:MK1:H71	1.89	0.55
1:A:50[2]:ILE:HD13	1:B:81:PRO:CD	2.38	0.54
1:B:32[2]:VAL:HG12	1:B:84[2]:VAL:HG22	1.91	0.52
4:A:546:HOH:O	1:B:49:GLY:CA	2.44	0.52
1:A:45:LYS:NZ	4:A:534:HOH:O	2.41	0.52
1:B:32[1]:VAL:HG23	3:B:902:MK1:H27	1.90	0.52
1:B:32[2]:VAL:HG12	1:B:84[2]:VAL:CG2	2.40	0.52
1:B:35[2]:GLU:HA	1:B:77:VAL:CG1	2.40	0.51
1:A:49:GLY:HA3	1:B:50[2]:ILE:HB	1.92	0.51
1:A:36:MET:HG3	1:A:38:LEU:HD13	1.92	0.51
1:A:11:VAL:HG21	1:A:24:LEU:HD21	1.93	0.51
1:B:25:ASP:HB3	1:B:84[1]:VAL:HG13	1.94	0.50
1:A:67:CYS:HA	4:A:533:HOH:O	2.12	0.48
1:A:61:GLN:HE21	1:A:61:GLN:HA	1.78	0.48
1:B:32[2]:VAL:CG1	1:B:84[2]:VAL:HG22	2.44	0.48
1:A:50[2]:ILE:HD13	1:B:81:PRO:HD2	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:30:ASP:HB2	3:B:902:MK1:H26	1.95	0.47
1:B:10:LEU:HD11	4:B:931:HOH:O	2.14	0.47
1:B:35[1]:GLU:HB2	4:B:934:HOH:O	2.15	0.47
1:B:25:ASP:CB	1:B:84[1]:VAL:HG13	2.45	0.46
1:A:57:ARG:NH1	1:A:59:TYR:OH	2.50	0.45
1:B:50[2]:ILE:HG13	3:B:902:MK1:H72	1.99	0.44
1:A:14:ARG:HG3	1:A:19:LEU:HD23	1.99	0.43
1:B:61:GLN:HE21	1:B:61:GLN:CA	2.11	0.43
1:A:13:ILE:HD12	1:A:64:ILE:HD11	2.01	0.43
1:A:43:LYS:NZ	4:A:549:HOH:O	2.52	0.43
1:A:50[2]:ILE:HD13	1:B:81:PRO:HD3	2.00	0.43
1:B:70:LYS:HB3	4:B:937:HOH:O	2.18	0.42
4:A:550:HOH:O	1:B:81:PRO:HD3	2.18	0.42
1:B:36:MET:HB2	2:B:502:ACT:O	2.19	0.42
1:A:61:GLN:HA	1:A:61:GLN:NE2	2.34	0.42
1:A:11:VAL:HG11	1:A:24:LEU:HD11	2.02	0.41
1:A:35[1]:GLU:OE1	1:A:57:ARG:NH2	2.53	0.41
1:B:35[1]:GLU:HA	1:B:77:VAL:CG1	2.50	0.41
1:A:34[1]:GLU:CD	4:A:537:HOH:O	2.59	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	102/99 (103%)	101 (99%)	1 (1%)	0	100	100
1	B	103/99 (104%)	100 (97%)	1 (1%)	2 (2%)	8	5
All	All	205/198 (104%)	201 (98%)	2 (1%)	2 (1%)	29	14

All (2) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	50[1]	ILE
1	B	50[2]	ILE

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	79/83 (95%)	77 (98%)	2 (2%)	47	60
1	B	78/83 (94%)	77 (99%)	1 (1%)	69	81
All	All	157/166 (95%)	154 (98%)	3 (2%)	57	71

All (3) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	38	LEU
1	A	67	CYS
1	B	67	CYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (4) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	58	GLN
1	A	61	GLN
1	B	37	ASN
1	B	61	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

6 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	ACT	A	501	-	1,3,3	3.08	1 (100%)	0,3,3	-	-
3	MK1	B	902	-	49,49,49	2.32	12 (24%)	59,69,69	2.29	12 (20%)
2	ACT	B	502	-	1,3,3	2.95	1 (100%)	0,3,3	-	-
2	ACT	B	504	-	1,3,3	4.51	1 (100%)	0,3,3	-	-
2	ACT	B	506	-	1,3,3	3.66	1 (100%)	0,3,3	-	-
2	ACT	A	503	-	1,3,3	3.03	1 (100%)	0,3,3	-	-

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	MK1	B	902	-	-	3/33/58/58	0/5/5/5

All (17) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	902	MK1	C25-C30	10.19	1.57	1.39
3	B	902	MK1	C6-C4	-5.18	1.40	1.52
2	B	504	ACT	CH3-C	4.51	1.54	1.48
3	B	902	MK1	C33-N5	4.48	1.44	1.34
2	B	506	ACT	CH3-C	3.66	1.53	1.48
3	B	902	MK1	C30-C22	3.53	1.55	1.51
2	A	501	ACT	CH3-C	3.08	1.52	1.48

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	B	902	MK1	C34-N5	3.04	1.42	1.33
2	A	503	ACT	CH3-C	3.03	1.52	1.48
2	B	502	ACT	CH3-C	2.95	1.52	1.48
3	B	902	MK1	C2-N3	2.87	1.52	1.48
3	B	902	MK1	C1-N1	2.69	1.51	1.46
3	B	902	MK1	C13-C21	2.61	1.56	1.51
3	B	902	MK1	C16-C15	2.17	1.43	1.38
3	B	902	MK1	C18-C17	2.13	1.43	1.38
3	B	902	MK1	C7-C4	-2.05	1.47	1.52
3	B	902	MK1	C28-C27	2.05	1.43	1.38

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
3	B	902	MK1	C24-C25-C30	-7.84	101.26	110.99
3	B	902	MK1	C25-C24-C23	7.26	110.48	103.50
3	B	902	MK1	O2-C11-C10	-5.29	94.69	109.87
3	B	902	MK1	C24-C25-C26	4.39	137.39	129.18
3	B	902	MK1	C4-N2-C3	4.07	131.78	126.09
3	B	902	MK1	C29-C30-C25	-4.02	116.00	120.80
3	B	902	MK1	C8-N3-C2	4.01	117.96	110.83
3	B	902	MK1	C32-C31-N1	3.76	120.43	113.12
3	B	902	MK1	C29-C30-C22	3.12	133.15	128.99
3	B	902	MK1	C34-N5-C33	-2.50	112.52	116.85
3	B	902	MK1	C7-C4-N2	2.32	114.47	108.75
3	B	902	MK1	C8-N3-C10	-2.24	107.61	111.81

There are no chirality outliers.

All (3) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	B	902	MK1	N3-C10-C11-C12
3	B	902	MK1	C11-C10-N3-C2
3	B	902	MK1	C11-C10-N3-C8

There are no ring outliers.

3 monomers are involved in 11 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	B	902	MK1	4	0
2	B	502	ACT	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	506	ACT	3	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.



5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.