

wwPDB X-ray Structure Validation Summary Report (i)

Nov 4, 2024 – 04:16 PM EST

PDB ID : 7K6C

Title : Crystal Structure of Dihydrofolate reductase (DHFR) from Mycobacterium

abscessus ATCC 19977 / DSM 44196 with NADP and inhibitor P218

Authors: Seattle Structural Genomics Center for Infectious Disease (SSGCID); Aben-

droth, J.; Santhakumar, V.; Walpole, C.; Lorimer, D.D.; Horanyi, P.S.; Ed-

wards, T.E.

Deposited on : 2020-09-19

Resolution : 2.00 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

 $Mol Probity \quad : \quad 4.02b\text{--}467$

Mogul : 2022.3.0, CSD as543be (2022)

 $Xtriage\ (Phenix) \quad : \quad 1.20.1$

EDS: 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.003 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)

Validation Pipeline (wwPDB-VP) : 2.39




1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X- $RAY\ DIFFRACTION$

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},\ {\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	164625	9409 (2.00-2.00)
Clashscore	180529	10737 (2.00-2.00)
Ramachandran outliers	177936	10628 (2.00-2.00)
Sidechain outliers	177891	10627 (2.00-2.00)
RSRZ outliers	164620	9409 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain		
1		1 7 1	2%		
1	A	171	89%	•	7%
1	В	171	88%	5%	6%
1	С	171	91%	•	7%
1	D	171	89%	•	7%
1	Е	171	84%	9%	7%



Continued from previous page...

Mol	Chain	Length	Quality of chain		
1	F	171	84%	9%	7%
1	G	171	87%	6%	6%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 9892 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dihydrofolate reductase.

Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	159	Total	С	N	О	S	0	2	0
1	A	109	1193	761	209	220	3	U		0
1	В	160	Total	С	N	О	S	0	3	0
1	Б	100	1215	775	213	224	3	U) 	
1	С	150	Total	С	N	О	S	0	4	0
1		159	1216	777	214	222	3	0	4	
1	D	159	Total	С	N	О	S	0	4	0
1	D	199	1230	785	212	230	3			U
1	Е	159	Total	С	N	О	S	0	2	0
1	15	159	1211	774	214	220	3	0	<u> </u>	0
1	F	150	Total	С	N	О	S	0	4	0
1	1 F	159	1206	772	216	215	3	U	4	0
1	G	160	Total	С	N	О	S	0	2	0
1	G	100	1222	779	217	223	3			0

There are 70 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-7	MET	-	initiating methionine	UNP B1MD46
A	-6	ALA	-	expression tag	UNP B1MD46
A	-5	HIS	-	expression tag	UNP B1MD46
A	-4	HIS	-	expression tag	UNP B1MD46
A	-3	HIS	-	expression tag	UNP B1MD46
A	-2	HIS	-	expression tag	UNP B1MD46
A	-1	HIS	-	expression tag	UNP B1MD46
A	0	HIS	-	expression tag	UNP B1MD46
A	84	SER	CYS	engineered mutation	UNP B1MD46
A	141	THR	GLU	engineered mutation	UNP B1MD46
В	-7	MET	-	initiating methionine	UNP B1MD46
В	-6	ALA	-	expression tag	UNP B1MD46
В	-5	HIS	=	expression tag	UNP B1MD46
В	-4	HIS	-	expression tag	UNP B1MD46
В	-3	HIS	-	expression tag	UNP B1MD46



Continued from previous page...


Chain	Residue	Modelled	Actual	Comment	Reference
В	-2	HIS	-	expression tag	UNP B1MD46
В	-1	HIS	-	expression tag	UNP B1MD46
В	0	HIS	-	expression tag	UNP B1MD46
В	84	SER	CYS	engineered mutation	UNP B1MD46
В	141	THR	GLU	engineered mutation	UNP B1MD46
С	-7	MET	_	initiating methionine	UNP B1MD46
С	-6	ALA	-	expression tag	UNP B1MD46
С	-5	HIS	-	expression tag	UNP B1MD46
С	-4	HIS	-	expression tag	UNP B1MD46
С	-3	HIS	-	expression tag	UNP B1MD46
С	-2	HIS	-	expression tag	UNP B1MD46
С	-1	HIS	-	expression tag	UNP B1MD46
С	0	HIS	-	expression tag	UNP B1MD46
С	84	SER	CYS	engineered mutation	UNP B1MD46
С	141	THR	GLU	engineered mutation	UNP B1MD46
D	-7	MET	-	initiating methionine	UNP B1MD46
D	-6	ALA	-	expression tag	UNP B1MD46
D	-5	HIS	-	expression tag	UNP B1MD46
D	-4	HIS	-	expression tag	UNP B1MD46
D	-3	HIS	-	expression tag	UNP B1MD46
D	-2	HIS	-	expression tag	UNP B1MD46
D	-1	HIS	-	expression tag	UNP B1MD46
D	0	HIS	-	expression tag	UNP B1MD46
D	84	SER	CYS	engineered mutation	UNP B1MD46
D	141	THR	GLU	engineered mutation	UNP B1MD46
Е	-7	MET	-	initiating methionine	UNP B1MD46
E	-6	ALA	-	expression tag	UNP B1MD46
Е	-5	HIS	ı	expression tag	UNP B1MD46
Е	-4	HIS	I	expression tag	UNP B1MD46
Е	-3	HIS	-	expression tag	UNP B1MD46
Е	-2	HIS	-	expression tag	UNP B1MD46
E	-1	HIS	ı	expression tag	UNP B1MD46
Е	0	HIS	-	expression tag	UNP B1MD46
Е	84	SER	CYS	engineered mutation	UNP B1MD46
Е	141	THR	GLU	engineered mutation	UNP B1MD46
F	-7	MET	-	initiating methionine	UNP B1MD46
F	-6	ALA	-	expression tag	UNP B1MD46
F	-5	HIS	-	expression tag	UNP B1MD46
F	-4	HIS	_	expression tag	UNP B1MD46
F	-3	HIS	-	expression tag	UNP B1MD46
F	-2	HIS	-	expression tag	UNP B1MD46
F	-1	HIS	-	expression tag	UNP B1MD46



Continued from previous page...

Chain	Residue	Modelled	Actual	Comment	Reference
F	0	HIS	-	expression tag	UNP B1MD46
F	84	SER	CYS	engineered mutation	UNP B1MD46
F	141	THR	GLU	engineered mutation	UNP B1MD46
G	-7	MET	-	initiating methionine	UNP B1MD46
G	-6	ALA	-	expression tag	UNP B1MD46
G	-5	HIS	-	expression tag	UNP B1MD46
G	-4	HIS	-	expression tag	UNP B1MD46
G	-3	HIS	-	expression tag	UNP B1MD46
G	-2	HIS	-	expression tag	UNP B1MD46
G	-1	HIS	-	expression tag	UNP B1MD46
G	0	HIS	-	expression tag	UNP B1MD46
G	84	SER	CYS	engineered mutation	UNP B1MD46
G	141	THR	GLU	engineered mutation	UNP B1MD46

• Molecule 2 is NADP NICOTINAMIDE-ADENINE-DINUCLEOTIDE PHOSPHATE (three-letter code: NAP) (formula: $C_{21}H_{28}N_7O_{17}P_3$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	
2	٨	1	Total C N O P		0					
2	2 A	1	48	21	7	17	3	U	0	
2	2 B	1	Total C N O P	0	0					
2			48	21	7	17	3	0	U	
2	С	1	Total	С	N	О	Р	0	0	
2		1	48	21	7	17	3	U		
2	9 D	1	Total	С	N	О	Р	0	0	
	ש	1	48	21	7	17	3	U	U	



Continued from previous page...

Mol	Chain	Residues	${f Atoms}$					ZeroOcc	AltConf	
2	2 E	1	Total	С	N	О	Р	0	0	
	1	48	21	7	17	3	U			
2	2 F	F 1	Total	С	N	О	Р	0	0	
2			48	21	7	17	3	0		
9	С	1	Total	С	N	О	Р	0	0	
2	G		48	21	7	17	3	U		

• Molecule 3 is 3-(2-{3-[(2,4-diamino-6-ethylpyrimidin-5-yl)oxy]propoxy}phenyl)propanoic acid (three-letter code: MMV) (formula: $C_{18}H_{24}N_4O_4$) (labeled as "Ligand of Interest" by depositor).




Mol	Chain	Residues	Α	ton	ıs		ZeroOcc	AltConf			
3	A	1	Total	С	N	О	0	0			
3	A	1	26	18	4	4	U	0			
3	В	1	Total	С	N	О	0	0			
9	Б	1	26	18	4	4	U	U			
3	С	1	Total	С	N	О	0	0			
9	3	1	26	18	4	4	U				
3	D	D	D	D	1	Total	С	N	О	0	0
J	D	1	26	18	4	4	U	U			
3	E	1	Total	С	N	О	0	0			
J	12	1	26	18	4	4	U	0			
3	F	1	Total	С	N	Ο	0	0			
	I'	1	26	18	4	4	U	U			
2	3 G	1	Total	С	N	О	0	0			
		1	26	18	4	4	U	U			



• Molecule 4 is CALCIUM ION (three-letter code: CA) (formula: Ca).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total Ca 1 1	0	0
4	В	2	Total Ca 2 2	0	0
4	С	1	Total Ca 1 1	0	0

• Molecule 5 is 1,2-ETHANEDIOL (three-letter code: EDO) (formula: $C_2H_6O_2$).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	В	1	Total C O 4 2 2	0	0
5	С	1	Total C O 4 2 2	0	0
5	D	1	Total C O 4 2 2	0	0
5	E	1	Total C O 4 2 2	0	0
5	F	1	Total C O 4 2 2	0	0
5	G	1	Total C O 4 2 2	0	0

• Molecule 6 is water.




Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	67	Total O 67 67	0	0
6	В	153	Total O 153 153	0	0
6	С	100	Total O 102 102	0	2
6	D	172	Total O 172 172	0	0
6	E	120	Total O 120 120	0	0
6	F	139	Total O 139 139	0	0
6	G	100	Total O 100 100	0	0



3 Residue-property plots (i)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Dihydrofolate reductase





• Molecule 1: Dihydrofolate reductase





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	75.63Å 62.49Å 124.55Å	Depositor
a, b, c, α , β , γ	90.00° 94.67° 90.00°	Depositor
Resolution (Å)	37.69 - 2.00	Depositor
Resolution (A)	37.69 - 2.00	EDS
% Data completeness	99.7 (37.69-2.00)	Depositor
(in resolution range)	99.7 (37.69-2.00)	EDS
R_{merge}	0.07	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	2.42 (at 2.00Å)	Xtriage
Refinement program	PHENIX 1.18.2	Depositor
P. P.	0.166 , 0.207	Depositor
R, R_{free}	0.166 , 0.207	DCC
R_{free} test set	2290 reflections (2.66%)	wwPDB-VP
Wilson B-factor (Å ²)	25.5	Xtriage
Anisotropy	0.441	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 47.1	EDS
L-test for twinning ²	$ < L > = 0.49, < L^2> = 0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	9892	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 8.81% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAP, EDO, MMV, CA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5
1	A	0.35	0/1228	0.60	0/1679
1	В	0.42	0/1254	0.64	0/1714
1	С	0.35	0/1258	0.61	0/1718
1	D	0.40	0/1269	0.62	0/1735
1	Е	0.37	0/1247	0.62	0/1703
1	F	0.38	0/1248	0.65	0/1706
1	G	0.36	0/1258	0.58	0/1717
All	All	0.38	0/8762	0.62	0/11972

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1193	0	1161	3	0
1	В	1215	0	1196	7	0
1	С	1216	0	1209	3	0
1	D	1230	0	1214	5	0
1	Е	1211	0	1201	11	0
1	F	1206	0	1196	9	0



Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	G	1222	0	1213	6	0
2	A	48	0	25	0	0
2	В	48	0	25	0	0
2	С	48	0	25	0	0
2	D	48	0	25	1	0
2	Ε	48	0	25	0	0
2	F	48	0	25	1	0
2	G	48	0	25	1	0
3	A	26	0	23	0	0
3	В	26	0	23	0	0
3	С	26	0	23	0	0
3	D	26	0	23	0	0
3	Е	26	0	23	0	0
3	F	26	0	23	0	0
3	G	26	0	23	0	0
4	A	1	0	0	0	0
4	В	2	0	0	0	0
4	С	1	0	0	0	0
5	В	4	0	6	0	0
5	С	4	0	6	0	0
5	D	4	0	6	0	0
5	Е	4	0	6	1	0
5	F	4	0	6	1	0
5	G	4	0	6	0	0
6	A	67	0	0	0	0
6	В	153	0	0	3	1
6	С	102	0	0	1	0
6	D	172	0	0	1	2
6	Е	120	0	0	3	1
6	F	139	0	0	1	2
6	G	100	0	0	0	0
All	All	9892	0	8762	42	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

The worst 5 of 42 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
1:F:36[A]:ARG:NH1	6:F:301:HOH:O	2.25	0.70
1:G:53:LEU:O	1:G:58:ARG:NH2	2.30	0.65



Continued from previous page...

Atom-1	Atom-2	Interatomic	Clash
1100111 1	1100111 2	$\operatorname{distance}\left(\mathrm{\AA}\right)$	overlap (Å)
1:E:71:ASP:OD2	6:E:301:HOH:O	2.14	0.64
1:B:33:ARG:NE	6:B:303:HOH:O	2.32	0.57
1:D:7:LEU:HD13	1:D:112:ALA:HB2	1.86	0.57

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} ({\rm \AA}) \end{array}$	$egin{aligned} ext{Clash} \ ext{overlap } (ext{Å}) \end{aligned}$
6:D:433:HOH:O	6:F:342:HOH:O[1_545]	2.03	0.17
6:B:375:HOH:O	6:E:331:HOH:O[2_555]	2.16	0.04
6:D:433:HOH:O	6:F:380:HOH:O[1_545]	2.19	0.01

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	A	159/171~(93%)	156 (98%)	3 (2%)	0	100	100
1	В	161/171~(94%)	158 (98%)	3 (2%)	0	100	100
1	\mathbf{C}	161/171~(94%)	158 (98%)	3 (2%)	0	100	100
1	D	161/171~(94%)	158 (98%)	3 (2%)	0	100	100
1	E	159/171~(93%)	156 (98%)	3 (2%)	0	100	100
1	F	161/171~(94%)	158 (98%)	3 (2%)	0	100	100
1	G	160/171~(94%)	157 (98%)	3 (2%)	0	100	100
All	All	1122/1197 (94%)	1101 (98%)	21 (2%)	0	100	100

There are no Ramachandran outliers to report.



5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Perce	ntiles
1	A	117/133 (88%)	117 (100%)	0	100	100
1	В	122/133~(92%)	122 (100%)	0	100	100
1	С	123/133 (92%)	123 (100%)	0	100	100
1	D	$126/133 \ (95\%)$	126 (100%)	0	100	100
1	E	121/133 (91%)	121 (100%)	0	100	100
1	F	119/133 (90%)	119 (100%)	0	100	100
1	G	123/133 (92%)	122 (99%)	1 (1%)	79	84
All	All	851/931 (91%)	850 (100%)	1 (0%)	92	95

All (1) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	G	148	GLU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no oligosaccharides in this entry.



5.6 Ligand geometry (i)

Of 24 ligands modelled in this entry, 4 are monoatomic - leaving 20 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Во	Bond lengths		В	ond ang	les
IVIOI	Type	Chain	nes	LIIIK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	EDO	D	203	-	3,3,3	0.42	0	2,2,2	0.71	0
2	NAP	F	201	-	46,52,52	0.63	0	61,80,80	0.74	1 (1%)
5	EDO	F	203	-	3,3,3	0.25	0	2,2,2	1.45	0
2	NAP	В	201	-	46,52,52	0.74	1 (2%)	61,80,80	0.81	1 (1%)
2	NAP	A	201	-	46,52,52	0.72	1 (2%)	61,80,80	0.71	1 (1%)
3	MMV	G	202	-	27,27,27	1.07	3 (11%)	32,35,35	1.21	5 (15%)
3	MMV	A	202	-	27,27,27	1.18	3 (11%)	32,35,35	1.22	5 (15%)
2	NAP	С	201	-	46,52,52	0.74	1 (2%)	61,80,80	0.77	1 (1%)
5	EDO	С	203	-	3,3,3	0.37	0	2,2,2	0.99	0
3	MMV	D	202	-	27,27,27	1.13	3 (11%)	32,35,35	0.95	1 (3%)
3	MMV	F	202	-	27,27,27	1.19	3 (11%)	32,35,35	1.30	5 (15%)
2	NAP	D	201	-	46,52,52	0.75	1 (2%)	61,80,80	0.74	1 (1%)
3	MMV	В	202	-	27,27,27	1.18	3 (11%)	32,35,35	1.04	0
2	NAP	G	201	-	46,52,52	0.68	1 (2%)	61,80,80	0.69	1 (1%)
2	NAP	Е	201	-	46,52,52	0.69	0	61,80,80	0.77	1 (1%)
5	EDO	Е	203	-	3,3,3	0.46	0	2,2,2	0.59	0
5	EDO	В	203	-	3,3,3	0.36	0	2,2,2	0.97	0
5	EDO	G	203	-	3,3,3	0.34	0	2,2,2	0.70	0
3	MMV	E	202	-	27,27,27	0.99	2 (7%)	32,35,35	1.34	6 (18%)
3	MMV	С	202	-	27,27,27	1.17	2 (7%)	32,35,35	1.05	3 (9%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	EDO	D	203	-	-	1/1/1/1	-



Continued from previous page...

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAP	F	201	-	-	2/31/67/67	0/5/5/5
5	EDO	F	203	-	-	1/1/1/1	-
2	NAP	В	201	-	-	3/31/67/67	0/5/5/5
2	NAP	A	201	-	-	4/31/67/67	0/5/5/5
3	MMV	G	202	-	-	3/15/15/15	0/2/2/2
3	MMV	A	202	-	-	1/15/15/15	0/2/2/2
2	NAP	C	201	-	-	3/31/67/67	0/5/5/5
5	EDO	С	203	-	-	1/1/1/1	-
3	MMV	D	202	-	-	1/15/15/15	0/2/2/2
3	MMV	F	202	-	-	2/15/15/15	0/2/2/2
2	NAP	D	201	-	-	5/31/67/67	0/5/5/5
3	MMV	В	202	-	-	2/15/15/15	0/2/2/2
2	NAP	G	201	-	-	4/31/67/67	0/5/5/5
2	NAP	Е	201	-	-	4/31/67/67	0/5/5/5
5	EDO	Е	203	-	-	1/1/1/1	-
5	EDO	В	203	-	-	1/1/1/1	-
5	EDO	G	203	-	-	1/1/1/1	-
3	MMV	Е	202	-	-	2/15/15/15	0/2/2/2
3	MMV	С	202	-	-	0/15/15/15	0/2/2/2

The worst 5 of 24 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	Observed(Å)	Ideal(A)
3	С	202	MMV	C6-C5	-3.29	1.35	1.40
3	G	202	MMV	C6-C5	-2.96	1.35	1.40
3	В	202	MMV	C6-C5	-2.96	1.35	1.40
3	A	202	MMV	C6-C5	-2.93	1.36	1.40
3	A	202	MMV	C23-C24	2.91	1.57	1.50

The worst 5 of 32 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^o)$	$\operatorname{Ideal}({}^{o})$
3	Е	202	MMV	O15-C16-C17	3.63	120.58	115.89
3	F	202	MMV	C3-N4-C5	3.25	118.63	116.26
3	F	202	MMV	N7-C1-N2	3.03	121.11	117.03
3	A	202	MMV	O15-C16-C17	2.80	119.51	115.89
3	G	202	MMV	N7-C1-N2	2.76	120.75	117.03

There are no chirality outliers.



5 of 42 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	A	201	NAP	O4D-C1D-N1N-C2N
2	A	201	NAP	O4D-C1D-N1N-C6N
2	В	201	NAP	O4D-C1D-N1N-C2N
2	В	201	NAP	O4D-C1D-N1N-C6N
2	С	201	NAP	O4D-C1D-N1N-C2N


There are no ring outliers.

5 monomers are involved in 5 short contacts:


Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	F	201	NAP	1	0
5	F	203	EDO	1	0
2	D	201	NAP	1	0
2	G	201	NAP	1	0
5	Е	203	EDO	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.
















































Rings

Torsions



5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<rsrz></rsrz>	# RSRZ > 2	$\mathrm{OWAB}(\mathrm{\AA}^2)$	Q<0.9
1	A	159/171 (92%)	0.15	4 (2%) 58 57	18, 40, 70, 84	2 (1%)
1	В	160/171 (93%)	-0.73	0 100 100	13, 21, 35, 55	3 (1%)
1	С	159/171 (92%)	-0.41	0 100 100	16, 31, 48, 59	4 (2%)
1	D	159/171 (92%)	-0.75	0 100 100	10, 22, 34, 65	4 (2%)
1	E	159/171 (92%)	-0.40	0 100 100	17, 28, 49, 58	2 (1%)
1	F	159/171 (92%)	-0.55	1 (0%) 85 85	12, 25, 43, 63	4 (2%)
1	G	160/171 (93%)	-0.41	0 100 100	16, 33, 49, 67	2 (1%)
All	All	1115/1197 (93%)	-0.44	5 (0%) 89 88	10, 28, 54, 84	21 (1%)

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	72	ALA	3.7
1	A	73	LEU	3.4
1	F	138	VAL	3.2
1	A	71	ASP	3.1
1	A	74	PHE	2.2

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.



6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	${f B ext{-}factors}({f \AA}^2)$	Q<0.9
5	EDO	D	203	4/4	0.86	0.11	35,36,41,45	0
5	EDO	С	203	4/4	0.87	0.13	33,35,36,44	0
5	EDO	Е	203	4/4	0.87	0.14	36,38,42,43	0
5	EDO	В	203	4/4	0.88	0.11	33,33,33,38	0
5	EDO	G	203	4/4	0.88	0.11	40,42,43,44	0
4	CA	С	204	1/1	0.89	0.09	57,57,57,57	0
3	MMV	A	202	26/26	0.89	0.10	32,41,49,53	0
5	EDO	F	203	4/4	0.90	0.12	28,33,34,38	0
2	NAP	A	201	48/48	0.92	0.10	27,40,44,51	0
4	CA	A	203	1/1	0.95	0.06	59,59,59,59	0
3	MMV	Е	202	26/26	0.95	0.07	17,24,38,45	0
3	MMV	В	202	26/26	0.96	0.06	11,17,26,31	0
2	NAP	G	201	48/48	0.96	0.06	22,34,41,44	0
3	MMV	F	202	26/26	0.96	0.06	15,21,28,35	0
3	MMV	G	202	26/26	0.96	0.06	19,27,35,37	0
3	MMV	D	202	26/26	0.97	0.05	13,20,29,33	0
2	NAP	С	201	48/48	0.97	0.05	18,24,31,32	0
3	MMV	С	202	26/26	0.97	0.05	16,23,34,35	0
4	CA	В	204	1/1	0.98	0.05	37,37,37,37	0
2	NAP	D	201	48/48	0.98	0.04	14,19,23,28	0
2	NAP	Е	201	48/48	0.98	0.05	20,26,32,33	0
2	NAP	F	201	48/48	0.98	0.04	15,21,25,27	0
4	CA	В	205	1/1	0.99	0.03	30,30,30,30	0
2	NAP	В	201	48/48	0.99	0.04	12,18,21,25	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.



Electron density around MMV A 202: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive) Electron density around NAP A 201: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around MMV B 202:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAP G 201:


 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





















Electron density around MMV D 202:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAP C 201:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around MMV C 202:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAP D 201:

 $2 \text{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\text{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around NAP E 201:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NAP F 201:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











6.5 Other polymers (i)

There are no such residues in this entry.

