



Full wwPDB X-ray Structure Validation Report ⓘ

Nov 17, 2024 – 05:08 PM EST

PDB ID : 4K71
Title : Crystal structure of a high affinity Human Serum Albumin variant bound to the Neonatal Fc Receptor
Authors : Schmidt, M.M.; Townson, S.A.; Andreucci, A.; Dombrowski, C.; Erbe, D.V.; King, B.; Kovalchin, J.T.; Masci, A.; Murillo, A.; Schirmer, E.B.; Furfine, E.S.; Barnes, T.M.
Deposited on : 2013-04-16
Resolution : 2.40 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 2022.3.0, CSD as543be (2022)
Xtriage (Phenix) : 1.20.1
EDS : 3.0
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
CCP4 : 9.0.003 (Gargrove)
Density-Fitness : 1.0.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.40 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.





Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	164625	4642 (2.40-2.40)
Clashscore	180529	5218 (2.40-2.40)
Ramachandran outliers	177936	5158 (2.40-2.40)
Sidechain outliers	177891	5159 (2.40-2.40)
RSRZ outliers	164620	4642 (2.40-2.40)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	585	
1	D	585	
2	B	274	
2	E	274	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
3	C	99	 <p>% 94% 6%</p>
3	F	99	 <p>3% 92% 8%</p>

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 15259 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Serum albumin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	583	Total 4631	C 2923	N 784	O 882	S 42	48	0	0
1	D	579	Total 4597	C 2903	N 778	O 874	S 42	107	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	418	MET	VAL	engineered mutation	UNP P02768
A	420	ALA	THR	engineered mutation	UNP P02768
A	505	GLY	GLU	engineered mutation	UNP P02768
A	547	ALA	VAL	engineered mutation	UNP P02768
D	418	MET	VAL	engineered mutation	UNP P02768
D	420	ALA	THR	engineered mutation	UNP P02768
D	505	GLY	GLU	engineered mutation	UNP P02768
D	547	ALA	VAL	engineered mutation	UNP P02768

- Molecule 2 is a protein called IgG receptor FcRn large subunit p51.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	264	Total 2078	C 1329	N 358	O 383	S 8	23	0	0
2	E	264	Total 2078	C 1329	N 358	O 383	S 8	10	0	0

- Molecule 3 is a protein called Beta-2-microglobulin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	C	99	Total 829	C 528	N 140	O 158	S 3	23	0	0
3	F	99	Total 829	C 528	N 140	O 158	S 3	3	0	0

- Molecule 4 is SULFATE ION (three-letter code: SO4) (formula: O₄S).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	A	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	B	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	C	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		

Continued on next page...

Continued from previous page...

Mol	Chain	Residues	Atoms			ZeroOcc	AltConf
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	D	1	Total	O	S	0	0
			5	4	1		
4	E	1	Total	O	S	0	0
			5	4	1		
4	F	1	Total	O	S	0	0
			5	4	1		


- Molecule 5 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
5	A	39	Total	O	0	0
			39	39		
5	B	28	Total	O	0	0
			28	28		
5	C	7	Total	O	0	0
			7	7		
5	D	30	Total	O	0	0
			30	30		
5	E	12	Total	O	0	0
			12	12		
5	F	6	Total	O	0	0
			6	6		


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Serum albumin




- Molecule 1: Serum albumin



- Molecule 2: IgG receptor FcRn large subunit p51




- Molecule 2: IgG receptor FcRn large subunit p51







- Molecule 3: Beta-2-microglobulin



- Molecule 3: Beta-2-microglobulin



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	127.89Å 203.54Å 100.56Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	108.29 – 2.40 108.29 – 2.40	Depositor EDS
% Data completeness (in resolution range)	99.8 (108.29-2.40) 99.7 (108.29-2.40)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.96 (at 2.39Å)	Xtrriage
Refinement program	REFMAC 5.5.0109	Depositor
R, R_{free}	0.212 , 0.255 0.212 , 0.250	Depositor DCC
R_{free} test set	5144 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	47.6	Xtrriage
Anisotropy	0.234	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 47.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.95	EDS
Total number of atoms	15259	wwPDB-VP
Average B, all atoms (Å ²)	52.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.48% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: SO4

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.67	0/4721	0.70	1/6364 (0.0%)
1	D	0.61	0/4686	0.68	3/6317 (0.0%)
2	B	0.69	0/2143	0.73	0/2912
2	E	0.64	0/2143	0.72	0/2912
3	C	0.67	0/852	0.76	0/1152
3	F	0.66	1/852 (0.1%)	0.74	0/1152
All	All	0.65	1/15397 (0.0%)	0.71	4/20809 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
3	F	80	CYS	CB-SG	-5.34	1.73	1.81

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	144	ARG	NE-CZ-NH1	5.89	123.24	120.30
1	A	144	ARG	NE-CZ-NH1	5.63	123.11	120.30
1	D	144	ARG	NE-CZ-NH2	-5.54	117.53	120.30
1	D	209	ARG	NE-CZ-NH1	5.15	122.88	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within

the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4631	0	4554	24	0
1	D	4597	0	4519	20	0
2	B	2078	0	1989	10	0
2	E	2078	0	1989	9	0
3	C	829	0	794	3	0
3	F	829	0	794	2	0
4	A	35	0	0	1	0
4	B	10	0	0	0	0
4	C	10	0	0	0	0
4	D	30	0	0	0	0
4	E	5	0	0	0	0
4	F	5	0	0	0	0
5	A	39	0	0	1	0
5	B	28	0	0	0	0
5	C	7	0	0	0	0
5	D	30	0	0	0	0
5	E	12	0	0	0	0
5	F	6	0	0	1	0
All	All	15259	0	14639	64	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (64) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:B:5:LEU:HD23	2:B:167:LEU:HD21	1.69	0.75
1:D:331:LEU:HD13	1:D:350:ALA:HB2	1.70	0.73
1:A:81:ARG:NH2	1:A:89:ASP:OD1	2.27	0.67
2:B:98:LEU:HD13	2:B:163:LEU:HD23	1.78	0.66
3:F:31:HIS:ND1	5:F:205:HOH:O	2.28	0.66
1:A:498:VAL:HG23	2:B:42:ARG:NH1	2.11	0.66
1:A:305:LEU:HD11	1:A:337:ARG:HD2	1.82	0.62
1:A:331:LEU:HD13	1:A:350:ALA:HB2	1.83	0.61
1:D:418:MET:CE	1:D:460:LEU:HD11	2.31	0.60
1:A:214:TRP:CD1	1:A:343:VAL:HG11	2.35	0.60
2:B:106:PRO:HG2	2:B:122:LEU:HD13	1.86	0.58
1:D:23:VAL:HG12	1:D:43:VAL:HG22	1.86	0.56
1:A:498:VAL:HG23	2:B:42:ARG:HH12	1.70	0.55
1:D:20:LYS:HE3	1:D:47:THR:HG21	1.89	0.54

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:322:ALA:HB1	1:D:325:VAL:HB	1.90	0.53
1:D:498:VAL:HG13	2:E:42:ARG:HH12	1.72	0.53
1:A:303:PRO:O	1:A:337:ARG:NH2	2.37	0.53
1:D:418:MET:HE2	1:D:460:LEU:HD11	1.92	0.52
3:F:25:CYS:HB2	3:F:39:LEU:HD21	1.91	0.52
1:D:498:VAL:HG13	2:E:42:ARG:NH1	2.24	0.52
1:A:66:LEU:HD13	1:A:251:LEU:HD12	1.92	0.51
1:A:305:LEU:HD11	1:A:337:ARG:CD	2.40	0.51
1:A:194:ALA:HB1	1:A:455:VAL:HG13	1.92	0.51
1:D:418:MET:HE1	1:D:460:LEU:HD11	1.92	0.51
1:D:383:GLU:HB3	1:D:384:PRO:HD3	1.93	0.50
1:A:498:VAL:HG22	5:A:714:HOH:O	2.10	0.50
2:B:221:THR:HG23	2:B:222:GLY:O	2.12	0.50
1:D:30:TYR:CD2	1:D:74:LEU:HD11	2.47	0.50
1:A:420:ALA:HB3	1:A:421:PRO:HD3	1.95	0.48
2:B:11:LEU:HD13	2:B:94:LEU:HD13	1.95	0.48
1:A:151:ALA:HB3	1:A:152:PRO:HD3	1.96	0.48
2:B:5:LEU:HD12	2:B:6:SER:H	1.79	0.48
1:D:194:ALA:HB1	1:D:455:VAL:CG1	2.44	0.47
1:D:331:LEU:HD21	1:D:347:LEU:CD2	2.44	0.47
2:B:98:LEU:HD13	2:B:163:LEU:CD2	2.43	0.47
1:D:20:LYS:CE	1:D:47:THR:HG21	2.44	0.47
2:E:186:ALA:HB2	2:E:196:LEU:CD1	2.45	0.46
1:D:36:PHE:CZ	1:D:40:VAL:HG21	2.51	0.46
1:A:113:PRO:O	1:A:145:ARG:NH1	2.48	0.45
2:E:54:GLU:OE2	2:E:166:HIS:ND1	2.38	0.44
1:A:458:ASN:O	1:A:462:VAL:HG13	2.17	0.44
1:D:209:ARG:NH2	1:D:327:LEU:HD13	2.32	0.44
3:C:64:LEU:HD22	3:C:66:TYR:CD1	2.53	0.44
1:D:420:ALA:HB3	1:D:421:PRO:HD3	2.00	0.44
3:C:39:LEU:HD13	3:C:68:THR:HG22	2.00	0.44
1:A:482:VAL:HG23	4:A:606:SO4:O4	2.18	0.43
1:A:554:PHE:CD2	1:A:575:LEU:HD22	2.53	0.43
1:D:139:LEU:CD2	1:D:154:LEU:HG	2.48	0.43
2:E:60:TYR:CD1	2:E:60:TYR:C	2.92	0.43
2:E:106:PRO:HG3	2:E:122:LEU:HD22	2.01	0.43
1:A:373:VAL:HG13	1:A:374:PHE:CD1	2.54	0.43
3:C:64:LEU:HD23	3:C:65:LEU:N	2.34	0.43
2:E:71:LYS:HE3	2:E:94:LEU:HD23	2.01	0.43
2:E:127:TRP:O	2:E:138:SER:OG	2.37	0.43
2:E:167:LEU:HD23	2:E:174:LEU:CD1	2.50	0.42

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:107:ASP:OD2	1:A:110:PRO:HA	2.20	0.42
1:A:418:MET:HE3	1:A:460:LEU:HD11	2.02	0.41
1:A:322:ALA:HB1	1:A:325:VAL:HB	2.02	0.41
1:D:485:ARG:HB3	1:D:486:PRO:HD3	2.02	0.41
2:B:267:LEU:H	2:B:267:LEU:HD12	1.85	0.41
1:A:344:VAL:HG23	1:A:451:ASP:OD1	2.21	0.41
1:A:23:VAL:HG12	1:A:43:VAL:HG22	2.03	0.40
1:A:383:GLU:HB3	1:A:384:PRO:HD3	2.02	0.40
1:D:348:ARG:HG3	1:D:482:VAL:CG1	2.51	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [\(i\)](#)

5.3.1 Protein backbone [\(i\)](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	581/585 (99%)	572 (98%)	9 (2%)	0	100	100
1	D	575/585 (98%)	556 (97%)	16 (3%)	3 (0%)	25	38
2	B	262/274 (96%)	253 (97%)	8 (3%)	1 (0%)	30	44
2	E	262/274 (96%)	254 (97%)	7 (3%)	1 (0%)	30	44
3	C	97/99 (98%)	96 (99%)	1 (1%)	0	100	100
3	F	97/99 (98%)	96 (99%)	1 (1%)	0	100	100
All	All	1874/1916 (98%)	1827 (98%)	42 (2%)	5 (0%)	37	51

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	113	PRO
1	D	115	LEU
2	E	102	ASN

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
2	B	100	PRO
1	D	366	PRO

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	507/508 (100%)	502 (99%)	5 (1%)	73	86
1	D	503/508 (99%)	495 (98%)	8 (2%)	58	76
2	B	218/226 (96%)	215 (99%)	3 (1%)	62	79
2	E	218/226 (96%)	215 (99%)	3 (1%)	62	79
3	C	94/94 (100%)	93 (99%)	1 (1%)	70	84
3	F	94/94 (100%)	90 (96%)	4 (4%)	25	42
All	All	1634/1656 (99%)	1610 (98%)	24 (2%)	60	77

All (24) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	57	GLU
1	A	435	SER
1	A	460	LEU
1	A	501	GLU
1	A	570	GLU
2	B	10	HIS
2	B	187	ARG
2	B	196	LEU
3	C	70	PHE
1	D	104	GLN
1	D	138	TYR
1	D	208	GLU
1	D	232	SER
1	D	245	CYS
1	D	276	LYS
1	D	498	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	D	571	GLU
2	E	5	LEU
2	E	140	ARG
2	E	143	GLN
3	F	64	LEU
3	F	70	PHE
3	F	88	SER
3	F	99	MET

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

5.6 Ligand geometry [i](#)

19 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	$\# Z > 2$	Counts	RMSZ	$\# Z > 2$
4	SO4	A	603	-	4,4,4	0.21	0	6,6,6	0.29	0
4	SO4	B	302	-	4,4,4	0.30	0	6,6,6	0.20	0
4	SO4	C	102	-	4,4,4	0.26	0	6,6,6	0.36	0

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
4	SO4	D	604	-	4,4,4	0.24	0	6,6,6	0.54	0
4	SO4	F	101	-	4,4,4	0.31	0	6,6,6	0.37	0
4	SO4	A	605	-	4,4,4	0.28	0	6,6,6	0.73	0
4	SO4	D	606	-	4,4,4	0.34	0	6,6,6	0.33	0
4	SO4	A	604	-	4,4,4	0.25	0	6,6,6	0.18	0
4	SO4	D	603	-	4,4,4	0.29	0	6,6,6	0.40	0
4	SO4	A	602	-	4,4,4	0.31	0	6,6,6	0.20	0
4	SO4	A	601	-	4,4,4	0.29	0	6,6,6	0.20	0
4	SO4	A	607	-	4,4,4	0.27	0	6,6,6	0.48	0
4	SO4	E	301	-	4,4,4	0.32	0	6,6,6	0.47	0
4	SO4	B	301	-	4,4,4	0.32	0	6,6,6	0.23	0
4	SO4	D	602	-	4,4,4	0.34	0	6,6,6	0.29	0
4	SO4	A	606	-	4,4,4	0.34	0	6,6,6	0.67	0
4	SO4	D	605	-	4,4,4	0.28	0	6,6,6	0.20	0
4	SO4	C	101	-	4,4,4	0.32	0	6,6,6	0.18	0
4	SO4	D	601	-	4,4,4	0.31	0	6,6,6	0.14	0

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	A	606	SO4	1	0

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	583/585 (99%)	-0.13	10 (1%) 69 65	27, 45, 78, 109	16 (2%)
1	D	579/585 (98%)	0.36	42 (7%) 22 21	27, 55, 104, 139	31 (5%)
2	B	264/274 (96%)	-0.21	4 (1%) 71 68	28, 41, 63, 98	7 (2%)
2	E	264/274 (96%)	-0.13	6 (2%) 61 58	28, 44, 68, 110	3 (1%)
3	C	99/99 (100%)	-0.08	1 (1%) 79 76	29, 44, 69, 83	8 (8%)
3	F	99/99 (100%)	0.01	3 (3%) 52 49	34, 52, 79, 86	1 (1%)
All	All	1888/1916 (98%)	0.02	66 (3%) 47 44	27, 47, 87, 139	66 (3%)

All (66) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	565	GLU	6.9
1	D	569	ALA	6.3
1	D	115	LEU	4.9
1	D	566	THR	4.8
1	D	568	PHE	4.4
1	D	4	LYS	4.4
2	E	5	LEU	4.2
1	A	585	LEU	4.1
1	D	368	GLU	4.1
1	D	511	ALA	4.1
1	D	507	PHE	4.0
1	D	2	ALA	4.0
1	D	321	GLU	4.0
1	D	364	ALA	3.9
1	D	111	ASN	3.8
1	D	559	CYS	3.8
1	D	570	GLU	3.7
1	D	301	ASP	3.7
2	E	100	PRO	3.7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	585	LEU	3.6
1	D	441	PRO	3.6
1	D	504	ALA	3.6
1	D	116	VAL	3.5
2	E	101	ASP	3.4
1	D	313	LYS	3.4
1	D	573	LYS	3.3
1	D	513	ILE	3.3
1	D	314	ASP	3.3
1	A	115	LEU	3.1
1	A	303	PRO	3.1
3	F	1	ILE	3.0
2	B	100	PRO	3.0
1	D	110	PRO	2.9
3	C	1	ILE	2.9
1	D	119	GLU	2.9
1	A	114	ARG	2.8
1	D	515	THR	2.8
2	B	268	GLU	2.8
1	D	303	PRO	2.7
1	D	121	ASP	2.7
1	D	300	ALA	2.6
2	B	5	LEU	2.6
1	D	554	PHE	2.6
1	A	304	SER	2.5
1	D	317	LYS	2.5
1	D	555	VAL	2.5
2	E	221	THR	2.5
1	D	505	GLY	2.5
2	E	168	GLU	2.5
1	A	4	LYS	2.4
1	A	561	ALA	2.4
1	D	516	LEU	2.4
2	B	85	LYS	2.3
3	F	48	LYS	2.3
1	D	571	GLU	2.3
1	A	109	ASN	2.3
3	F	73	THR	2.2
1	D	113	PRO	2.2
1	A	519	LYS	2.2
1	D	567	CYS	2.2
1	A	515	THR	2.1

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	D	435	SER	2.1
1	D	305	LEU	2.1
1	D	365	ASP	2.1
2	E	248	HIS	2.1
1	D	514	CYS	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95th percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å ²)	Q<0.9
4	SO4	A	601	5/5	0.60	0.14	96,101,103,106	0
4	SO4	B	301	5/5	0.67	0.13	90,94,100,102	0
4	SO4	D	601	5/5	0.68	0.13	86,90,95,97	0
4	SO4	A	604	5/5	0.73	0.10	91,96,100,101	0
4	SO4	D	606	5/5	0.74	0.12	86,91,95,97	0
4	SO4	C	101	5/5	0.79	0.10	79,91,97,97	0
4	SO4	F	101	5/5	0.81	0.11	78,85,94,95	0
4	SO4	E	301	5/5	0.86	0.15	58,61,76,76	0
4	SO4	A	602	5/5	0.87	0.10	87,88,91,97	0
4	SO4	B	302	5/5	0.88	0.17	64,68,74,76	0
4	SO4	A	603	5/5	0.88	0.12	76,78,81,85	0
4	SO4	C	102	5/5	0.89	0.14	57,57,65,73	0
4	SO4	D	603	5/5	0.91	0.11	66,74,81,87	0
4	SO4	D	602	5/5	0.91	0.15	62,64,66,67	0
4	SO4	A	607	5/5	0.92	0.12	68,71,79,79	0
4	SO4	A	605	5/5	0.94	0.08	36,46,48,55	0
4	SO4	D	604	5/5	0.95	0.07	49,49,62,63	0
4	SO4	A	606	5/5	0.96	0.11	48,52,62,64	0

Continued on next page...

Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(\AA^2)	Q<0.9
4	SO4	D	605	5/5	0.97	0.07	46,57,59,67	0

6.5 Other polymers [i](#)

There are no such residues in this entry.