



# Full wwPDB X-ray Structure Validation Report ⓘ

Oct 9, 2023 – 11:58 AM EDT

PDB ID : 7K7D  
Title : Crystal structure of diphtheria toxin from crystals obtained at pH 6.0  
Authors : Lovell, S.; Kashipathy, M.M.; Battaile, K.P.; Rodnin, M.V.; Ladokhin, A.S.  
Deposited on : 2020-09-22  
Resolution : 2.10 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Xtriage (Phenix) : 1.13  
EDS : 2.35.1  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.35.1


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.10 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	5197 (2.10-2.10)
Clashscore	141614	5710 (2.10-2.10)
Ramachandran outliers	138981	5647 (2.10-2.10)
Sidechain outliers	138945	5648 (2.10-2.10)
RSRZ outliers	127900	5083 (2.10-2.10)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	538	 6% 87% 7% 6%
1	B	538	 6% 85% 9% 6%

## 2 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 7718 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Diphtheria toxin.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	507	3753	2384	629	728	12	0	0	0
1	B	505	3738	2371	630	725	12	0	0	0

There are 8 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	51	GLU	LYS	engineered mutation	UNP Q5PY51
A	148	LYS	GLU	engineered mutation	UNP Q5PY51
A	536	MET	-	cloning artifact	UNP Q5PY51
A	537	ALA	-	cloning artifact	UNP Q5PY51
B	51	GLU	LYS	engineered mutation	UNP Q5PY51
B	148	LYS	GLU	engineered mutation	UNP Q5PY51
B	536	MET	-	cloning artifact	UNP Q5PY51
B	537	ALA	-	cloning artifact	UNP Q5PY51


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	119	Total 119	O 119	0	0
2	B	108	Total 108	O 108	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Diphtheria toxin



- Molecule 1: Diphtheria toxin



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	69.44Å 69.61Å 73.12Å 117.90° 93.86° 97.93°	Depositor
Resolution (Å)	36.43 – 2.10 46.57 – 2.10	Depositor EDS
% Data completeness (in resolution range)	97.4 (36.43-2.10) 97.4 (46.57-2.10)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.15 (at 2.10Å)	Xtrriage
Refinement program	PHENIX 1.14rc3_3208	Depositor
R, $R_{free}$	0.210 , 0.258 0.215 , 0.258	Depositor DCC
$R_{free}$ test set	3177 reflections (4.72%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	41.2	Xtrriage
Anisotropy	0.465	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.32 , 48.6	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	7718	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	56.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 10.44% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.50	0/3828	0.59	0/5206
1	B	0.48	0/3813	0.58	0/5185
All	All	0.49	0/7641	0.59	0/10391

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	3753	0	3583	19	0
1	B	3738	0	3578	27	0
2	A	119	0	0	3	0
2	B	108	0	0	1	0
All	All	7718	0	7161	44	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 3.

All (44) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:238:VAL:HG22	1:A:239:SER:H	1.52	0.73

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:247:LEU:HD11	1:B:343:ALA:HB2	1.85	0.59
1:B:440:LYS:O	1:B:493:ARG:HA	2.03	0.59
1:B:510:GLY:HA3	1:B:526:LYS:HE3	1.86	0.58
1:B:238:VAL:HG22	1:B:239:SER:H	1.69	0.58
1:A:351:VAL:HG12	1:A:351:VAL:O	2.06	0.55
1:B:238:VAL:HG22	1:B:239:SER:N	2.21	0.55
1:A:474:LYS:O	1:B:73:LEU:HD23	2.08	0.53
1:A:49:ASP:OD1	1:A:103:LYS:NZ	2.33	0.53
1:A:406:ILE:HB	1:A:533:ILE:HD13	1.88	0.53
1:B:246:TYR:HD2	1:B:346:LEU:HD11	1.76	0.51
1:B:251:HIS:NE2	1:B:255:LEU:HD22	2.27	0.49
1:A:523:VAL:HG22	1:A:524:ASN:N	2.28	0.49
1:B:94:LEU:HD11	1:B:136:LEU:HD13	1.96	0.47
1:A:95:LYS:HA	2:A:625:HOH:O	2.13	0.47
1:B:347:VAL:O	1:B:347:VAL:HG12	2.14	0.47
1:A:52:GLY:HA2	1:A:153:TRP:CD1	2.51	0.46
1:A:182:MET:HE2	2:A:716:HOH:O	2.16	0.46
1:B:254:ALA:O	1:B:260:LEU:HD12	2.16	0.46
1:B:407:ARG:HA	1:B:534:LYS:O	2.15	0.46
1:B:213:THR:O	1:B:217:ILE:HG13	2.16	0.45
1:A:238:VAL:HG22	1:A:239:SER:N	2.27	0.45
1:B:349:GLU:H	1:B:350:LEU:HD23	1.82	0.45
1:A:315:GLY:HA3	1:A:323:HIS:CD2	2.52	0.45
1:B:440:LYS:HE2	1:B:499:ILE:HG13	1.99	0.44
1:B:281:TRP:O	1:B:285:VAL:HG23	2.17	0.44
1:A:28:VAL:HG11	1:A:166:ASN:ND2	2.33	0.44
1:B:94:LEU:HB2	1:B:138:LEU:HD21	2.00	0.44
1:A:352:ASP:C	1:A:354:GLY:N	2.71	0.43
1:B:526:LYS:HE2	2:B:601:HOH:O	2.18	0.43
1:B:315:GLY:HA3	1:B:323:HIS:CD2	2.54	0.43
1:B:346:LEU:HB2	1:B:347:VAL:HG23	2.01	0.42
1:A:85:TYR:CG	1:A:133:ARG:HD2	2.53	0.42
1:B:108:LEU:HB3	1:B:117:GLN:OE1	2.20	0.42
1:A:504:ILE:HG23	1:A:504:ILE:O	2.19	0.42
1:B:228:ASN:O	1:B:231:SER:HB2	2.19	0.42
1:B:246:TYR:CD2	1:B:346:LEU:HD11	2.54	0.41
1:B:351:VAL:O	1:B:351:VAL:HG23	2.20	0.41
1:A:200:SER:O	1:A:201:CYS:C	2.59	0.41
1:A:250:PHE:CE1	1:A:342:GLN:HG3	2.55	0.41
1:A:387:GLN:HG2	2:A:604:HOH:O	2.19	0.41
1:A:476:PRO:HD3	1:B:73:LEU:HD22	2.01	0.41

*Continued on next page...*

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:283:VAL:O	1:B:287:GLN:HG3	2.21	0.40
1:B:344:ILE:N	1:B:345:PRO:HD2	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	497/538 (92%)	475 (96%)	19 (4%)	3 (1%)	25	21
1	B	495/538 (92%)	473 (96%)	17 (3%)	5 (1%)	15	11
All	All	992/1076 (92%)	948 (96%)	36 (4%)	8 (1%)	19	15

All (8) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	B	201	CYS
1	A	235	ASN
1	B	223	HIS
1	B	68	ASP
1	A	68	ASP
1	A	438	PRO
1	B	438	PRO
1	B	107	GLY

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was



analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	388/454 (86%)	381 (98%)	7 (2%)	59	65
1	B	389/454 (86%)	383 (98%)	6 (2%)	65	71
All	All	777/908 (86%)	764 (98%)	13 (2%)	60	67

All (13) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	18	SER
1	A	219	SER
1	A	352	ASP
1	A	446	SER
1	A	451	SER
1	A	495	SER
1	A	516	LYS
1	B	9	SER
1	B	30	SER
1	B	133	ARG
1	B	237	THR
1	B	442	ASP
1	B	516	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	507/538 (94%)	0.43	31 (6%) 21 26	33, 54, 90, 124	0
1	B	505/538 (93%)	0.39	31 (6%) 21 26	33, 52, 87, 110	0
All	All	1012/1076 (94%)	0.41	62 (6%) 21 26	33, 53, 89, 124	0

All (62) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	228	ASN	5.8
1	A	123	PHE	5.5
1	A	124	ILE	5.1
1	B	124	ILE	4.9
1	B	108	LEU	4.7
1	A	346	LEU	4.6
1	B	123	PHE	4.5
1	B	233	SER	4.4
1	A	102	ILE	4.4
1	A	238	VAL	3.9
1	B	226	ILE	3.8
1	A	111	THR	3.8
1	B	350	LEU	3.8
1	A	226	ILE	3.6
1	B	102	ILE	3.3
1	A	120	THR	3.3
1	A	347	VAL	3.3
1	B	111	THR	3.2
1	A	110	LEU	3.2
1	A	225	PRO	3.1
1	A	122	GLU	3.1
1	A	344	ILE	3.1
1	A	106	LEU	3.0
1	B	126	ARG	3.0

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	B	422	ALA	3.0
1	A	93	ALA	2.9
1	A	241	GLU	2.9
1	A	233	SER	2.9
1	B	479	VAL	2.8
1	B	420	ILE	2.8
1	B	223	HIS	2.8
1	B	346	LEU	2.7
1	B	396	VAL	2.6
1	A	485	ALA	2.5
1	B	395	ALA	2.5
1	A	92	LEU	2.4
1	B	92	LEU	2.4
1	A	234	PRO	2.4
1	B	120	THR	2.4
1	A	114	LEU	2.3
1	A	350	LEU	2.3
1	B	234	PRO	2.3
1	A	134	VAL	2.3
1	B	421	THR	2.3
1	B	118	VAL	2.3
1	A	118	VAL	2.3
1	A	126	ARG	2.3
1	B	397	SER	2.3
1	B	222	GLU	2.2
1	B	483	VAL	2.2
1	A	94	LEU	2.1
1	B	255	LEU	2.1
1	B	349	GLU	2.1
1	B	485	ALA	2.1
1	B	394	TYR	2.1
1	A	422	ALA	2.1
1	B	225	PRO	2.1
1	A	388	PRO	2.1
1	B	398	TRP	2.1
1	A	420	ILE	2.1
1	B	119	GLY	2.1
1	A	125	LYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

### 6.4 Ligands [i](#)

There are no ligands in this entry.

### 6.5 Other polymers [i](#)

There are no such residues in this entry.