



# Full wwPDB X-ray Structure Validation Report ⓘ

Jan 15, 2019 – 10:41 PM EST

PDB ID : 5K8A  
Title : NIST FAB  
Authors : Gallagher, D.T.; Galvin, C.V.; Karageorgos, I.; Marino, J.P.  
Deposited on : 2016-05-27  
Resolution : 2.00 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.7.3 (157068), CSD as539be (2018)  
Xtriage (Phenix) : 1.13  
EDS : rb-20031633  
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)  
Refmac : 5.8.0158  
CCP4 : 7.0 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : rb-20031633


# 1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 2.00 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	111664	7193 (2.00-2.00)
Clashscore	122126	8267 (2.00-2.00)
Ramachandran outliers	120053	8166 (2.00-2.00)
Sidechain outliers	120020	8165 (2.00-2.00)
RSRZ outliers	108989	7011 (2.00-2.00)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	213	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">11%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="text-align: left;">92%</div> <div style="text-align: right;">8% •</div> </div>
1	E	213	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">10%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="text-align: left;">87%</div> <div style="text-align: right;">12% •</div> </div>
1	L	213	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">8%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="text-align: left;">95%</div> <div style="text-align: right;">•</div> </div>
1	M	213	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">9%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="text-align: left;">95%</div> <div style="text-align: right;">5%</div> </div>
2	B	227	<div style="display: flex; justify-content: space-between; align-items: center;"> <div style="text-align: right;">8%</div> <div style="width: 100%; height: 15px; background: linear-gradient(to right, red, orange, yellow, green, grey);"></div> <div style="text-align: left;">91%</div> <div style="text-align: right;">7% •</div> </div>

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Length	Quality of chain
2	F	227	 <p>7% 92% 6% ••</p>
2	H	227	 <p>9% 92% 6% •</p>
2	V	227	 <p>12% 86% 11% •</p>

## 2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 13629 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called humanized Fab Lite chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	L	213	Total 1626	C 1020	N 270	O 329	S 7	0	0	0
1	M	213	Total 1626	C 1020	N 270	O 329	S 7	0	0	0
1	A	213	Total 1626	C 1020	N 270	O 329	S 7	0	0	0
1	E	213	Total 1626	C 1020	N 270	O 329	S 7	0	0	0

- Molecule 2 is a protein called Heavy chain.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	H	223	Total 1690	C 1074	N 279	O 329	S 8	0	0	0
2	V	223	Total 1690	C 1074	N 279	O 329	S 8	0	0	0
2	B	223	Total 1690	C 1074	N 279	O 329	S 8	0	0	0
2	F	223	Total 1690	C 1074	N 279	O 329	S 8	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	L	36	Total 36	O 36	0	0
3	H	52	Total 52	O 52	0	0
3	M	38	Total 38	O 38	0	0
3	V	54	Total 54	O 54	0	0

*Continued on next page...*


*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Residues</b>	<b>Atoms</b>		<b>ZeroOcc</b>	<b>AltConf</b>
3	A	30	Total 30	O 30	0	0
3	B	56	Total 56	O 56	0	0
3	E	47	Total 47	O 47	0	0
3	F	52	Total 52	O 52	0	0


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: humanized Fab Lite chain




- Molecule 1: humanized Fab Lite chain




- Molecule 1: humanized Fab Lite chain




- Molecule 1: humanized Fab Lite chain






- Molecule 2: Heavy chain






HIS

- Molecule 2: Heavy chain




- Molecule 2: Heavy chain



HIS

- Molecule 2: Heavy chain



## 4 Data and refinement statistics i

Property	Value	Source
Space group	I 2 2 2	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	149.16Å 149.20Å 195.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	19.87 – 2.00 19.87 – 2.00	Depositor EDS
% Data completeness (in resolution range)	99.8 (19.87-2.00) 99.7 (19.87-2.00)	Depositor EDS
$R_{merge}$	0.08	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	2.67 (at 2.01Å)	Xtrriage
Refinement program	REFMAC 5.8.0135	Depositor
R, $R_{free}$	0.164 , 0.189 0.202 , 0.225	Depositor DCC
$R_{free}$ test set	7286 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	40.5	Xtrriage
Anisotropy	0.091	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.34 , 38.7	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.53$ , $\langle L^2 \rangle = 0.37$	Xtrriage
Estimated twinning fraction	0.000 for -k,-h,-l	Xtrriage
Reported twinning fraction	0.670 for H, K, L 0.330 for K, H, -L	Depositor
Outliers	0 of 145860 reflections	Xtrriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	13629	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	44.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 3.38% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



## 5 Model quality

### 5.1 Standard geometry

Bond lengths and bond angles in the following residue types are not validated in this section: PCA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.58	0/1663	0.70	0/2255
1	E	0.61	0/1663	0.76	1/2255 (0.0%)
1	L	0.59	0/1663	0.77	3/2255 (0.1%)
1	M	0.57	0/1663	0.71	0/2255
2	B	0.60	0/1727	0.74	0/2362
2	F	0.64	0/1727	0.78	2/2362 (0.1%)
2	H	0.58	0/1727	0.75	1/2362 (0.0%)
2	V	0.65	0/1727	0.76	0/2362
All	All	0.60	0/13560	0.75	7/18468 (0.0%)

There are no bond length outliers.

All (7) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	L	28	ARG	NE-CZ-NH2	-8.65	115.97	120.30
2	F	196	LEU	CA-CB-CG	7.10	131.63	115.30
1	L	46	LEU	CA-CB-CG	6.80	130.94	115.30
1	E	60	ARG	NE-CZ-NH2	-6.07	117.27	120.30
1	L	28	ARG	NE-CZ-NH1	5.62	123.11	120.30
2	F	68	ARG	NE-CZ-NH1	5.51	123.05	120.30
2	H	68	ARG	NE-CZ-NH2	-5.37	117.61	120.30

There are no chirality outliers.

There are no planarity outliers.

### 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen

atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1626	0	1574	10	0
1	E	1626	0	1574	12	0
1	L	1626	0	1574	5	0
1	M	1626	0	1574	6	0
2	B	1690	0	1670	5	0
2	F	1690	0	1670	5	0
2	H	1690	0	1670	3	0
2	V	1690	0	1670	14	0
3	A	30	0	0	1	0
3	B	56	0	0	0	0
3	E	47	0	0	0	0
3	F	52	0	0	0	0
3	H	52	0	0	0	0
3	L	36	0	0	0	0
3	M	38	0	0	0	0
3	V	54	0	0	0	0
All	All	13629	0	12976	53	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (53) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:L:26:SER:HB3	1:M:5:THR:HG21	1.64	0.80
1:A:38:LYS:HB2	1:A:41:LYS:HD2	1.65	0.78
1:L:5:THR:HG21	1:M:26:SER:HB3	1.70	0.72
2:B:206:ASN:ND2	2:B:213:LYS:HG2	2.20	0.56
2:V:217:ARG:HB3	1:E:151:ASN:HD22	1.70	0.56
1:A:39:PRO:HB3	1:A:164:GLU:HG3	1.87	0.56
2:V:88:ASP:HB2	2:V:89:PRO:HD2	1.88	0.56
1:A:5:THR:HG23	1:E:5:THR:HG23	1.88	0.55
2:F:30:SER:OG	2:F:75:THR:HG21	2.09	0.53
1:M:11:LEU:HD11	1:M:19:VAL:HG13	1.91	0.53
1:A:119:PRO:HD3	1:A:131:VAL:HG22	1.89	0.53
2:V:35:SER:HB2	2:V:54:TRP:CE3	2.44	0.52
1:A:28:ARG:NH2	3:A:301:HOH:O	2.44	0.50
1:E:18:ARG:HH11	1:E:18:ARG:HB2	1.76	0.50
1:E:150:ASP:HA	1:E:190:VAL:HG22	1.95	0.48

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
2:F:88:ASP:HB2	2:F:89:PRO:HD2	1.95	0.48
1:E:141:ARG:HH21	1:E:162:VAL:HG11	1.77	0.48
2:B:35:SER:HB2	2:B:54:TRP:CE3	2.49	0.47
1:M:48:TYR:O	1:M:52:LYS:HB2	2.14	0.47
1:E:107:ARG:HH11	1:E:108:THR:HG23	1.79	0.47
1:A:107:ARG:HG2	1:A:170:SER:HB2	1.97	0.46
2:V:102:ILE:O	2:V:105:PHE:CE1	2.68	0.45
1:E:185:TYR:CZ	1:E:210:ARG:HG3	2.51	0.45
1:E:192:ALA:HB2	1:E:207:SER:HB3	1.99	0.45
2:F:12:VAL:HG23	2:F:118:VAL:HG22	1.99	0.45
2:H:35:SER:HB2	2:H:54:TRP:CE3	2.52	0.45
2:V:2:VAL:HG13	2:V:27:PHE:CD2	2.52	0.45
2:V:157:VAL:CG1	2:V:185:LEU:HD21	2.48	0.43
2:V:41:GLN:HB2	2:V:47:LEU:HD23	1.99	0.43
2:V:32:ALA:HA	2:V:55:TRP:CD2	2.53	0.43
1:A:212:GLU:OE2	2:B:136:LYS:HE3	2.18	0.43
2:V:63:PRO:HA	2:V:66:LYS:HB2	2.00	0.43
1:A:90:GLY:HA2	1:A:95:PHE:CD1	2.54	0.43
1:L:5:THR:HG21	1:M:26:SER:CB	2.43	0.42
1:M:190:VAL:HG12	1:M:209:ASN:OD1	2.19	0.42
2:V:158:THR:OG1	2:V:206:ASN:HB3	2.20	0.42
2:H:138:THR:HA	2:H:142:THR:O	2.20	0.42
1:L:79:PRO:HA	1:L:105:ILE:HG13	2.01	0.42
2:B:12:VAL:O	2:B:118:VAL:HA	2.20	0.42
2:V:12:VAL:O	2:V:118:VAL:HA	2.19	0.41
2:V:29:LEU:HB3	2:V:75:THR:HG22	2.02	0.41
1:L:46:LEU:HD11	1:L:61:PHE:CD2	2.54	0.41
1:A:123:GLN:HG2	1:A:128:THR:HG23	2.01	0.41
2:B:92:THR:HG23	2:B:117:THR:HA	2.03	0.41
1:E:4:MET:HG2	1:E:23:CYS:SG	2.60	0.41
2:F:124:LYS:NZ	2:F:151:ASP:HB3	2.35	0.41
1:A:22:THR:HG22	1:A:71:THR:HG22	2.02	0.41
2:F:12:VAL:O	2:F:118:VAL:HA	2.20	0.41
1:E:19:VAL:HB	1:E:74:ILE:HB	2.02	0.41
2:V:40:ARG:HB3	2:V:50:LEU:HD11	2.02	0.41
2:H:12:VAL:O	2:H:118:VAL:HA	2.20	0.40
1:E:160:GLU:HA	1:E:175:SER:O	2.21	0.40
2:V:217:ARG:HD3	1:E:151:ASN:ND2	2.36	0.40

There are no symmetry-related clashes.

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	211/213 (99%)	203 (96%)	8 (4%)	0	100	100
1	E	211/213 (99%)	207 (98%)	3 (1%)	1 (0%)	31	25
1	L	211/213 (99%)	205 (97%)	6 (3%)	0	100	100
1	M	211/213 (99%)	206 (98%)	5 (2%)	0	100	100
2	B	221/227 (97%)	212 (96%)	9 (4%)	0	100	100
2	F	221/227 (97%)	215 (97%)	6 (3%)	0	100	100
2	H	221/227 (97%)	216 (98%)	5 (2%)	0	100	100
2	V	221/227 (97%)	213 (96%)	8 (4%)	0	100	100
All	All	1728/1760 (98%)	1677 (97%)	50 (3%)	1 (0%)	53	51

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	E	167	SER

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	185/185 (100%)	182 (98%)	3 (2%)	65	70
1	E	185/185 (100%)	177 (96%)	8 (4%)	32	28
1	L	185/185 (100%)	181 (98%)	4 (2%)	55	58
1	M	185/185 (100%)	181 (98%)	4 (2%)	55	58

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
2	B	194/198 (98%)	185 (95%)	9 (5%)	29	25
2	F	194/198 (98%)	187 (96%)	7 (4%)	38	36
2	H	194/198 (98%)	187 (96%)	7 (4%)	38	36
2	V	194/198 (98%)	191 (98%)	3 (2%)	67	72
All	All	1516/1532 (99%)	1471 (97%)	45 (3%)	44	44

All (45) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	L	24	SER
1	L	46	LEU
1	L	142	GLU
1	L	201	SER
2	H	65	LEU
2	H	76	SER
2	H	88	ASP
2	H	102	ILE
2	H	200	THR
2	H	212	THR
2	H	222	SER
1	M	5	THR
1	M	64	SER
1	M	72	LEU
1	M	144	LYS
2	V	122	SER
2	V	142	THR
2	V	157	VAL
1	A	32	MET
1	A	38	LYS
1	A	107	ARG
2	B	5	ARG
2	B	25	SER
2	B	65	LEU
2	B	83	LYS
2	B	102	ILE
2	B	134	SER
2	B	157	VAL
2	B	196	LEU
2	B	206	ASN
1	E	9	SER
1	E	38	LYS

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type
1	E	104	GLU
1	E	128	THR
1	E	169	ASP
1	E	179	THR
1	E	190	VAL
1	E	207	SER
2	F	67	ASP
2	F	88	ASP
2	F	101	MET
2	F	122	SER
2	F	196	LEU
2	F	216	LYS
2	F	217	ARG

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (5) such sidechains are listed below:

Mol	Chain	Res	Type
1	M	198	GLN
1	A	36	GLN
1	A	198	GLN
1	E	151	ASN
2	F	112	GLN

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

4 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	PCA	B	1	2	8,8,9	0.68	0	9,10,12	1.10	1 (11%)
2	PCA	F	1	2	8,8,9	0.46	0	9,10,12	1.55	3 (33%)
2	PCA	H	1	2	8,8,9	0.42	0	9,10,12	1.25	1 (11%)
2	PCA	V	1	2	8,8,9	0.48	0	9,10,12	1.40	1 (11%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	PCA	B	1	2	-	0/0/11/13	0/1/1/1
2	PCA	F	1	2	-	0/0/11/13	0/1/1/1
2	PCA	H	1	2	-	0/0/11/13	0/1/1/1
2	PCA	V	1	2	-	0/0/11/13	0/1/1/1

There are no bond length outliers.

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	F	1	PCA	O-C-CA	-2.94	118.24	125.09
2	V	1	PCA	CB-CA-C	-2.93	108.67	112.70
2	F	1	PCA	OE-CD-CG	-2.50	122.32	126.83
2	H	1	PCA	O-C-CA	-2.17	120.03	125.09
2	B	1	PCA	OE-CD-CG	-2.12	123.00	126.83
2	F	1	PCA	CB-CA-C	-2.02	109.92	112.70

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry [i](#)

There are no ligands in this entry.

## 5.7 Other polymers [i](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.



## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	213/213 (100%)	0.80	23 (10%) 6 5	33, 46, 61, 111	0
1	E	213/213 (100%)	0.75	22 (10%) 6 6	30, 43, 63, 116	0
1	L	213/213 (100%)	0.75	18 (8%) 11 10	31, 44, 63, 113	0
1	M	213/213 (100%)	0.71	19 (8%) 9 9	29, 46, 67, 101	0
2	B	222/227 (97%)	0.69	19 (8%) 10 10	30, 40, 66, 113	0
2	F	222/227 (97%)	0.50	15 (6%) 17 17	30, 39, 55, 124	0
2	H	222/227 (97%)	0.64	20 (9%) 9 9	32, 43, 57, 112	0
2	V	222/227 (97%)	0.87	28 (12%) 3 3	30, 40, 77, 131	0
All	All	1740/1760 (98%)	0.71	164 (9%) 8 8	29, 42, 63, 131	0

All (164) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	L	213	CYS	14.0
1	E	213	CYS	12.9
1	A	213	CYS	11.8
2	F	223	CYS	11.6
2	B	223	CYS	11.2
2	H	223	CYS	11.1
2	V	137	SER	10.7
1	M	213	CYS	10.0
2	V	136	LYS	9.6
2	V	222	SER	8.9
2	V	140	GLY	8.7
2	V	138	THR	8.6
2	B	222	SER	8.2
2	V	223	CYS	7.9
2	V	139	SER	6.4
2	H	222	SER	5.7

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	H	36	VAL	5.6
2	V	135	SER	5.4
2	V	198	THR	5.1
2	F	222	SER	5.1
1	L	40	GLY	4.7
2	B	197	GLY	4.6
1	E	156	GLY	4.5
1	A	201	SER	4.3
1	L	1	ASP	4.3
1	E	40	GLY	4.1
1	A	156	GLY	4.1
2	V	221	LYS	4.0
2	B	198	THR	4.0
2	B	139	SER	4.0
2	B	221	LYS	4.0
1	M	155	SER	4.0
1	M	156	GLY	3.9
2	B	140	GLY	3.9
1	E	201	SER	3.8
1	A	40	GLY	3.7
1	E	212	GLU	3.7
2	B	138	THR	3.7
2	F	36	VAL	3.6
2	B	135	SER	3.6
1	M	40	GLY	3.5
1	M	201	SER	3.5
2	V	66	LYS	3.5
1	L	167	SER	3.5
1	M	204	VAL	3.4
1	M	1	ASP	3.4
1	E	125	LYS	3.4
1	M	211	GLY	3.4
2	B	102	ILE	3.4
1	A	155	SER	3.4
2	V	197	GLY	3.3
2	V	102	ILE	3.3
2	V	51	ALA	3.3
2	B	36	VAL	3.3
2	F	51	ALA	3.3
2	F	221	LYS	3.3
2	H	38	TRP	3.2
1	A	88	PHE	3.2

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	H	147	CYS	3.2
1	E	82	PHE	3.2
1	A	39	PRO	3.2
1	E	135	LEU	3.1
1	M	39	PRO	3.1
1	E	134	LEU	3.1
2	V	145	LEU	3.1
1	L	202	SER	3.0
2	V	55	TRP	3.0
1	L	46	LEU	3.0
1	A	85	TYR	3.0
2	V	112	GLN	3.0
1	M	202	SER	2.9
2	H	169	GLY	2.9
1	A	153	LEU	2.9
2	F	140	GLY	2.9
2	H	37	GLY	2.9
1	A	167	SER	2.8
1	A	1	ASP	2.8
1	A	142	GLU	2.8
2	V	45	LYS	2.8
1	A	23	CYS	2.8
2	H	44	GLY	2.8
1	A	82	PHE	2.8
1	L	125	LYS	2.8
1	A	168	LYS	2.8
2	B	38	TRP	2.8
2	F	147	CYS	2.8
1	E	35	TYR	2.7
1	L	34	TRP	2.7
2	B	51	ALA	2.7
2	F	37	GLY	2.7
2	H	221	LYS	2.7
1	M	152	ALA	2.7
2	V	116	VAL	2.7
2	F	50	LEU	2.7
1	A	18	ARG	2.7
2	H	51	ALA	2.6
1	L	156	GLY	2.6
1	M	85	TYR	2.6
1	L	201	SER	2.6
1	A	72	LEU	2.6

*Continued on next page...*

*Continued from previous page...*

<b>Mol</b>	<b>Chain</b>	<b>Res</b>	<b>Type</b>	<b>RSRZ</b>
2	V	211	ASN	2.6
1	M	190	VAL	2.6
1	L	212	GLU	2.5
2	V	67	ASP	2.5
2	H	53	ILE	2.5
2	F	39	ILE	2.5
1	L	132	VAL	2.5
1	A	104	GLU	2.5
1	E	202	SER	2.5
2	F	44	GLY	2.5
1	L	72	LEU	2.5
1	E	146	GLN	2.4
1	L	122	GLU	2.4
1	A	164	GLU	2.4
2	V	36	VAL	2.4
1	E	164	GLU	2.4
2	B	50	LEU	2.4
1	L	23	CYS	2.4
2	H	112	GLN	2.4
1	E	155	SER	2.4
1	E	132	VAL	2.4
1	M	153	LEU	2.3
1	M	198	GLN	2.3
2	B	39	ILE	2.3
2	V	65	LEU	2.3
1	E	142	GLU	2.3
1	E	153	LEU	2.3
1	A	76	SER	2.3
2	V	49	TRP	2.3
1	E	169	ASP	2.3
2	H	211	ASN	2.3
1	E	211	GLY	2.2
1	M	164	GLU	2.2
2	V	147	CYS	2.2
1	M	144	LYS	2.2
2	H	149	VAL	2.2
2	F	66	LYS	2.2
2	H	217	ARG	2.2
2	V	38	TRP	2.2
2	H	189	VAL	2.2
2	F	148	LEU	2.2
1	A	202	SER	2.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
2	B	142	THR	2.1
2	V	39	ILE	2.1
2	F	38	TRP	2.1
1	L	144	LYS	2.1
1	L	211	GLY	2.1
1	E	180	LEU	2.1
1	M	82	PHE	2.1
2	F	45	LYS	2.1
2	H	39	ILE	2.1
2	H	76	SER	2.1
2	B	168	SER	2.1
2	V	146	GLY	2.1
1	E	85	TYR	2.1
2	B	136	LYS	2.1
2	H	82	LEU	2.1
1	M	200	LEU	2.0
1	A	46	LEU	2.0
1	L	104	GLU	2.0
1	A	35	TYR	2.0
2	B	211	ASN	2.0
2	H	195	SER	2.0
1	E	87	CYS	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	PCA	B	1	8/9	0.81	0.31	48,52,61,64	0
2	PCA	F	1	8/9	0.83	0.29	48,60,70,73	0
2	PCA	V	1	8/9	0.90	0.27	46,54,58,61	0
2	PCA	H	1	8/9	0.95	0.22	52,63,69,73	0

## 6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

## 6.4 Ligands

There are no ligands in this entry.

## 6.5 Other polymers

There are no such residues in this entry.