



# Full wwPDB X-ray Structure Validation Report i

May 16, 2020 – 03:55 pm BST

PDB ID : 5K91  
Title : Crystal structure of dimeric chlorite dismutase from Cyanothec sp. PCC7425 in complex with fluoride  
Authors : Puehringer, D.; Schaffner, I.; Mlynek, G.; Obinger, C.; Djinovic-Carugo, K.  
Deposited on : 2016-05-31  
Resolution : 1.18 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the i symbol.

---

The following versions of software and data (see [references](#) i) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.11  
buster-report : 1.1.7 (2018)  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.11


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

## X-RAY DIFFRACTION


The reported resolution of this entry is 1.18 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1123 (1.20-1.16)
Clashscore	141614	1182 (1.20-1.16)
Ramachandran outliers	138981	1134 (1.20-1.16)
Sidechain outliers	138945	1134 (1.20-1.16)
RSRZ outliers	127900	1102 (1.20-1.16)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.



## 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 13105 atoms, of which 6004 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Chlorite dismutase.

Mol	Chain	Residues	Atoms						ZeroOcc	AltConf	Trace	
1	A	174	Total	C 2922	H 937	N 1457	O 265	S 258	5	0	6	0
1	B	175	Total	C 2961	H 948	N 1474	O 269	S 265	5	0	10	0
1	C	175	Total	C 2928	H 938	N 1458	O 266	S 260	6	0	6	0
1	D	175	Total	C 2957	H 947	N 1469	O 269	S 267	5	0	10	0

There are 28 discrepancies between the modelled and reference sequences:


Chain	Residue	Modelled	Actual	Comment	Reference
A	-5	GLY	-	expression tag	UNP B8HNS6
A	-4	PRO	-	expression tag	UNP B8HNS6
A	-3	GLY	-	expression tag	UNP B8HNS6
A	-2	TYR	-	expression tag	UNP B8HNS6
A	-1	GLN	-	expression tag	UNP B8HNS6
A	0	ASP	-	expression tag	UNP B8HNS6
A	1	PRO	-	expression tag	UNP B8HNS6
B	-5	GLY	-	expression tag	UNP B8HNS6
B	-4	PRO	-	expression tag	UNP B8HNS6
B	-3	GLY	-	expression tag	UNP B8HNS6
B	-2	TYR	-	expression tag	UNP B8HNS6
B	-1	GLN	-	expression tag	UNP B8HNS6
B	0	ASP	-	expression tag	UNP B8HNS6
B	1	PRO	-	expression tag	UNP B8HNS6
C	-5	GLY	-	expression tag	UNP B8HNS6
C	-4	PRO	-	expression tag	UNP B8HNS6
C	-3	GLY	-	expression tag	UNP B8HNS6
C	-2	TYR	-	expression tag	UNP B8HNS6
C	-1	GLN	-	expression tag	UNP B8HNS6
C	0	ASP	-	expression tag	UNP B8HNS6
C	1	PRO	-	expression tag	UNP B8HNS6

*Continued on next page...*

*Continued from previous page...*

Chain	Residue	Modelled	Actual	Comment	Reference
D	-5	GLY	-	expression tag	UNP B8HNS6
D	-4	PRO	-	expression tag	UNP B8HNS6
D	-3	GLY	-	expression tag	UNP B8HNS6
D	-2	TYR	-	expression tag	UNP B8HNS6
D	-1	GLN	-	expression tag	UNP B8HNS6
D	0	ASP	-	expression tag	UNP B8HNS6
D	1	PRO	-	expression tag	UNP B8HNS6

- Molecule 2 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: C<sub>34</sub>H<sub>32</sub>FeN<sub>4</sub>O<sub>4</sub>).



Mol	Chain	Residues	Atoms						ZeroOcc	AltConf
2	A	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		
2	B	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		
2	C	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		
2	D	1	Total	C	Fe	H	N	O	0	0
			73	34	1	30	4	4		

- Molecule 3 is FLUORIDE ION (three-letter code: F) (formula: F).


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	B	1	Total F 1 1	0	0

*Continued on next page...*

*Continued from previous page...*


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total F 1 1	0	0
3	D	1	Total F 1 1	0	0
3	C	1	Total F 1 1	0	0

- Molecule 4 is GLYCEROL (three-letter code: GOL) (formula: C<sub>3</sub>H<sub>8</sub>O<sub>3</sub>).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	1	Total C H O 12 3 6 3	0	0
4	B	1	Total C H O 12 3 6 3	0	0
4	C	1	Total C H O 13 3 7 3	0	0
4	D	1	Total C H O 13 3 7 3	0	0

- Molecule 5 is SULFATE ION (three-letter code: SO4) (formula: O<sub>4</sub>S).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total    O    S 5    4    1	0	0
5	D	1	Total    O    S 5    4    1	0	0


- Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	A	243	Total    O 243    243	0	0
6	B	249	Total    O 249    249	0	0
6	C	246	Total    O 246    246	0	0
6	D	243	Total    O 243    243	0	0


### 3 Residue-property plots

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.


- Molecule 1: Chlorite dismutase




- Molecule 1: Chlorite dismutase



- Molecule 1: Chlorite dismutase



- Molecule 1: Chlorite dismutase



## 4 Data and refinement statistics i

Property	Value	Source
Space group	P 1	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	51.15 Å    54.70 Å    94.27 Å 99.06°    94.84°    99.04°	Depositor
Resolution (Å)	46.34 – 1.18 46.34 – 1.18	Depositor EDS
% Data completeness (in resolution range)	90.2 (46.34-1.18) 90.4 (46.34-1.18)	Depositor EDS
$R_{merge}$	0.06	Depositor
$R_{sym}$	(Not available)	Depositor
$\langle I/\sigma(I) \rangle^1$	1.43 (at 1.18 Å)	Xtriage
Refinement program	PHENIX dev_2712	Depositor
$R$ , $R_{free}$	0.190 , 0.222 0.190 , 0.222	Depositor DCC
$R_{free}$ test set	14770 reflections (5.01%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	15.1	Xtriage
Anisotropy	0.142	Xtriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.43 , 50.9	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.51$ , $\langle L^2 \rangle = 0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
$F_o, F_c$ correlation	0.98	EDS
Total number of atoms	13105	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	22.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The analyses of the Patterson function reveals a significant off-origin peak that is 97.71 % of the origin peak, indicating pseudo-translational symmetry. The chance of finding a peak of this or larger height randomly in a structure without pseudo-translational symmetry is equal to 1.0686e-10. The detected translational NCS is most likely also responsible for the elevated intensity ratio.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality i

### 5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: GOL, F, SO4, HEM

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	A	1.16	6/1521 (0.4%)	1.71	13/2052 (0.6%)
1	B	1.11	4/1559 (0.3%)	1.51	17/2103 (0.8%)
1	C	1.15	6/1527 (0.4%)	1.99	18/2060 (0.9%)
1	D	1.09	5/1564 (0.3%)	1.44	25/2109 (1.2%)
All	All	1.12	21/6171 (0.3%)	1.67	73/8324 (0.9%)

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	60	ARG	CZ-NH2	-17.63	1.10	1.33
1	A	60	ARG	CZ-NH2	11.98	1.48	1.33
1	A	60	ARG	CZ-NH1	8.53	1.44	1.33
1	B	60	ARG	CZ-NH1	8.04	1.43	1.33
1	D	60	ARG	NE-CZ	7.55	1.42	1.33
1	C	55	PHE	CE1-CZ	7.44	1.51	1.37
1	D	60	ARG	CZ-NH2	7.39	1.42	1.33
1	C	60	ARG	NE-CZ	7.29	1.42	1.33
1	C	172	GLU	CG-CD	7.14	1.62	1.51
1	C	60	ARG	CZ-NH1	6.93	1.42	1.33
1	C	172	GLU	CB-CG	-6.81	1.39	1.52
1	A	172	GLU	CG-CD	6.79	1.62	1.51
1	A	172	GLU	CB-CG	-6.60	1.39	1.52
1	D	172	GLU	CG-CD	6.25	1.61	1.51
1	B	81	GLU	CG-CD	6.21	1.61	1.51
1	B	172	GLU	CG-CD	6.03	1.60	1.51
1	D	138	GLU	CD-OE1	5.98	1.32	1.25
1	C	150	GLU	CD-OE2	5.46	1.31	1.25
1	A	102	ASP	CB-CG	5.27	1.62	1.51
1	A	141	PHE	CB-CG	-5.25	1.42	1.51
1	D	60	ARG	C-N	-5.02	1.22	1.34

All (73) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	60	ARG	NE-CZ-NH2	59.12	149.86	120.30
1	A	60	ARG	NE-CZ-NH2	50.78	145.69	120.30
1	C	60	ARG	NE-CZ-NH1	-40.96	99.82	120.30
1	B	60	ARG	NE-CZ-NH1	-37.34	101.63	120.30
1	D	60	ARG	NE-CZ-NH2	27.09	133.85	120.30
1	A	60	ARG	NH1-CZ-NH2	-21.60	95.64	119.40
1	B	60	ARG	NH1-CZ-NH2	18.31	139.54	119.40
1	D	134	ASP	CB-CG-OD2	15.26	132.03	118.30
1	D	60	ARG	NE-CZ-NH1	-14.98	112.81	120.30
1	A	60	ARG	CD-NE-CZ	14.13	143.38	123.60
1	C	60	ARG	CD-NE-CZ	12.90	141.66	123.60
1	B	134	ASP	CB-CG-OD2	-10.30	109.03	118.30
1	A	134	ASP	CB-CG-OD1	10.20	127.48	118.30
1	A	60	ARG	CG-CD-NE	-10.09	90.62	111.80
1	B	60	ARG	CD-NE-CZ	9.89	137.45	123.60
1	C	105	ARG	NE-CZ-NH2	-9.71	115.44	120.30
1	C	173	ARG	NE-CZ-NH2	-9.69	115.45	120.30
1	D	55	PHE	CB-CG-CD2	-9.41	114.21	120.80
1	B	163[A]	ARG	NE-CZ-NH2	-8.80	115.90	120.30
1	B	163[B]	ARG	NE-CZ-NH2	-8.80	115.90	120.30
1	C	60	ARG	NH1-CZ-NH2	-8.53	110.02	119.40
1	D	162[A]	MET	CG-SD-CE	8.43	113.68	100.20
1	D	162[B]	MET	CG-SD-CE	8.43	113.68	100.20
1	D	79[A]	ARG	NE-CZ-NH2	-8.41	116.10	120.30
1	D	79[B]	ARG	NE-CZ-NH2	-8.41	116.10	120.30
1	A	162[A]	MET	CG-SD-CE	8.01	113.02	100.20
1	A	162[B]	MET	CG-SD-CE	8.01	113.02	100.20
1	D	163[A]	ARG	NE-CZ-NH2	-7.72	116.44	120.30
1	D	163[B]	ARG	NE-CZ-NH2	-7.72	116.44	120.30
1	B	55	PHE	CB-CG-CD2	7.50	126.05	120.80
1	B	162[A]	MET	CG-SD-CE	7.47	112.15	100.20
1	B	162[B]	MET	CG-SD-CE	7.47	112.15	100.20
1	D	134	ASP	OD1-CG-OD2	-7.37	109.29	123.30
1	C	162[A]	MET	CG-SD-CE	6.91	111.25	100.20
1	C	162[B]	MET	CG-SD-CE	6.91	111.25	100.20
1	D	102	ASP	CB-CG-OD2	-6.85	112.14	118.30
1	D	59	ILE	CA-CB-CG1	-6.75	98.18	111.00
1	C	55	PHE	CD1-CE1-CZ	-6.70	112.06	120.10
1	B	58	ASN	O-C-N	-6.59	112.15	122.70
1	D	60	ARG	CD-NE-CZ	6.56	132.79	123.60
1	C	21	ARG	NE-CZ-NH1	-6.54	117.03	120.30
1	C	79	ARG	NE-CZ-NH2	-6.53	117.03	120.30

Continued on next page...

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	D	55	PHE	CB-CG-CD1	6.41	125.28	120.80
1	A	60	ARG	CB-CG-CD	-6.24	95.37	111.60
1	A	102	ASP	CB-CG-OD2	-6.22	112.70	118.30
1	A	133	ARG	NE-CZ-NH2	-6.20	117.20	120.30
1	B	61	TYR	CB-CG-CD2	6.14	124.68	121.00
1	B	60	ARG	NE-CZ-NH2	-6.08	117.26	120.30
1	C	32	GLU	OE1-CD-OE2	5.90	130.38	123.30
1	A	58	ASN	CB-CG-OD1	-5.88	109.83	121.60
1	D	32	GLU	OE1-CD-OE2	5.81	130.27	123.30
1	D	133	ARG	NE-CZ-NH2	-5.80	117.40	120.30
1	D	61	TYR	N-CA-CB	5.66	120.79	110.60
1	D	99[A]	MET	CG-SD-CE	-5.60	91.24	100.20
1	D	99[B]	MET	CG-SD-CE	-5.60	91.24	100.20
1	D	60	ARG	NH1-CZ-NH2	-5.59	113.25	119.40
1	C	60	ARG	CG-CD-NE	5.56	123.47	111.80
1	B	99	MET	CG-SD-CE	-5.46	91.47	100.20
1	A	163[A]	ARG	NE-CZ-NH2	-5.45	117.58	120.30
1	A	163[B]	ARG	NE-CZ-NH2	-5.45	117.58	120.30
1	D	60	ARG	O-C-N	5.38	131.31	122.70
1	D	133	ARG	NE-CZ-NH1	5.36	122.98	120.30
1	B	57	SER	CA-C-N	-5.34	105.45	117.20
1	C	104	ARG	NE-CZ-NH1	5.28	122.94	120.30
1	B	55	PHE	CG-CD1-CE1	5.18	126.50	120.80
1	D	59	ILE	CA-CB-CG2	5.15	121.21	110.90
1	C	60	ARG	CB-CG-CD	5.13	124.95	111.60
1	C	105	ARG	NE-CZ-NH1	5.10	122.85	120.30
1	C	60	ARG	CB-CA-C	-5.09	100.22	110.40
1	B	58	ASN	N-CA-CB	5.07	119.72	110.60
1	C	79	ARG	NE-CZ-NH1	5.03	122.82	120.30
1	D	59	ILE	CG1-CB-CG2	5.03	122.47	111.40
1	B	11	ARG	NE-CZ-NH1	5.01	122.80	120.30

There are no chirality outliers.

There are no planarity outliers.

## 5.2 Too-close contacts

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1465	1457	1427	8	0
1	B	1487	1474	1424	5	0
1	C	1470	1458	1426	3	0
1	D	1488	1469	1413	7	0
2	A	43	30	30	0	0
2	B	43	30	30	0	0
2	C	43	30	30	0	0
2	D	43	30	30	0	0
3	A	1	0	0	0	0
3	B	1	0	0	0	0
3	C	1	0	0	0	0
3	D	1	0	0	0	0
4	A	6	6	6	0	0
4	B	6	6	7	0	0
4	C	6	7	7	0	0
4	D	6	7	7	0	0
5	A	5	0	0	0	0
5	D	5	0	0	0	0
6	A	243	0	0	5	1
6	B	249	0	0	1	0
6	C	246	0	0	2	1
6	D	243	0	0	5	2
All	All	7101	6004	5837	23	2

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All (23) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:D:78[A]:ASN:OD1	6:D:301:HOH:O	2.14	0.65
1:D:98[B]:GLU:OE1	6:D:302:HOH:O	2.15	0.64
1:C:60:ARG:NH2	6:C:301:HOH:O	2.31	0.63
1:B:150:GLU:HG2	6:B:414:HOH:O	1.99	0.61
1:A:95:GLN:NE2	6:A:303:HOH:O	2.34	0.60
1:A:48:SER:N	6:A:304:HOH:O	2.38	0.56
1:A:59:ILE:C	1:A:60:ARG:HD3	2.27	0.55
1:A:98:GLU:HG3	6:A:393:HOH:O	2.08	0.54
1:B:114:HIS:CE1	1:B:162[B]:MET:HE3	2.47	0.50
1:C:98:GLU:HG3	6:C:396:HOH:O	2.12	0.50
1:A:60:ARG:N	1:A:60:ARG:HD3	2.27	0.48
1:B:8[A]:ILE:CD1	1:B:51:ARG:HG3	2.44	0.47

*Continued on next page...*

*Continued from previous page...*

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:94:ALA:O	1:B:98:GLU:HG3	2.16	0.45
1:A:138:GLU:OE1	6:A:301:HOH:O	2.21	0.45
1:D:114:HIS:CE1	1:D:162[B]:MET:HE3	2.51	0.45
1:D:98[A]:GLU:HG3	6:D:406:HOH:O	2.16	0.44
1:D:98[A]:GLU:OE1	6:D:302:HOH:O	2.21	0.44
1:D:21:ARG:NH1	6:D:314:HOH:O	2.49	0.43
1:C:8:ILE:CD1	1:C:51:ARG:HG3	2.49	0.43
1:D:59:ILE:HA	1:D:59:ILE:HD13	1.80	0.42
1:A:20:ILE:HD12	1:A:30:LEU:HA	2.00	0.42
1:B:8[A]:ILE:O	1:B:34:VAL:HA	2.20	0.42
1:A:151:HIS:HE1	6:A:412:HOH:O	2.04	0.41

All (2) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
6:A:460:HOH:O	6:D:486:HOH:O[1_454]	1.97	0.23
6:C:480:HOH:O	6:D:433:HOH:O[1_565]	2.19	0.01

## 5.3 Torsion angles (i)

### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	176/188 (94%)	172 (98%)	4 (2%)	0	100 100
1	B	181/188 (96%)	178 (98%)	3 (2%)	0	100 100
1	C	177/188 (94%)	174 (98%)	3 (2%)	0	100 100
1	D	181/188 (96%)	178 (98%)	3 (2%)	0	100 100
All	All	715/752 (95%)	702 (98%)	13 (2%)	0	100 100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [\(i\)](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	153/159 (96%)	152 (99%)	1 (1%)	84 57
1	B	157/159 (99%)	156 (99%)	1 (1%)	86 61
1	C	154/159 (97%)	153 (99%)	1 (1%)	86 61
1	D	158/159 (99%)	157 (99%)	1 (1%)	86 61
All	All	622/636 (98%)	618 (99%)	4 (1%)	84 61

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	168	TRP
1	B	168	TRP
1	C	168	TRP
1	D	168	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [\(i\)](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [\(i\)](#)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates [\(i\)](#)

There are no carbohydrates in this entry.

## 5.6 Ligand geometry (i)

Of 14 ligands modelled in this entry, 4 are monoatomic - leaving 10 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z  > 2$	Counts	RMSZ	# $ Z  > 2$
4	GOL	B	203	-	5,5,5	1.32	1 (20%)	5,5,5	0.77	0
4	GOL	A	203	-	5,5,5	1.19	0	5,5,5	0.52	0
4	GOL	D	203	-	5,5,5	1.40	1 (20%)	5,5,5	0.73	0
4	GOL	C	203	-	5,5,5	1.02	0	5,5,5	0.81	0
2	HEM	B	201	1	27,50,50	1.86	5 (18%)	17,82,82	2.51	8 (47%)
5	SO4	D	204	-	4,4,4	0.40	0	6,6,6	0.35	0
2	HEM	D	201	1	27,50,50	1.65	4 (14%)	17,82,82	2.09	9 (52%)
5	SO4	A	204	-	4,4,4	0.26	0	6,6,6	0.57	0
2	HEM	C	201	1	27,50,50	1.47	4 (14%)	17,82,82	2.71	9 (52%)
2	HEM	A	201	1	27,50,50	2.20	6 (22%)	17,82,82	2.12	5 (29%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
4	GOL	B	203	-	-	0/4/4/4	-
4	GOL	A	203	-	-	0/4/4/4	-
4	GOL	D	203	-	-	2/4/4/4	-
4	GOL	C	203	-	-	0/4/4/4	-
2	HEM	B	201	1	-	0/6/54/54	-
2	HEM	D	201	1	-	0/6/54/54	-
2	HEM	C	201	1	-	0/6/54/54	-
2	HEM	A	201	1	-	0/6/54/54	-

All (21) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	A	201	HEM	C3B-C2B	-8.10	1.29	1.40

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	201	HEM	C3B-C2B	-5.10	1.33	1.40
2	D	201	HEM	C3C-C2C	-4.93	1.33	1.40
2	D	201	HEM	C3B-C2B	-4.60	1.34	1.40
2	B	201	HEM	C3C-C2C	-4.39	1.34	1.40
2	C	201	HEM	C3C-C2C	-4.30	1.34	1.40
2	B	201	HEM	CAA-C2A	4.23	1.58	1.52
2	A	201	HEM	C3C-C2C	-3.54	1.35	1.40
2	A	201	HEM	C3B-CAB	3.48	1.55	1.47
2	D	201	HEM	C3C-CAC	2.95	1.53	1.47
2	A	201	HEM	C1A-NA	2.74	1.41	1.36
2	C	201	HEM	C3B-C2B	-2.51	1.36	1.40
2	C	201	HEM	C4B-NB	2.49	1.41	1.36
2	B	201	HEM	C3B-CAB	2.46	1.52	1.47
4	D	203	GOL	O2-C2	-2.42	1.36	1.43
2	A	201	HEM	C3D-C2D	-2.27	1.31	1.37
4	B	203	GOL	O1-C1	-2.21	1.33	1.42
2	C	201	HEM	C3D-C2D	-2.20	1.31	1.37
2	D	201	HEM	C1A-NA	2.11	1.40	1.36
2	A	201	HEM	C1A-CHA	-2.06	1.35	1.41
2	B	201	HEM	C3C-CAC	2.00	1.51	1.47

All (31) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	C	201	HEM	CMD-C2D-C1D	-5.97	119.30	128.46
2	B	201	HEM	CMD-C2D-C1D	-4.89	120.95	128.46
2	A	201	HEM	CAA-CBA-CGA	-4.75	104.71	112.67
2	B	201	HEM	C1D-C2D-C3D	4.64	110.22	107.00
2	A	201	HEM	CMA-C3A-C4A	-4.34	121.79	128.46
2	C	201	HEM	C4C-C3C-C2C	4.06	109.73	106.90
2	C	201	HEM	C1D-C2D-C3D	3.65	109.54	107.00
2	C	201	HEM	CMB-C2B-C3B	3.65	131.50	124.68
2	C	201	HEM	CMC-C2C-C3C	3.47	131.17	124.68
2	A	201	HEM	C1D-C2D-C3D	-3.33	104.68	107.00
2	C	201	HEM	CMD-C2D-C3D	3.27	131.10	124.94
2	C	201	HEM	CMA-C3A-C4A	-3.26	123.46	128.46
2	B	201	HEM	C4A-C3A-C2A	3.07	109.13	107.00
2	D	201	HEM	C4C-C3C-C2C	3.05	109.03	106.90
2	B	201	HEM	CMA-C3A-C4A	-3.01	123.84	128.46
2	B	201	HEM	CAA-CBA-CGA	-2.97	107.69	112.67
2	D	201	HEM	CMA-C3A-C4A	-2.95	123.93	128.46
2	B	201	HEM	CMB-C2B-C3B	2.92	130.15	124.68

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	D	201	HEM	CAA-CBA-CGA	-2.80	107.97	112.67
2	B	201	HEM	CBA-CAA-C2A	-2.80	107.32	112.49
2	A	201	HEM	CMA-C3A-C2A	2.76	130.14	124.94
2	D	201	HEM	CBD-CAD-C3D	-2.74	107.43	112.48
2	D	201	HEM	C3C-C4C-NC	-2.53	106.16	110.94
2	D	201	HEM	CMA-C3A-C2A	2.50	129.65	124.94
2	D	201	HEM	CMD-C2D-C1D	-2.46	124.68	128.46
2	B	201	HEM	CMC-C2C-C3C	2.24	128.88	124.68
2	C	201	HEM	CMA-C3A-C2A	2.15	129.00	124.94
2	D	201	HEM	CAD-CBD-CGD	2.10	116.19	112.67
2	C	201	HEM	C3C-C4C-NC	-2.08	107.02	110.94
2	A	201	HEM	CMD-C2D-C3D	2.05	128.80	124.94
2	D	201	HEM	CMD-C2D-C3D	2.02	128.76	124.94

There are no chirality outliers.


All (2) torsion outliers are listed below:


Mol	Chain	Res	Type	Atoms
4	D	203	GOL	O1-C1-C2-O2
4	D	203	GOL	O1-C1-C2-C3


There are no ring outliers.


No monomer is involved in short contacts.

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less than 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.









## 5.7 Other polymers [\(i\)](#)

There are no such residues in this entry.

## 5.8 Polymer linkage issues [\(i\)](#)

There are no chain breaks in this entry.

## 6 Fit of model and data i

### 6.1 Protein, DNA and RNA chains i

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	174/188 (92%)	-0.18	4 (2%) 60 61	11, 18, 30, 42	0
1	B	175/188 (93%)	-0.15	4 (2%) 60 61	11, 19, 30, 41	0
1	C	175/188 (93%)	-0.15	3 (1%) 70 70	11, 18, 30, 44	0
1	D	175/188 (93%)	-0.11	6 (3%) 45 47	10, 19, 30, 42	0
All	All	699/752 (92%)	-0.15	17 (2%) 59 59	10, 18, 31, 44	0

All (17) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	40	ALA	6.5
1	B	40	ALA	5.2
1	C	39	GLY	4.8
1	C	47	ASP	4.6
1	A	39	GLY	4.1
1	C	40	ALA	4.0
1	A	40	ALA	3.2
1	D	2	ASN	3.1
1	D	47	ASP	2.8
1	B	2	ASN	2.7
1	B	39	GLY	2.6
1	D	39	GLY	2.5
1	A	2	ASN	2.2
1	B	47	ASP	2.2
1	D	38	ASN	2.1
1	A	110	ARG	2.1
1	D	51	ARG	2.1

### 6.2 Non-standard residues in protein, DNA, RNA chains i

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [\(i\)](#)


There are no carbohydrates in this entry.


## 6.4 Ligands [\(i\)](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.


Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
4	GOL	D	203	6/6	0.90	0.12	31,38,45,47	0
3	F	A	202	1/1	0.90	0.10	35,35,35,35	0
4	GOL	C	203	6/6	0.91	0.14	31,38,45,46	0
4	GOL	A	203	6/6	0.91	0.10	30,38,44,44	0
3	F	D	202	1/1	0.92	0.12	33,33,33,33	0
3	F	C	202	1/1	0.92	0.09	34,34,34,34	0
3	F	B	202	1/1	0.92	0.11	29,29,29,29	0
4	GOL	B	203	6/6	0.93	0.09	32,37,43,43	0
5	SO4	D	204	5/5	0.94	0.17	57,58,58,58	0
5	SO4	A	204	5/5	0.96	0.15	54,55,55,55	0
2	HEM	D	201	43/43	0.99	0.07	11,13,18,20	0
2	HEM	B	201	43/43	0.99	0.06	11,13,17,19	0
2	HEM	C	201	43/43	0.99	0.07	11,13,17,20	0
2	HEM	A	201	43/43	0.99	0.06	11,13,18,20	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.









## 6.5 Other polymers (i)

There are no such residues in this entry.