



# Full wwPDB X-ray Structure Validation Report ⓘ

Nov 15, 2023 – 11:33 PM JST

PDB ID : 6K9R  
Title : Crystal Structure Analysis of Endo-beta-1,4-xylanase II Complexed with Xylotriose  
Authors : Li, C.; Wan, Q.  
Deposited on : 2019-06-17  
Resolution : 1.30 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at [validation@mail.wwpdb.org](mailto:validation@mail.wwpdb.org)

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

---

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467  
Mogul : 1.8.5 (274361), CSD as541be (2020)  
Xtriage (Phenix) : 1.13  
EDS : 2.36  
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)  
Refmac : 5.8.0158  
CCP4 : 7.0.044 (Gargrove)  
Ideal geometry (proteins) : Engh & Huber (2001)  
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)  
Validation Pipeline (wwPDB-VP) : 2.36


# 1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

*X-RAY DIFFRACTION*

The reported resolution of this entry is 1.30 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
$R_{free}$	130704	1058 (1.30-1.30)
Clashscore	141614	1101 (1.30-1.30)
Ramachandran outliers	138981	1058 (1.30-1.30)
Sidechain outliers	138945	1058 (1.30-1.30)
RSRZ outliers	127900	1029 (1.30-1.30)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for  $\geq 3$ , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions  $\leq 5\%$ . The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	189	
2	B	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
2	XYP	B	1	X	-	-	-

## 2 Entry composition i

There are 4 unique types of molecules in this entry. The entry contains 3231 atoms, of which 1438 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.


- Molecule 1 is a protein called Endo-1,4-beta-xylanase 2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	H	N	O				S
1	A	189	2944	963	1410	261	309	1	0	18	0

There are 2 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	44	GLU	ASN	engineered mutation	UNP P36217
A	177	GLN	GLU	engineered mutation	UNP P36217

- Molecule 2 is an oligosaccharide called beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	H	O			
2	B	3	56	15	28	13	0	0	0

- Molecule 3 is IODIDE ION (three-letter code: IOD) (formula: I).

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	4	Total	I	0	0
			4	4		


- Molecule 4 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
4	A	227	Total	O	0	0
			227	227		


### 3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ( $RSRZ > 2$ ). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: Endo-1,4-beta-xylanase 2



- Molecule 2: beta-D-xylopyranose-(1-4)-beta-D-xylopyranose-(1-4)-beta-D-xylopyranose



## 4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, $\alpha$ , $\beta$ , $\gamma$	48.92Å 58.59Å 69.34Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	39.97 – 1.30 39.97 – 1.13	Depositor EDS
% Data completeness (in resolution range)	85.4 (39.97-1.30) 82.8 (39.97-1.13)	Depositor EDS
$R_{merge}$	(Not available)	Depositor
$R_{sym}$	0.03	Depositor
$\langle I/\sigma(I) \rangle$ <sup>1</sup>	1.77 (at 1.13Å)	Xtrriage
Refinement program	PHENIX 1.11.1_2575	Depositor
R, $R_{free}$	0.140 , 0.158 0.141 , 0.158	Depositor DCC
$R_{free}$ test set	3107 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å <sup>2</sup> )	7.2	Xtrriage
Anisotropy	0.034	Xtrriage
Bulk solvent $k_{sol}$ (e/Å <sup>3</sup> ), $B_{sol}$ (Å <sup>2</sup> )	0.41 , 54.5	EDS
L-test for twinning <sup>2</sup>	$\langle  L  \rangle = 0.48$ , $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
$F_o, F_c$ correlation	0.96	EDS
Total number of atoms	3231	wwPDB-VP
Average B, all atoms (Å <sup>2</sup> )	8.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 8.09% of the height of the origin peak. No significant pseudotranslation is detected.*

<sup>1</sup>Intensities estimated from amplitudes.

<sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

## 5 Model quality [i](#)

### 5.1 Standard geometry [i](#)

Bond lengths and bond angles in the following residue types are not validated in this section: IOD, XYP

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 5$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z  >5	RMSZ	# Z  >5
1	A	0.42	0/1646	0.64	0/2245

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	75[A]	SER	Peptide

### 5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1534	1410	1338	10	0
2	B	28	28	0	1	0

*Continued on next page...*

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	4	0	0	0	0
4	A	227	0	0	7	3
All	All	1793	1438	1338	11	3

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (11) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:44[A]:GLU:HG3	4:A:368:HOH:O	1.68	0.90
1:A:151:ASN:OD1	4:A:301:HOH:O	2.01	0.78
1:A:44[B]:GLU:HG3	4:A:360:HOH:O	1.93	0.69
1:A:44[B]:GLU:HG2	1:A:179:TYR:CD1	2.45	0.52
1:A:2:THR:CG2	4:A:427:HOH:O	2.62	0.47
1:A:2:THR:N	4:A:306:HOH:O	2.48	0.47
1:A:86:GLU:O	1:A:136:GLN:HA	2.18	0.43
1:A:42:SER:O	1:A:180:PHE:HA	2.19	0.42
1:A:44[A]:GLU:CG	4:A:368:HOH:O	2.44	0.42
1:A:2:THR:HA	1:A:28[B]:THR:HG22	2.00	0.41
4:A:302:HOH:O	2:B:3:XYP:O4	2.22	0.41

All (3) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
4:A:416:HOH:O	4:A:511:HOH:O[3_554]	1.95	0.25
4:A:490:HOH:O	4:A:515:HOH:O[3_644]	1.98	0.22
4:A:407:HOH:O	4:A:503:HOH:O[2_564]	2.07	0.13

## 5.3 Torsion angles [i](#)

### 5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.



Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	205/189 (108%)	202 (98%)	3 (2%)	0	100	100

There are no Ramachandran outliers to report.

### 5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	172/155 (111%)	172 (100%)	0	100	100

There are no protein residues with a non-rotameric sidechain to report.

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

### 5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

## 5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 5.5 Carbohydrates [i](#)

3 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with  $|Z| > 2$  is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z  > 2	Counts	RMSZ	# Z  > 2
2	XYP	B	1	2	10,10,10	4.06	6 (60%)	14,14,14	3.63	7 (50%)
2	XYP	B	2	2	9,9,10	1.21	1 (11%)	10,12,14	1.33	1 (10%)
2	XYP	B	3	2	9,9,10	2.96	3 (33%)	10,12,14	2.20	4 (40%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	XYP	B	1	2	1/1/4/4	-	0/1/1/1
2	XYP	B	2	2	-	-	0/1/1/1
2	XYP	B	3	2	-	-	0/1/1/1

All (10) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	B	1	XYP	O1-C1	-7.43	1.15	1.39
2	B	1	XYP	C4-C3	6.77	1.62	1.52
2	B	3	XYP	C4-C3	-6.29	1.43	1.52
2	B	1	XYP	O5-C5	-5.58	1.34	1.43
2	B	3	XYP	O4-C4	4.46	1.52	1.43
2	B	1	XYP	C1-C2	-3.77	1.43	1.52
2	B	2	XYP	C2-C3	-2.88	1.48	1.52
2	B	3	XYP	C2-C3	-2.49	1.48	1.52
2	B	1	XYP	O5-C1	2.36	1.46	1.43
2	B	1	XYP	C3-C2	2.26	1.58	1.52

All (12) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	1	XYP	O1-C1-C2	10.34	138.13	109.03
2	B	1	XYP	O5-C1-C2	-5.51	101.25	109.43
2	B	3	XYP	C5-C4-C3	4.87	115.65	109.67
2	B	1	XYP	O1-C1-O5	3.58	119.04	109.72
2	B	2	XYP	C5-O5-C1	-3.37	106.34	111.52
2	B	3	XYP	C4-C3-C2	-2.93	107.45	110.92
2	B	1	XYP	O3-C3-C4	2.63	115.03	109.99
2	B	1	XYP	O5-C5-C4	2.42	114.51	110.77
2	B	1	XYP	O2-C2-C3	2.36	115.80	110.35
2	B	3	XYP	C5-O5-C1	2.36	115.15	111.52

*Continued on next page...*

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	B	3	XYP	O3-C3-C2	-2.17	105.83	109.99
2	B	1	XYP	O4-C4-C3	2.01	114.17	110.14

All (1) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
2	B	1	XYP	C1


There are no torsion outliers.

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	B	3	XYP	1	0

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



## 5.6 Ligand geometry [i](#)

Of 4 ligands modelled in this entry, 4 are monoatomic - leaving 0 for Mogul analysis.

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no torsion outliers.

There are no ring outliers.

No monomer is involved in short contacts.

## 5.7 Other polymers

There are no such residues in this entry.

## 5.8 Polymer linkage issues

There are no chain breaks in this entry.

## 6 Fit of model and data

### 6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95<sup>th</sup> percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å <sup>2</sup> )	Q<0.9
1	A	189/189 (100%)	2.76	143 (75%) <b>0</b>   <b>0</b>	2, 6, 13, 26	3 (1%)

All (143) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	2	THR	11.0
1	A	3	ILE	6.6
1	A	18	TRP	5.8
1	A	25[A]	VAL	5.6
1	A	129	ILE	4.6
1	A	135	TYR	4.6
1	A	159	TRP	4.5
1	A	85	ILE	4.5
1	A	137	TYR	4.3
1	A	123	VAL	4.2
1	A	138	TRP	4.2
1	A	22	HIS	4.2
1	A	39	TRP	4.1
1	A	51	TRP	4.1
1	A	9	TYR	4.1
1	A	115	TYR	4.1
1	A	37[A]	VAL	4.0
1	A	156	PHE	3.9
1	A	87	TYR	3.9
1	A	88	TYR	3.9
1	A	27	TYR	3.9
1	A	164	LEU	3.9
1	A	179	TYR	3.9
1	A	28[A]	THR	3.8
1	A	140	VAL	3.8
1	A	89	ILE	3.8
1	A	57	ASN	3.8

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	173	ILE	3.7
1	A	13	TYR	3.7
1	A	118	TYR	3.7
1	A	31	PRO	3.6
1	A	77	TYR	3.6
1	A	79	TRP	3.6
1	A	35	PHE	3.6
1	A	15	TYR	3.6
1	A	90	VAL	3.6
1	A	96	TYR	3.6
1	A	165	THR	3.6
1	A	93	PHE	3.6
1	A	76[A]	VAL	3.5
1	A	174	VAL	3.5
1	A	21	GLY	3.5
1	A	128	ILE	3.5
1	A	187	ILE	3.5
1	A	171	TYR	3.5
1	A	119	ARG	3.4
1	A	108	VAL	3.4
1	A	17	TYR	3.4
1	A	66[A]	TYR	3.4
1	A	117	ILE	3.4
1	A	134	PHE	3.4
1	A	75[A]	SER	3.3
1	A	38	ASN	3.3
1	A	126	PRO	3.3
1	A	158	ALA	3.3
1	A	60	ILE	3.3
1	A	176	VAL	3.2
1	A	74	LEU	3.2
1	A	84	LEU	3.2
1	A	62	PHE	3.2
1	A	73	TYR	3.1
1	A	168[B]	THR	3.1
1	A	45	PHE	3.1
1	A	160	ALA	3.1
1	A	101	GLY	3.1
1	A	166	LEU	3.1
1	A	55	THR	3.1
1	A	100	THR	3.1
1	A	12	GLY	3.1

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	23	GLY	3.1
1	A	46	VAL	3.0
1	A	5	PRO	3.0
1	A	122	ARG	2.9
1	A	169[B]	MET	2.9
1	A	153	ALA	2.9
1	A	190	SER	2.9
1	A	50	GLY	2.9
1	A	185	ALA	2.9
1	A	61	ASN	2.9
1	A	109	THR	2.9
1	A	175	ALA	2.8
1	A	59	VAL	2.8
1	A	103	THR	2.8
1	A	143	ASN	2.8
1	A	102	ALA	2.8
1	A	11	ASN	2.8
1	A	149[A]	SER	2.8
1	A	120	THR	2.7
1	A	163	GLY	2.7
1	A	105	LEU	2.7
1	A	130	GLY	2.7
1	A	180	PHE	2.7
1	A	33	GLY	2.6
1	A	14	PHE	2.6
1	A	150[A]	VAL	2.6
1	A	95	THR	2.6
1	A	189	VAL	2.6
1	A	4	GLN	2.5
1	A	152	THR	2.5
1	A	67	ASN	2.5
1	A	106	GLY	2.5
1	A	48	GLY	2.5
1	A	70	GLY	2.5
1	A	114	VAL	2.5
1	A	188	THR	2.4
1	A	36[A]	SER	2.4
1	A	127	SER	2.4
1	A	40	SER	2.4
1	A	83	PRO	2.4
1	A	98	PRO	2.4
1	A	112	GLY	2.4

*Continued on next page...*

*Continued from previous page...*

Mol	Chain	Res	Type	RSRZ
1	A	178	GLY	2.4
1	A	72	SER	2.4
1	A	69	ASN	2.3
1	A	53	PRO	2.3
1	A	65[A]	SER	2.3
1	A	167	GLY	2.3
1	A	80	SER	2.3
1	A	26[A]	THR	2.3
1	A	78	GLY	2.3
1	A	131	THR	2.3
1	A	32	GLY	2.3
1	A	7	THR	2.3
1	A	124	ASN	2.2
1	A	157	ASN	2.2
1	A	111	ASP	2.2
1	A	44[A]	GLU	2.2
1	A	132	ALA	2.2
1	A	139	SER	2.2
1	A	155	HIS	2.1
1	A	41	ASN	2.1
1	A	19	ASN	2.1
1	A	154	ASN	2.1
1	A	125	GLN	2.1
1	A	184	SER	2.1
1	A	30	GLY	2.1
1	A	94	GLY	2.1
1	A	145	ARG	2.0
1	A	68	PRO	2.0
1	A	42	SER	2.0
1	A	146[A]	SER	2.0
1	A	147	SER	2.0
1	A	161	GLN	2.0

## 6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

## 6.3 Carbohydrates [i](#)


In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
2	XYP	B	1	10/10	0.82	0.33	7,15,20,23	0
2	XYP	B	3	9/10	0.91	0.20	9,14,17,18	0
2	XYP	B	2	9/10	0.94	0.18	5,6,7,8	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.



## 6.4 Ligands [i](#)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95<sup>th</sup> percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	B-factors(Å <sup>2</sup> )	Q<0.9
3	IOD	A	205	1/1	0.98	0.22	26,26,26,26	0
3	IOD	A	206	1/1	0.99	0.20	35,35,35,35	0
3	IOD	A	207	1/1	0.99	0.18	20,20,20,20	0
3	IOD	A	204	1/1	1.00	0.17	5,5,5,5	0

## 6.5 Other polymers [i](#)

There are no such residues in this entry.