

# Full wwPDB X-ray Structure Validation Report (i)

#### Jul 29, 2024 – 10:09 AM JST

PDB ID	:	8KAQ
Title	:	Glycoside hydrolase family 1 beta-glucosidase (E318G mutant) from Strepto-
		myces griseus (sophorose complex)
Authors	:	Kumakura, H.; Motouchi, S.; Nakai, H.; Nakajima, M.
Deposited on	:	2023-08-03
Resolution	:	2.13 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix)	:	1.13
$\mathrm{EDS}$	:	2.37.1
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

# 1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 2.13 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} {\rm Whole \ archive} \\ (\#{\rm Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
$R_{free}$	130704	2523 (2.16-2.12)
Clashscore	141614	2653 (2.16-2.12)
Ramachandran outliers	138981	2618 (2.16-2.12)
Sidechain outliers	138945	2617 (2.16-2.12)
RSRZ outliers	127900	2485 (2.16-2.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain	
1	А	414	2% <b>88</b> %	8% ••
1	В	414	86%	8% • 5%
1	С	414	86%	10% ••
1	D	414	7% 12%	• 12%

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit crite-



ria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	CL	D	502	-	-	Х	-



#### 8KAQ

# 2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 12837 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
1	Δ	401	Total	С	Ν	0	$\mathbf{S}$	0	0	0
	A	401	3114	1968	548	590	8	0	0	0
1	р	205	Total	С	Ν	0	S	0	0	0
	D	395	3067	1939	538	582	8	0	0	0
1	C	1 400	Total	С	Ν	0	S	0	0	0
	400	3110	1967	545	590	8	0	0		
1	1 D	264	Total	С	Ν	0	S	0	0	0
	304	2854	1807	502	537	8	0	0	0	

• Molecule 1 is a protein called Putative beta-glucosidase.

There are 36 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	318	GLY	GLU	engineered mutation	UNP B1W1N0
А	407	LEU	-	expression tag	UNP B1W1N0
А	408	GLU	-	expression tag	UNP B1W1N0
А	409	HIS	-	expression tag	UNP B1W1N0
А	410	HIS	-	expression tag	UNP B1W1N0
А	411	HIS	-	expression tag	UNP B1W1N0
А	412	HIS	-	expression tag	UNP B1W1N0
А	413	HIS	-	expression tag	UNP B1W1N0
А	414	HIS	-	expression tag	UNP B1W1N0
В	318	GLY	GLU	engineered mutation	UNP B1W1N0
В	407	LEU	-	expression tag	UNP B1W1N0
В	408	GLU	-	expression tag	UNP B1W1N0
В	409	HIS	-	expression tag	UNP B1W1N0
В	410	HIS	-	expression tag	UNP B1W1N0
В	411	HIS	-	expression tag	UNP B1W1N0
В	412	HIS	-	expression tag	UNP B1W1N0
В	413	HIS	-	expression tag	UNP B1W1N0
В	414	HIS	-	expression tag	UNP B1W1N0
С	318	GLY	GLU	engineered mutation	UNP B1W1N0
С	407	LEU	-	expression tag	UNP B1W1N0
С	408	GLU	-	expression tag	UNP B1W1N0



Chain	Residue	Modelled	Actual	Comment	Reference
С	409	HIS	-	expression tag	UNP B1W1N0
С	410	HIS	-	expression tag	UNP B1W1N0
С	411	HIS	-	expression tag	UNP B1W1N0
С	412	HIS	-	expression tag	UNP B1W1N0
С	413	HIS	-	expression tag	UNP B1W1N0
С	414	HIS	-	expression tag	UNP B1W1N0
D	318	GLY	GLU	engineered mutation	UNP B1W1N0
D	407	LEU	-	expression tag	UNP B1W1N0
D	408	GLU	-	expression tag	UNP B1W1N0
D	409	HIS	-	expression tag	UNP B1W1N0
D	410	HIS	-	expression tag	UNP B1W1N0
D	411	HIS	-	expression tag	UNP B1W1N0
D	412	HIS	-	expression tag	UNP B1W1N0
D	413	HIS	-	expression tag	UNP B1W1N0
D	414	HIS	-	expression tag	UNP B1W1N0

• Molecule 2 is beta-D-glucopyranose (three-letter code: BGC) (formula:  $C_6H_{12}O_6$ ) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	А	1	Total         C         O           12         6         6	0	0
2	А	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 11  6  5 \end{array}$	0	0
2	А	1	Total         C         O           11         6         5	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
2	В	1	Total         C         O           11         6         5	0	0
2	С	1	Total         C         O           11         6         5	0	0
2	С	1	Total         C         O           12         6         6	0	0
2	С	1	Total         C         O           11         6         5	0	0
2	D	1	$\begin{array}{ccc} \text{Total}  \text{C}  \text{O} \\ 12  6  6 \end{array}$	0	0

• Molecule 3 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula:  $C_6H_{14}O_4$ ).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	А	1	Total C O 10 6 4	0	0
3	В	1	Total         C         O           10         6         4	0	0
3	С	1	Total         C         O           10         6         4	0	0

• Molecule 4 is alpha-D-glucopyranose (three-letter code: GLC) (formula:  $C_6H_{12}O_6$ ) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	А	1	Total         C         O           12         6         6	0	0
4	В	1	Total         C         O           12         6         6	0	0
4	С	1	Total         C         O           12         6         6	0	0

• Molecule 5 is CHLORIDE ION (three-letter code: CL) (formula: Cl).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	А	2	Total Cl 2 2	0	0
5	В	2	Total Cl 2 2	0	0
5	С	1	Total Cl 1 1	0	0
5	D	1	Total Cl 1 1	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	169	Total O 169 169	0	0
6	В	137	Total O 137 137	0	0



Continued from previous page...

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	С	143	Total O 143 143	0	0
6	D	80	Total         O           80         80	0	0



# 3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Putative beta-glucosidase





# 4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	91.79Å 100.18Å $96.05$ Å	Deperitor
a, b, c, $\alpha$ , $\beta$ , $\gamma$	$90.00^{\circ}$ $102.87^{\circ}$ $90.00^{\circ}$	Depositor
$\mathbf{P}_{\text{assolution}}(\hat{\mathbf{A}})$	46.82 - 2.13	Depositor
Resolution (A)	46.82 - 2.13	EDS
% Data completeness	98.5 (46.82-2.13)	Depositor
(in resolution range)	98.5(46.82-2.13)	EDS
R <sub>merge</sub>	(Not available)	Depositor
$R_{sym}$	(Not available)	Depositor
$< I/\sigma(I) > 1$	$2.20 (at 2.14 \text{\AA})$	Xtriage
Refinement program	REFMAC 5.8.0415	Depositor
D D.	0.183 , $0.223$	Depositor
$\Pi, \Pi_{free}$	0.192 , $0.229$	DCC
$R_{free}$ test set	4704 reflections $(5.04%)$	wwPDB-VP
Wilson B-factor $(Å^2)$	25.6	Xtriage
Anisotropy	0.027	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.38 , $37.8$	EDS
L-test for twinning <sup>2</sup>	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	0.013 for l,-k,h	Xtriage
$F_o, F_c$ correlation	0.95	EDS
Total number of atoms	12837	wwPDB-VP
Average B, all atoms $(Å^2)$	29.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 4.42% of the height of the origin peak. No significant pseudotranslation is detected.

<sup>&</sup>lt;sup>2</sup>Theoretical values of  $\langle |L| \rangle$ ,  $\langle L^2 \rangle$  for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



<sup>&</sup>lt;sup>1</sup>Intensities estimated from amplitudes.

# 5 Model quality (i)

## 5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: GLC, CL, BGC, PGE

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles	
	Chain	RMSZ	# Z  > 5	RMSZ	# Z  > 5
1	А	0.39	0/3205	0.71	0/4386
1	В	0.40	0/3156	0.71	0/4319
1	С	0.42	0/3202	0.72	0/4383
1	D	0.39	0/2933	0.70	0/4005
All	All	0.40	0/12496	0.71	0/17093

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	А	0	2
1	В	0	1
1	С	0	1
1	D	0	2
All	All	0	6

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (6) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	А	245	ARG	Sidechain
1	А	269	ARG	Sidechain
1	В	245	ARG	Sidechain
1	С	113	ARG	Sidechain
1	D	269	ARG	Sidechain



Continued from previous page...

Mol	Chain	Res	Type	Group
1	D	383	ARG	Sidechain

### 5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	3114	0	2945	19	0
1	В	3067	0	2904	20	0
1	С	3110	0	2939	27	0
1	D	2854	0	2703	28	0
2	А	34	0	32	4	0
2	В	11	0	10	5	0
2	С	34	0	31	0	0
2	D	12	0	12	0	0
3	А	10	0	14	2	0
3	В	10	0	14	0	0
3	С	10	0	14	0	0
4	А	12	0	11	0	0
4	В	12	0	12	5	0
4	С	12	0	11	0	0
5	А	2	0	0	0	0
5	В	2	0	0	2	0
5	С	1	0	0	1	0
5	D	1	0	0	2	0
6	А	169	0	0	1	0
6	В	137	0	0	1	0
6	C	143	0	0	1	0
6	D	80	0	0	2	0
All	All	12837	0	11652	102	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 4.

All (102) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.



8KAQ	
OILIG	

Atom_1	Atom_2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
4:B:501:GLC:O2	2:B:502:BGC:C1	1.68	1.39
2:A:501:BGC:O2	2:A:502:BGC:C1	1.82	1.25
1:B:105:ARG:NH2	5:D:502:CL:CL	2.48	0.84
1:A:161:SER:OG	3:A:503:PGE:H2	1.78	0.83
1:A:289:THR:O	1:A:290:THR:HB	1.76	0.83
1:C:253:ILE:HG23	1:C:309:VAL:HG13	1.59	0.83
4:B:501:GLC:C2	2:B:502:BGC:C1	2.57	0.82
1:B:289:THR:O	1:B:290:THR:HB	1.80	0.78
1:C:289:THR:O	1:C:290:THR:HB	1.84	0.75
1:C:253:ILE:CG2	1:C:309:VAL:HG13	2.16	0.74
2:A:501:BGC:HB	2:A:502:BGC:C1	2.01	0.72
1:C:174:VAL:HG12	1:C:175:MET:CE	2.23	0.68
2:A:501:BGC:C2	2:A:502:BGC:C1	2.72	0.67
1:D:20:LEU:HB2	1:D:352:GLY:HA3	1.79	0.64
1:A:20:LEU:HB2	1:A:352:GLY:HA3	1.80	0.63
1:B:174:VAL:HG12	1:B:175:MET:HE2	1.80	0.62
1:A:105:ARG:NH2	6:A:602:HOH:O	2.33	0.61
1:C:363:GLU:OE1	5:C:506:CL:CL	2.56	0.60
1:C:174:VAL:HG12	1:C:175:MET:HE2	1.82	0.60
1:D:250:ASP:OD1	1:D:304:HIS:NE2	2.36	0.59
1:C:234:LEU:HD21	1:C:271:ARG:HD2	1.87	0.57
1:B:174:VAL:HG12	1:B:175:MET:CE	2.34	0.57
1:D:230:VAL:HG12	1:D:230:VAL:O	2.04	0.56
1:B:156:ILE:O	5:B:505:CL:CL	2.60	0.56
1:A:161:SER:HA	3:A:503:PGE:H32	1.88	0.56
1:A:289:THR:O	1:A:290:THR:CB	2.52	0.55
1:B:178:GLN:NE2	1:B:185:SER:O	2.38	0.55
4:B:501:GLC:O2	2:B:502:BGC:O5	2.24	0.55
1:D:53:PRO:HG2	1:D:55:LEU:HD22	1.87	0.54
1:D:59:ASP:OD2	5:D:502:CL:CL	2.63	0.54
1:D:245:ARG:O	1:D:245:ARG:HG2	2.08	0.54
4:B:501:GLC:C2	2:B:502:BGC:O5	2.56	0.53
1:D:267:TYR:O	1:D:332:TYR:OH	2.25	0.52
1:B:20:LEU:HB2	1:B:352:GLY:HA3	1.90	0.52
5:B:504:CL:CL	6:B:703:HOH:O	2.56	0.52
1:D:105:ARG:NH2	6:D:603:HOH:O	2.42	0.52
1:D:15:PHE:CD2	1:D:399:ARG:HD3	2.45	0.52
1:B:15:PHE:HE1	1:B:401:LEU:HD13	1.74	0.51
1:D:20:LEU:HD22	1:D:315:ILE:HD13	1.93	0.51
1:C:64:TRP:N	1:C:65:PRO:CD	2.74	0.50
1:B:280:ALA:HB1	1:B:284:VAL:HG11	1.92	0.50
1:C:20:LEU:HB2	1:C:352:GLY:HA3	1.94	0.50



	lous pagem	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
4:B:501:GLC:O2	2:B:502:BGC:C2	2.52	0.50
1:B:15:PHE:CE1	1:B:401:LEU:HD13	2.47	0.49
1:C:305:THR:O	1:C:309:VAL:HG22	2.12	0.49
1:A:225:THR:HG22	1:A:263:GLY:HA3	1.95	0.49
1:B:313:PRO:HB3	1:B:351:ARG:HG3	1.95	0.49
1:B:170:ASN:HB3	1:B:252:PHE:CD2	2.48	0.49
1:C:8:ASP:OD1	1:C:11:THR:HG22	2.14	0.48
1:B:64:TRP:N	1:B:65:PRO:CD	2.76	0.48
1:C:253:ILE:HG23	1:C:309:VAL:CG1	2.36	0.48
1:A:64:TRP:N	1:A:65:PRO:CD	2.76	0.48
1:C:53:PRO:HG2	1:C:55:LEU:HD22	1.95	0.48
1:A:10:LYS:HE2	1:A:407:LEU:HD22	1.95	0.48
1:C:178:GLN:NE2	1:C:185:SER:O	2.47	0.48
1:C:8:ASP:OD1	1:C:11:THR:CG2	2.62	0.47
1:C:15:PHE:HE1	1:C:401:LEU:HD13	1.78	0.47
1:C:278:ILE:HG22	1:C:279:PRO:O	2.14	0.47
1:D:166:ILE:O	1:D:224:TRP:HB2	2.16	0.46
1:A:153:ALA:N	1:A:154:PRO:HD2	2.30	0.46
1:C:225:THR:HG22	1:C:263:GLY:HA3	1.97	0.46
1:D:289:THR:O	1:D:290:THR:HB	2.15	0.46
1:C:170:ASN:HB3	1:C:252:PHE:CD2	2.50	0.46
1:A:53:PRO:HB2	1:A:55:LEU:HD13	1.97	0.46
1:A:83:VAL:HG12	1:A:103:TYR:CD1	2.50	0.46
1:D:326:ASP:O	1:D:329:ARG:N	2.48	0.46
1:D:170:ASN:HB3	1:D:252:PHE:CD2	2.51	0.45
1:C:395:LEU:HA	1:C:398:THR:HB	1.97	0.45
1:D:333:TYR:CD2	1:D:392:LEU:HD11	2.52	0.45
1:C:105:ARG:NH2	6:C:606:HOH:O	2.49	0.45
1:D:83:VAL:HG12	1:D:103:TYR:CD1	2.52	0.45
1:A:35:THR:HA	1:A:40:TRP:CG	2.52	0.44
1:D:97:ARG:NH1	6:D:605:HOH:O	2.44	0.44
1:C:322:ALA:HB2	1:C:371:PHE:CE2	2.52	0.44
1:C:234:LEU:HB3	1:C:235:PRO:CD	2.48	0.44
1:A:313:PRO:HB3	1:A:351:ARG:HG3	2.01	0.43
1:A:363:GLU:OE2	2:A:502:BGC:O4	2.31	0.43
1:A:169:PRO:HD2	1:A:225:THR:O	2.19	0.43
1:D:64:TRP:N	1:D:65:PRO:CD	2.82	0.43
1:B:225:THR:HG22	1:B:263:GLY:HA3	2.00	0.43
1:C:12:PRO:HB3	1:C:400:GLU:HG2	2.01	0.42
1:D:10:LYS:HB2	1:D:11:THR:HG23	2.01	0.42
1:D:394:SER:O	1:D:396:GLY:N	2.53	0.42



Atom 1	Atom 2	Interatomic	Clash
Atom-1	Atom-2	$distance ( { m \AA} )$	overlap (Å)
1:C:15:PHE:CE1	1:C:401:LEU:HD13	2.54	0.42
1:D:153:ALA:N	1:D:154:PRO:HD2	2.35	0.42
1:B:83:VAL:HG12	1:B:103:TYR:CD1	2.54	0.42
1:D:253:ILE:CG2	1:D:309:VAL:HG13	2.50	0.42
1:D:35:THR:HA	1:D:40:TRP:CG	2.55	0.42
1:B:289:THR:O	1:B:290:THR:CB	2.56	0.42
1:D:12:PRO:HB3	1:D:400:GLU:HG2	2.01	0.41
1:C:265:GLN:HA	1:C:318:GLY:O	2.20	0.41
1:D:21:TRP:CZ2	1:D:395:LEU:HB2	2.55	0.41
1:A:265:GLN:HA	1:A:318:GLY:O	2.19	0.41
1:B:206:HIS:HB2	1:B:224:TRP:CZ3	2.56	0.41
1:A:181:ARG:O	1:B:211:LYS:HE2	2.21	0.41
1:D:257:ARG:HG3	1:D:309:VAL:O	2.20	0.41
1:C:356:TRP:HA	1:C:357:SER:HA	1.86	0.41
1:B:35:THR:HA	1:B:40:TRP:CG	2.56	0.40
1:D:214:ARG:HH12	1:D:260:ASP:CG	2.25	0.40
1:A:170:ASN:HB3	1:A:252:PHE:CD2	2.55	0.40
1:D:316:VAL:HG11	1:D:353:TYR:CE1	2.56	0.40
1:B:286:ARG:HD2	1:B:290:THR:O	2.21	0.40

There are no symmetry-related clashes.

### 5.3 Torsion angles (i)

#### 5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Perce	ntiles
1	А	399/414~(96%)	385~(96%)	14 (4%)	0	100	100
1	В	393/414~(95%)	381~(97%)	12 (3%)	0	100	100
1	С	398/414~(96%)	387~(97%)	11 (3%)	0	100	100
1	D	356/414~(86%)	339~(95%)	17 (5%)	0	100	100
All	All	1546/1656~(93%)	1492 (96%)	54 (4%)	0	100	100



There are no Ramachandran outliers to report.

#### 5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percen	tiles
1	А	310/322~(96%)	301~(97%)	9~(3%)	42	40
1	В	306/322~(95%)	296~(97%)	10 (3%)	38	35
1	С	310/322~(96%)	303~(98%)	7 (2%)	50	51
1	D	285/322~(88%)	276~(97%)	9~(3%)	39	37
All	All	1211/1288 (94%)	1176 (97%)	35 (3%)	42	40

All (35) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	10	LYS
1	А	55	LEU
1	А	73	SER
1	А	186	PHE
1	А	222	VAL
1	А	224	TRP
1	А	269	ARG
1	А	337	LEU
1	А	401	LEU
1	В	10	LYS
1	В	11	THR
1	В	55	LEU
1	В	224	TRP
1	В	269	ARG
1	В	284	VAL
1	В	309	VAL
1	В	337	LEU
1	В	401	LEU
1	В	404	THR
1	С	11	THR
1	С	55	LEU
1	С	73	SER



Mol	Chain	Res	Type
1	С	224	TRP
1	С	269	ARG
1	С	337	LEU
1	С	401	LEU
1	D	10	LYS
1	D	55	LEU
1	D	171	MET
1	D	224	TRP
1	D	269	ARG
1	D	284	VAL
1	D	337	LEU
1	D	383	ARG
1	D	398	THR

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	С	49	HIS

#### 5.3.3 RNA (i)

There are no RNA molecules in this entry.

### 5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

### 5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

### 5.6 Ligand geometry (i)

Of 20 ligands modelled in this entry, 6 are monoatomic - leaving 14 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mal	Tune	Chain	Dec	Tiple	Bo	Bond lengths			ond ang	gles
10101	туре	Unam	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z  > 2
2	BGC	С	504	-	11,11,12	0.54	0	$15,\!15,\!17$	1.16	1 (6%)
3	PGE	С	503	-	9,9,9	0.24	0	8,8,8	0.27	0
4	GLC	С	505	-	12,12,12	0.62	0	17,17,17	0.88	1 (5%)
2	BGC	С	501	-	11,11,12	0.65	0	15,15,17	2.30	3 (20%)
2	BGC	А	504	-	11,11,12	0.41	0	15,15,17	0.85	0
2	BGC	А	501	-	12,12,12	0.69	0	17,17,17	0.80	0
3	PGE	А	503	-	9,9,9	0.24	0	8,8,8	0.17	0
3	PGE	В	503	-	9,9,9	0.28	0	8,8,8	0.22	0
4	GLC	В	501	-	12,12,12	0.65	0	17,17,17	1.03	2 (11%)
4	GLC	А	505	-	12,12,12	0.32	0	17,17,17	0.42	0
2	BGC	D	501	-	12,12,12	0.42	0	17,17,17	1.52	3 (17%)
2	BGC	А	502	-	11,11,12	0.66	0	15,15,17	1.42	3 (20%)
2	BGC	В	502	-	11,11,12	0.61	0	15,15,17	1.08	1 (6%)
2	BGC	С	502	-	12,12,12	0.65	0	17,17,17	0.90	0

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	BGC	С	504	-	-	0/2/19/22	0/1/1/1
3	PGE	С	503	-	-	4/7/7/7	-
4	GLC	С	505	-	-	0/2/22/22	0/1/1/1
2	BGC	С	501	-	-	0/2/19/22	0/1/1/1
2	BGC	А	504	-	-	0/2/19/22	0/1/1/1
2	BGC	А	501	-	-	2/2/22/22	0/1/1/1
3	PGE	А	503	-	-	4/7/7/7	-
3	PGE	В	503	-	-	2/7/7/7	-
4	GLC	В	501	-	-	0/2/22/22	0/1/1/1
4	GLC	А	505	-	-	0/2/22/22	0/1/1/1
2	BGC	D	501	-	-	0/2/22/22	0/1/1/1
2	BGC	А	502	-	-	2/2/19/22	0/1/1/1
2	BGC	В	502	-	-	0/2/19/22	0/1/1/1
2	BGC	С	502	-	-	0/2/22/22	0/1/1/1



There are no bond length outliers.

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
2	С	501	BGC	C1-O5-C5	5.70	119.92	112.19
2	С	501	BGC	O5-C1-C2	5.04	118.54	110.77
2	С	501	BGC	C1-C2-C3	4.20	114.83	109.67
2	D	501	BGC	C1-O5-C5	4.05	121.31	113.66
2	А	502	BGC	C1-O5-C5	3.35	116.74	112.19
2	С	504	BGC	O2-C2-C3	3.16	116.47	110.14
2	А	502	BGC	C1-C2-C3	2.76	113.06	109.67
2	В	502	BGC	C1-C2-C3	2.61	112.88	109.67
2	А	502	BGC	O5-C1-C2	2.43	114.52	110.77
2	D	501	BGC	O5-C1-C2	2.36	114.50	110.28
4	С	505	GLC	O4-C4-C3	-2.29	105.06	110.35
2	D	501	BGC	C1-C2-C3	2.14	114.75	110.31
4	В	501	GLC	C1-O5-C5	2.07	117.58	113.66
4	B	501	GLC	C1-C2-C3	2.05	114.57	110.31

All (14) bond angle outliers are listed below:

There are no chirality outliers.

All (14) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	А	502	BGC	C4-C5-C6-O6
3	В	503	PGE	O2-C3-C4-O3
2	А	501	BGC	C4-C5-C6-O6
3	А	503	PGE	O2-C3-C4-O3
2	А	502	BGC	O5-C5-C6-O6
2	А	501	BGC	O5-C5-C6-O6
3	А	503	PGE	O1-C1-C2-O2
3	С	503	PGE	O3-C5-C6-O4
3	А	503	PGE	O3-C5-C6-O4
3	В	503	PGE	O1-C1-C2-O2
3	С	503	PGE	O1-C1-C2-O2
3	А	503	PGE	C3-C4-O3-C5
3	С	503	PGE	C4-C3-O2-C2
3	С	503	PGE	O2-C3-C4-O3

There are no ring outliers.

5 monomers are involved in 11 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	А	501	BGC	3	0
				<i>a</i>	1 ,



	0	-	1 0		
Mol	Chain	Res	Type	Clashes	Symm-Clashes
3	А	503	PGE	2	0
4	В	501	GLC	5	0
2	А	502	BGC	4	0
2	В	502	BGC	5	0

Continued from previous page...

The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and similar rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.











































## 5.7 Other polymers (i)

There are no such residues in this entry.

## 5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



# 6 Fit of model and data (i)

## 6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median,  $95^{th}$  percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	< <b>RSRZ</b> >	#RSR	RZ>2	$OWAB(Å^2)$	Q<0.9
1	А	401/414~(96%)	-0.21	7 (1%) 7	70 75	17, 23, 45, 79	0
1	В	395/414~(95%)	-0.23	2 (0%) 9	91 93	16, 26, 44, 83	0
1	С	400/414~(96%)	-0.21	1 (0%) 9	94 95	16, 24, 48, 63	0
1	D	364/414~(87%)	0.39	30 (8%)	11 15	18, 36, 63, 95	0
All	All	1560/1656~(94%)	-0.07	40 (2%) 3	56 62	16, 26, 52, 95	0

All (40) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	405	ALA	4.2
1	D	11	THR	3.8
1	D	296	THR	3.6
1	В	282	ASP	3.6
1	D	294	TYR	3.5
1	В	10	LYS	3.2
1	D	401	LEU	3.2
1	D	270	THR	3.2
1	D	284	VAL	3.0
1	С	273	GLY	3.0
1	А	409	HIS	3.0
1	D	307	ALA	3.0
1	А	282	ASP	2.8
1	D	10	LYS	2.8
1	D	180	LYS	2.8
1	D	283	ASP	2.8
1	А	235	PRO	2.8
1	D	231	TYR	2.8
1	D	404	THR	2.7
1	D	347	GLY	2.6
1	D	309	VAL	2.6



Mol	Chain	Res	Type	RSRZ
1	D	244	TYR	2.6
1	D	399	ARG	2.6
1	А	281	ALA	2.6
1	D	397	ARG	2.5
1	D	336	ALA	2.4
1	D	193	THR	2.4
1	D	14	PRO	2.4
1	D	287	THR	2.3
1	D	290	THR	2.2
1	А	275	GLU	2.1
1	D	251	VAL	2.1
1	D	15	PHE	2.1
1	D	190	GLY	2.1
1	D	189	ALA	2.1
1	D	293	TYR	2.1
1	А	188	PRO	2.1
1	D	391	TRP	2.0
1	D	331	ASP	2.0
1	D	400	GLU	2.0

### 6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

#### 6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

#### 6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median,  $95^{th}$  percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-factors}(\mathbf{A}^2)$	Q<0.9
2	BGC	С	502	12/12	0.84	0.21	42,47,52,57	0
2	BGC	А	501	12/12	0.85	0.19	$45,\!53,\!58,\!58$	0
2	BGC	В	502	11/12	0.86	0.12	30,37,44,46	0
3	PGE	А	503	10/10	0.86	0.17	$38,\!51,\!56,\!57$	0



Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B$ -factors( $Å^2$ )	Q<0.9
2	BGC	D	501	12/12	0.88	0.12	$35,\!45,\!52,\!53$	0
3	PGE	В	503	10/10	0.88	0.20	33,36,40,41	0
2	BGC	С	501	11/12	0.89	0.15	29,40,45,47	0
3	PGE	С	503	10/10	0.90	0.19	30,31,33,35	0
2	BGC	А	502	11/12	0.91	0.16	$23,\!40,\!47,\!50$	0
4	GLC	В	501	12/12	0.91	0.15	33,41,44,45	0
5	CL	В	505	1/1	0.92	0.25	59, 59, 59, 59, 59	0
4	GLC	А	505	12/12	0.93	0.14	$29,\!33,\!35,\!36$	0
2	BGC	С	504	11/12	0.94	0.16	27,32,34,35	0
4	GLC	С	505	12/12	0.94	0.18	35,37,39,39	0
2	BGC	А	504	11/12	0.94	0.13	31,34,38,41	0
5	CL	В	504	1/1	0.95	0.18	$50,\!50,\!50,\!50$	0
5	CL	А	507	1/1	0.96	0.16	60,60,60,60	0
5	CL	D	502	1/1	0.96	0.14	42,42,42,42	0
5	CL	C	506	1/1	0.99	0.07	44,44,44,44	0
5	CL	А	506	1/1	0.99	0.12	44,44,44	0

The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.













































## 6.5 Other polymers (i)

There are no such residues in this entry.

