



Full wwPDB X-ray Structure Validation Report ⓘ

Mar 8, 2018 – 10:11 pm GMT

PDB ID : 3KD8
Title : Cofactor-Independent Phosphoglycerate mutase from *Thermoplasma Acidophilum* DSM 1728
Authors : Joachimiak, A.; Duke, N.E.C.; Marshall, N.; Buck, K.; Midwest Center for Structural Genomics (MCSG)
Deposited on : 2009-10-22
Resolution : 2.60 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : trunk30967
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
Refmac : 5.8.0158
CCP4 : 7.0 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30967


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 2.60 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	111664	2767 (2.60-2.60)
Clashscore	122126	3110 (2.60-2.60)
Ramachandran outliers	120053	3062 (2.60-2.60)
Sidechain outliers	120020	3062 (2.60-2.60)
RSRZ outliers	108989	2706 (2.60-2.60)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	399	<p>2% 68% 19% 8%</p>
1	B	399	<p>2% 69% 18% 9%</p>

2 Entry composition [i](#)

There are 2 unique types of molecules in this entry. The entry contains 5761 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called 2,3-bisphosphoglycerate-independent phosphoglycerate mutase.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace	
			Total	C	N	O	S				Se
1	A	367	2817	1758	508	535	3	13	0	0	0
1	B	365	2799	1748	506	529	3	13	0	0	0

There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-2	SER	-	expression tag	UNP Q9HL27
A	-1	ASN	-	expression tag	UNP Q9HL27
A	0	ALA	-	expression tag	UNP Q9HL27
B	-2	SER	-	expression tag	UNP Q9HL27
B	-1	ASN	-	expression tag	UNP Q9HL27
B	0	ALA	-	expression tag	UNP Q9HL27


- Molecule 2 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
2	A	75	Total 75	O 75	0	0
2	B	70	Total 70	O 70	0	0


3 Residue-property plots [i](#)

These plots are drawn for all protein, RNA and DNA chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: 2,3-bisphosphoglycerate-independent phosphoglycerate mutase



- Molecule 1: 2,3-bisphosphoglycerate-independent phosphoglycerate mutase



4 Data and refinement statistics

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	87.91Å 137.08Å 67.09Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	73.92 – 2.60 33.54 – 2.60	Depositor EDS
% Data completeness (in resolution range)	99.8 (73.92-2.60) 99.9 (33.54-2.60)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	4.47 (at 2.61Å)	Xtrriage
Refinement program	REFMAC 5.5.0102	Depositor
R, R_{free}	0.207 , 0.288 0.210 , 0.285	Depositor DCC
R_{free} test set	1306 reflections (5.09%)	wwPDB-VP
Wilson B-factor (Å ²)	38.3	Xtrriage
Anisotropy	0.523	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.36 , 44.5	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	5761	wwPDB-VP
Average B, all atoms (Å ²)	33.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.78% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.69	0/2856	0.82	5/3837 (0.1%)
1	B	0.69	1/2837 (0.0%)	0.80	1/3810 (0.0%)
All	All	0.69	1/5693 (0.0%)	0.81	6/7647 (0.1%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
1	B	356	VAL	CB-CG1	-6.00	1.40	1.52

All (6) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	221	ARG	NE-CZ-NH2	-5.70	117.45	120.30
1	A	195	ARG	CG-CD-NE	-5.64	99.97	111.80
1	A	356	VAL	CB-CA-C	-5.62	100.72	111.40
1	A	221	ARG	NE-CZ-NH1	5.33	122.96	120.30
1	A	177	ASP	CB-CG-OD1	5.11	122.89	118.30
1	B	23	ARG	NE-CZ-NH2	-5.00	117.80	120.30

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2817	0	2804	52	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	B	2799	0	2785	62	0
2	A	75	0	0	2	0
2	B	70	0	0	2	0
All	All	5761	0	5589	114	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

All (114) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:80:GLU:HG3	1:B:96:ARG:HG2	1.41	1.03
1:A:103:ARG:HH11	1:A:103:ARG:HG3	1.23	1.02
1:A:149:VAL:HG21	1:A:187:MSE:HE1	1.42	1.00
1:A:80:GLU:HG3	1:A:96:ARG:HG2	1.44	0.96
1:A:14:ASP:O	1:A:15:ARG:HB2	1.66	0.94
1:B:103:ARG:HG3	1:B:103:ARG:HH11	1.32	0.91
1:B:120:GLU:O	1:B:121:GLU:HB2	1.72	0.90
1:B:249:LEU:O	1:B:250:LYS:HB2	1.71	0.89
1:A:278:ILE:O	1:A:282:VAL:HG23	1.74	0.88
1:B:249:LEU:O	1:B:250:LYS:CB	2.21	0.84
1:B:244:VAL:HG11	1:B:249:LEU:HB3	1.61	0.83
1:B:80:GLU:O	1:B:81:ALA:CB	2.29	0.81
1:B:244:VAL:HG11	1:B:249:LEU:CB	2.12	0.79
1:B:90:PRO:HA	1:B:226:VAL:HG22	1.65	0.78
1:A:80:GLU:HG3	1:A:96:ARG:CG	2.17	0.73
1:B:103:ARG:HD2	2:B:455:HOH:O	1.90	0.70
1:B:48:PRO:O	1:B:49:ILE:HB	1.90	0.69
1:B:80:GLU:HG3	1:B:96:ARG:CG	2.22	0.69
1:B:351:ASP:N	1:B:352:PRO:HD2	2.08	0.69
1:A:306:ASN:HD22	1:A:306:ASN:C	1.97	0.67
1:B:13:GLY:HA2	1:B:372:GLU:OE2	1.94	0.67
1:A:141:VAL:HG23	1:A:144:ARG:HH11	1.59	0.67
1:A:394:LEU:O	1:A:395:ALA:HB3	1.94	0.67
1:A:394:LEU:O	1:A:395:ALA:CB	2.43	0.67
1:B:238:MSE:HE1	1:B:388:MSE:SE	2.45	0.67
1:A:350:GLY:O	1:A:351:ASP:HB2	1.94	0.66
1:B:90:PRO:HB3	1:B:227:PRO:O	1.95	0.66
1:A:141:VAL:HG23	1:A:144:ARG:NH1	2.11	0.65
1:B:336:THR:HB	1:B:355:ILE:HG13	1.79	0.65
1:B:236:ASN:HB3	1:B:238:MSE:HE2	1.79	0.64

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:103:ARG:NH1	1:A:103:ARG:HG3	1.99	0.63
1:B:80:GLU:O	1:B:81:ALA:HB3	1.96	0.63
1:B:351:ASP:N	1:B:352:PRO:CD	2.63	0.62
1:A:149:VAL:CG2	1:A:187:MSE:HE1	2.24	0.61
1:A:321:MSE:HG2	1:A:324:LEU:HD12	1.84	0.60
1:B:319:ARG:NH1	2:B:432:HOH:O	2.37	0.57
1:A:31:ARG:NH1	1:A:370:PHE:O	2.37	0.57
1:B:241:ALA:HB2	1:B:288:HIS:CD2	2.40	0.56
1:B:244:VAL:HG11	1:B:249:LEU:HB2	1.85	0.56
1:B:334:CYS:HB2	1:B:391:LEU:HD21	1.87	0.56
1:B:176:THR:O	1:B:177:ASP:HB3	2.06	0.56
1:A:95:PHE:CD1	1:A:220:VAL:HG13	2.41	0.55
1:B:33:ASN:ND2	1:B:318:ASP:O	2.28	0.55
1:B:48:PRO:HB3	1:B:60:SER:OG	2.07	0.54
1:B:351:ASP:H	1:B:352:PRO:CD	2.22	0.53
1:B:296:LYS:CB	1:B:296:LYS:NZ	2.72	0.53
1:A:46:MSE:HE3	1:A:48:PRO:HG3	1.89	0.52
1:A:356:VAL:HG22	1:A:370:PHE:CE1	2.45	0.52
1:B:103:ARG:NH1	1:B:103:ARG:HG3	2.11	0.51
1:A:24:THR:OG1	1:A:27:GLN:HG3	2.10	0.51
1:A:59:THR:O	1:A:60:SER:HB2	2.10	0.51
1:A:98:ASN:OD1	1:A:221:ARG:NH2	2.43	0.51
1:A:324:LEU:HD22	1:A:333:ILE:HD12	1.92	0.51
1:A:327:ILE:HD11	1:A:333:ILE:HD11	1.91	0.51
1:A:277:LYS:NZ	1:A:294:ASN:O	2.43	0.51
1:B:173:ILE:HG13	1:B:187:MSE:HB3	1.92	0.51
1:B:5:ILE:HD12	1:B:281:ALA:HB1	1.94	0.50
1:A:2:LYS:HB2	1:A:395:ALA:HA	1.93	0.50
1:A:110:ARG:NH1	1:A:162:ASP:OD2	2.45	0.50
1:A:306:ASN:ND2	1:A:308:PRO:HD2	2.26	0.50
1:B:80:GLU:O	1:B:81:ALA:HB2	2.11	0.50
1:A:203:VAL:O	1:A:207:ARG:HG3	2.12	0.49
1:B:134:SER:OG	1:B:150:SER:HB2	2.12	0.49
1:B:385:TYR:O	1:B:385:TYR:CG	2.65	0.49
1:B:15:ARG:O	1:B:307:TYR:OH	2.24	0.49
1:B:94:ALA:HA	1:B:147:LEU:O	2.12	0.49
1:A:11:GLY:HA3	2:A:445:HOH:O	2.13	0.49
1:B:79:PHE:CE2	1:B:248:TRP:CD1	3.01	0.48
1:B:79:PHE:HE2	1:B:248:TRP:CD1	2.32	0.48
1:B:111:ARG:HD3	1:B:143:HIS:CE1	2.49	0.48
1:A:177:ASP:OD1	1:A:177:ASP:C	2.52	0.48

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:246:SER:HB2	1:A:249:LEU:HB2	1.97	0.47
1:B:5:ILE:HD13	1:B:291:VAL:HB	1.96	0.47
1:B:276:GLY:O	1:B:280:LYS:HB2	2.15	0.47
1:B:44:GLY:HA3	1:B:380:LEU:HB2	1.97	0.47
1:A:73:TYR:HA	1:A:74:PRO:HD3	1.71	0.47
1:A:104:ASP:N	2:A:463:HOH:O	2.28	0.46
1:A:95:PHE:CE1	1:A:223:ALA:HB2	2.51	0.46
1:B:138:LYS:HE2	1:B:138:LYS:HB2	1.75	0.46
1:B:296:LYS:HB3	1:B:296:LYS:HZ3	1.81	0.46
1:A:334:CYS:HB2	1:A:391:LEU:HD21	1.99	0.44
1:B:306:ASN:N	1:B:308:PRO:HD2	2.32	0.44
1:A:97:ALA:HB1	1:A:218:LEU:HD11	1.99	0.44
1:A:45:ILE:HG12	1:A:376:ALA:HA	2.00	0.44
1:A:68:ASP:HA	1:A:69:PRO:HD3	1.84	0.43
1:A:72:TYR:HD1	1:A:229:ILE:HD13	1.84	0.43
1:B:29:ALA:HB1	1:B:315:GLU:HG2	1.99	0.43
1:B:31:ARG:NH1	1:B:371:ASP:HB3	2.34	0.43
1:B:67:TYR:CZ	1:B:256:LEU:HD22	2.54	0.43
1:A:391:LEU:O	1:A:394:LEU:O	2.36	0.43
1:B:120:GLU:O	1:B:121:GLU:CB	2.55	0.43
1:A:306:ASN:ND2	1:A:306:ASN:C	2.69	0.42
1:A:111:ARG:HG2	1:A:143:HIS:CD2	2.54	0.42
1:B:252:LEU:HD22	1:B:256:LEU:HD11	2.02	0.42
1:A:80:GLU:CG	1:A:96:ARG:HE	2.33	0.42
1:B:226:VAL:HA	1:B:227:PRO:HD3	1.75	0.42
1:A:41:GLY:HA2	1:A:362:VAL:HG11	2.02	0.42
1:B:73:TYR:CD1	1:B:74:PRO:HD2	2.55	0.42
1:B:4:ILE:HD13	1:B:391:LEU:HB3	2.01	0.42
1:A:49:ILE:HD11	1:A:60:SER:HA	2.01	0.42
1:A:12:LEU:HB3	1:A:13:GLY:H	1.74	0.41
1:B:5:ILE:HD11	1:B:285:THR:CG2	2.51	0.41
1:A:203:VAL:O	1:A:207:ARG:CG	2.68	0.41
1:A:351:ASP:N	1:A:352:PRO:HD2	2.35	0.41
1:B:141:VAL:O	1:B:144:ARG:HB2	2.19	0.41
1:B:25:PRO:HD3	1:B:307:TYR:CE2	2.55	0.41
1:B:248:TRP:C	1:B:249:LEU:O	2.56	0.41
1:B:73:TYR:HA	1:B:74:PRO:HD3	1.90	0.41
1:A:306:ASN:ND2	1:A:309:LEU:H	2.19	0.41
1:A:97:ALA:HA	1:A:219:LEU:O	2.21	0.41
1:B:89:ARG:O	1:B:226:VAL:HG11	2.21	0.41
1:A:72:TYR:HB3	1:A:229:ILE:HD12	2.03	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:B:111:ARG:HG2	1:B:143:HIS:CD2	2.56	0.40
1:B:321:MSE:HB3	1:B:324:LEU:HD12	2.02	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	357/399 (90%)	334 (94%)	17 (5%)	6 (2%)	10	20
1	B	355/399 (89%)	334 (94%)	13 (4%)	8 (2%)	7	13
All	All	712/798 (89%)	668 (94%)	30 (4%)	14 (2%)	8	16

All (14) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	60	SER
1	A	351	ASP
1	B	59	THR
1	B	81	ALA
1	B	121	GLU
1	A	12	LEU
1	A	49	ILE
1	A	395	ALA
1	A	15	ARG
1	B	112	ALA
1	B	250	LYS
1	B	14	ASP
1	B	177	ASP
1	B	351	ASP

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	307/318 (96%)	272 (89%)	35 (11%)	6	11
1	B	303/318 (95%)	278 (92%)	25 (8%)	12	24
All	All	610/636 (96%)	550 (90%)	60 (10%)	9	16

All (60) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	9	LEU
1	A	12	LEU
1	A	14	ASP
1	A	18	SER
1	A	31	ARG
1	A	39	SER
1	A	51	PRO
1	A	62	MSE
1	A	63	SER
1	A	70	LYS
1	A	79	PHE
1	A	84	LEU
1	A	103	ARG
1	A	138	LYS
1	A	150	SER
1	A	177	ASP
1	A	192	GLU
1	A	200	ASP
1	A	202	ARG
1	A	212	ARG
1	A	220	VAL
1	A	237	ARG
1	A	250	LYS
1	A	252	LEU
1	A	257	ARG
1	A	306	ASN
1	A	313	VAL

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	321	MSE
1	A	353	VAL
1	A	356	VAL
1	A	360	ASP
1	A	362	VAL
1	A	381	ARG
1	A	393	GLN
1	A	394	LEU
1	B	60	SER
1	B	89	ARG
1	B	103	ARG
1	B	120	GLU
1	B	121	GLU
1	B	138	LYS
1	B	141	VAL
1	B	150	SER
1	B	177	ASP
1	B	200	ASP
1	B	202	ARG
1	B	212	ARG
1	B	220	VAL
1	B	221	ARG
1	B	226	VAL
1	B	237	ARG
1	B	246	SER
1	B	248	TRP
1	B	252	LEU
1	B	296	LYS
1	B	307	TYR
1	B	322	GLU
1	B	329	ASP
1	B	353	VAL
1	B	364	ASN

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (3) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	119	ASN
1	A	210	ASN
1	A	306	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data [i](#)

6.1 Protein, DNA and RNA chains [i](#)

In the following table, the column labelled ‘#RSRZ > 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q < 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	354/399 (88%)	-0.27	9 (2%) 57 50	17, 32, 48, 67	0
1	B	352/399 (88%)	-0.18	7 (1%) 65 59	17, 33, 50, 62	0
All	All	706/798 (88%)	-0.22	16 (2%) 60 54	17, 32, 49, 67	0

All (16) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	B	271	GLY	5.8
1	A	51	PRO	4.2
1	B	266	ALA	4.1
1	B	396	GLY	3.7
1	A	50	SER	3.5
1	B	265	GLY	3.4
1	B	79	PHE	3.2
1	A	305	GLY	3.0
1	A	202	ARG	3.0
1	A	298	THR	2.6
1	A	297	ALA	2.6
1	A	350	GLY	2.5
1	A	385	TYR	2.3
1	A	49	ILE	2.2
1	B	58	ASP	2.2
1	B	73	TYR	2.2

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.