



wwPDB X-ray Structure Validation Summary Report ⓘ

Aug 9, 2020 – 01:55 PM BST

PDB ID : 1KEN
Title : INFLUENZA VIRUS HEMAGGLUTININ COMPLEXED WITH AN AN-
TIBODY THAT PREVENTS THE HEMAGGLUTININ LOW PH FUSO-
GENIC TRANSITION
Authors : Barbey-Martin, C.; Gigant, B.; Bizebard, T.; Calder, L.J.; Wharto, S.A.;
Skehel, J.J.; Knossow, M.
Deposited on : 2001-11-16
Resolution : 3.50 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Mogul : 1.8.5 (274361), CSD as541be (2020)
Xtriage (Phenix) : **NOT EXECUTED**
EDS : **NOT EXECUTED**
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.13.1


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 3.50 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
Clashscore	141614	1036 (3.58-3.42)
Ramachandran outliers	138981	1005 (3.58-3.42)
Sidechain outliers	138945	1006 (3.58-3.42)





The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$.

Note EDS was not executed.

Mol	Chain	Length	Quality of chain
1	A	328	30% 53% 13% ..
1	C	328	32% 52% 13% ..
1	E	328	27% 57% 13% ..
2	B	175	18% 57% 22% .
2	D	175	15% 51% 32% .
2	F	175	16% 56% 26% .
3	L	213	21% 55% 21% .
3	U	213	20% 60% 19% .

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
4	H	221	
4	T	221	
5	G	3	
5	I	3	
5	J	3	

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:

Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
5	MAN	G	3	X	-	-	-
5	MAN	I	3	X	-	-	-
5	MAN	J	3	X	-	-	-

2 Entry composition [i](#)

There are 5 unique types of molecules in this entry. The entry contains 18492 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called hemagglutinin HA1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	320	2472	1547	434	478	13	0	0	0
1	C	320	2472	1547	434	478	13	0	0	0
1	E	320	2472	1547	434	478	13	0	0	0

- Molecule 2 is a protein called hemagglutinin HA2.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
2	B	175	1421	882	250	283	6	0	0	0
2	D	175	1421	882	250	283	6	0	0	0
2	F	175	1421	882	250	283	6	0	0	0

- Molecule 3 is a protein called influenza virus infectivity neutralizing antibody (light chain).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
3	L	213	1638	1028	272	332	6	0	0	0
3	U	213	1638	1028	272	332	6	0	0	0

- Molecule 4 is a protein called influenza virus infectivity neutralizing antibody (heavy chain).

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
4	H	221	1720	1102	272	340	6	0	0	0
4	T	219	1700	1092	267	335	6	0	0	0

- Molecule 5 is an oligosaccharide called alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
5	G	3	39	22	2	15	0	0	0
5	I	3	39	22	2	15	0	0	0
5	J	3	39	22	2	15	0	0	0


3 Residue-property plots

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

Note EDS was not executed.


- Molecule 1: hemagglutinin HA1


Chain A: 




- Molecule 1: hemagglutinin HA1

Chain C: 









• Molecule 1: hemagglutinin HA1



• Molecule 2: hemagglutinin HA2




• Molecule 2: hemagglutinin HA2






- Molecule 2: hemagglutinin HA2

Chain F: 16% 56% 26%




- Molecule 3: influenza virus infectivity neutralizing antibody (light chain)

Chain L: 21% 55% 21%





- Molecule 3: influenza virus infectivity neutralizing antibody (light chain)

Chain U: 20% 60% 19%




- Molecule 4: influenza virus infectivity neutralizing antibody (heavy chain)

Chain H:  20% 58% 18%



- Molecule 4: influenza virus infectivity neutralizing antibody (heavy chain)

Chain T:  19% 59% 18%



- Molecule 5: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:  67% 33%

MAG1
MAG2
MAN3

- Molecule 5: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain I:  33% 67%

MAG1
MAG2
MAN3

- Molecule 5: alpha-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain J:  100%

MAC1
MAC2
MAIN3

4 Data and refinement statistics

Xtrriage (Phenix) and EDS were not executed - this section is therefore incomplete.

Property	Value	Source
Space group	P 21 21 2	Depositor
Cell constants a, b, c, α , β , γ	143.04Å 315.59Å 97.03Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	25.00 – 3.50	Depositor
% Data completeness (in resolution range)	91.8 (25.00-3.50)	Depositor
R_{merge}	(Not available)	Depositor
R_{sym}	0.13	Depositor
Refinement program	CNS 1.0	Depositor
R, R_{free}	0.255 , 0.323	Depositor
Estimated twinning fraction	No twinning to report.	Xtrriage
Total number of atoms	18492	wwPDB-VP
Average B, all atoms (Å ²)	47.0	wwPDB-VP

5 Model quality i

5.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, MAN

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.71	0/2528	0.91	3/3443 (0.1%)
1	C	0.77	1/2528 (0.0%)	0.96	2/3443 (0.1%)
1	E	0.71	0/2528	0.93	2/3443 (0.1%)
2	B	0.74	0/1445	0.86	0/1939
2	D	0.77	1/1445 (0.1%)	0.89	3/1939 (0.2%)
2	F	0.73	0/1445	0.84	0/1939
3	L	0.82	2/1679 (0.1%)	1.05	7/2281 (0.3%)
3	U	0.76	0/1679	0.97	3/2281 (0.1%)
4	H	0.85	2/1774 (0.1%)	1.00	2/2431 (0.1%)
4	T	0.78	1/1754 (0.1%)	1.00	2/2406 (0.1%)
All	All	0.76	7/18805 (0.0%)	0.95	24/25545 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
3	L	0	1
3	U	0	1
4	H	0	1
4	T	0	2
All	All	0	5

The worst 5 of 7 bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
4	H	1	ASP	CB-CG	7.47	1.67	1.51
3	L	155	GLU	CB-CG	7.10	1.65	1.52
3	L	155	GLU	CG-CD	6.60	1.61	1.51
4	H	61	ASN	CB-CG	5.86	1.64	1.51

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	D	92	TRP	CB-CG	-5.42	1.40	1.50

The worst 5 of 24 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
4	T	64	LEU	CA-CB-CG	-10.84	90.37	115.30
3	L	111	ASP	N-CA-C	-9.85	84.40	111.00
4	H	64	LEU	CA-CB-CG	-7.90	97.13	115.30
3	U	111	ASP	N-CA-C	-7.69	90.24	111.00
2	D	79	ASP	CB-CG-OD2	7.20	124.78	118.30

There are no chirality outliers.

All (5) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
4	H	101	TYR	Sidechain
3	L	141	TYR	Sidechain
4	T	33	TYR	Sidechain
4	T	95	TYR	Sidechain
3	U	72	TYR	Sidechain

5.2 Too-close contacts [\(i\)](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	2472	0	2424	336	0
1	C	2472	0	2424	328	0
1	E	2472	0	2424	334	0
2	B	1421	0	1346	259	0
2	D	1421	0	1346	273	0
2	F	1421	0	1346	276	0
3	L	1638	0	1578	232	0
3	U	1638	0	1578	260	0
4	H	1720	0	1639	264	0
4	T	1700	0	1622	276	0
5	G	39	0	34	3	0
5	I	39	0	34	4	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
5	J	39	0	34	0	0
All	All	18492	0	17829	2549	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 70.

The worst 5 of 2549 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:74:PRO:HA	1:A:141:ARG:HH12	1.11	1.16
3:U:199:HIS:HB3	3:U:201:THR:HG22	1.28	1.11
3:U:134:VAL:HG22	3:U:179:THR:HG23	1.30	1.11
1:C:77:ASP:O	1:C:80:GLN:HG2	1.47	1.10
1:C:96:ASN:HD21	1:C:140:LYS:HE2	1.10	1.10

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	318/328 (97%)	247 (78%)	43 (14%)	28 (9%)	1 8
1	C	318/328 (97%)	241 (76%)	53 (17%)	24 (8%)	1 11
1	E	318/328 (97%)	250 (79%)	41 (13%)	27 (8%)	1 9
2	B	173/175 (99%)	96 (56%)	44 (25%)	33 (19%)	0 2
2	D	173/175 (99%)	89 (51%)	48 (28%)	36 (21%)	0 1
2	F	173/175 (99%)	89 (51%)	49 (28%)	35 (20%)	0 1
3	L	211/213 (99%)	141 (67%)	46 (22%)	24 (11%)	0 6
3	U	211/213 (99%)	139 (66%)	51 (24%)	21 (10%)	0 7

Continued on next page...

Continued from previous page...

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
4	H	219/221 (99%)	158 (72%)	38 (17%)	23 (10%)	0	7
4	T	217/221 (98%)	147 (68%)	47 (22%)	23 (11%)	0	7
All	All	2331/2377 (98%)	1597 (68%)	460 (20%)	274 (12%)	0	5

5 of 274 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	29	ILE
1	A	81	ASN
1	A	210	GLN
1	A	263	GLY
1	A	287	SER

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	282/289 (98%)	252 (89%)	30 (11%)	6	30
1	C	282/289 (98%)	249 (88%)	33 (12%)	5	26
1	E	282/289 (98%)	248 (88%)	34 (12%)	5	24
2	B	149/149 (100%)	120 (80%)	29 (20%)	1	7
2	D	149/149 (100%)	116 (78%)	33 (22%)	1	5
2	F	149/149 (100%)	118 (79%)	31 (21%)	1	6
3	L	187/187 (100%)	147 (79%)	40 (21%)	1	5
3	U	187/187 (100%)	152 (81%)	35 (19%)	1	8
4	H	196/196 (100%)	163 (83%)	33 (17%)	2	12
4	T	194/196 (99%)	167 (86%)	27 (14%)	3	20
All	All	2057/2080 (99%)	1732 (84%)	325 (16%)	2	15

5 of 325 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	E	239	PRO
2	F	123	ARG
4	T	16	GLN
1	E	250	ASN
2	F	21	TRP

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. 5 of 102 such sidechains are listed below:

Mol	Chain	Res	Type
2	D	28	ASN
1	E	80	GLN
4	T	58	ASN
2	D	49	ASN
2	D	154	ASN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

9 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 2$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# $ Z > 2$	Counts	RMSZ	# $ Z > 2$
5	NAG	G	1	1,5	14,14,15	0.68	0	17,19,21	1.40	2 (11%)
5	NAG	G	2	5	14,14,15	0.51	0	17,19,21	0.93	1 (5%)
5	MAN	G	3	5	11,11,12	1.00	1 (9%)	15,15,17	0.94	1 (6%)
5	NAG	I	1	1,5	14,14,15	0.65	0	17,19,21	1.04	2 (11%)

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	NAG	I	2	5	14,14,15	0.71	0	17,19,21	0.96	1 (5%)
5	MAN	I	3	5	11,11,12	1.16	2 (18%)	15,15,17	1.04	1 (6%)
5	NAG	J	1	1,5	14,14,15	0.80	0	17,19,21	1.26	2 (11%)
5	NAG	J	2	5	14,14,15	1.19	1 (7%)	17,19,21	1.33	3 (17%)
5	MAN	J	3	5	11,11,12	0.89	0	15,15,17	1.12	1 (6%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	G	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	G	2	5	-	3/6/23/26	0/1/1/1
5	MAN	G	3	5	1/1/4/5	1/2/19/22	0/1/1/1
5	NAG	I	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	I	2	5	-	3/6/23/26	0/1/1/1
5	MAN	I	3	5	1/1/4/5	2/2/19/22	0/1/1/1
5	NAG	J	1	1,5	-	2/6/23/26	0/1/1/1
5	NAG	J	2	5	-	2/6/23/26	0/1/1/1
5	MAN	J	3	5	1/1/4/5	0/2/19/22	0/1/1/1

All (4) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
5	J	2	NAG	C1-C2	-3.04	1.47	1.52
5	G	3	MAN	C4-C5	2.20	1.57	1.53
5	I	3	MAN	C4-C5	2.07	1.57	1.53
5	I	3	MAN	C6-C5	2.04	1.58	1.51

The worst 5 of 14 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
5	G	1	NAG	C2-N2-C7	-3.74	117.58	122.90
5	G	1	NAG	C4-C3-C2	-3.21	106.31	111.02
5	I	3	MAN	C6-C5-C4	3.01	120.06	113.00
5	J	3	MAN	C6-C5-C4	2.82	119.61	113.00
5	J	2	NAG	C4-C3-C2	2.81	115.14	111.02

All (3) chirality outliers are listed below:

Mol	Chain	Res	Type	Atom
5	J	3	MAN	C1
5	G	3	MAN	C1
5	I	3	MAN	C1

5 of 17 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	J	1	NAG	O5-C5-C6-O6
5	J	2	NAG	C8-C7-N2-C2
5	I	1	NAG	O5-C5-C6-O6
5	J	1	NAG	C4-C5-C6-O6
5	G	2	NAG	C8-C7-N2-C2

There are no ring outliers.

3 monomers are involved in 7 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	G	2	NAG	3	0
5	I	1	NAG	2	0
5	I	2	NAG	2	0

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

EDS was not executed - this section is therefore empty.

6.2 Non-standard residues in protein, DNA, RNA chains

EDS was not executed - this section is therefore empty.

6.3 Carbohydrates

EDS was not executed - this section is therefore empty.

6.4 Ligands

EDS was not executed - this section is therefore empty.

6.5 Other polymers

EDS was not executed - this section is therefore empty.