



Full wwPDB X-ray Structure Validation Report ⓘ

Oct 17, 2021 – 05:41 AM EDT

PDB ID : 1KL3
Title : an engineered streptavidin with improved affinity for the strep-tag II peptide
: SAm1-StrepII
Authors : Korndorfer, I.P.; Skerra, A.
Deposited on : 2001-12-11
Resolution : 1.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.23.2
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.23.2


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.






Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	4298 (1.70-1.70)
Clashscore	141614	4695 (1.70-1.70)
Ramachandran outliers	138981	4610 (1.70-1.70)
Sidechain outliers	138945	4610 (1.70-1.70)
RSRZ outliers	127900	4222 (1.70-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	127	
1	B	127	
1	C	127	
1	D	127	
2	E	9	

Continued on next page...

Continued from previous page...

Mol	Chain	Length	Quality of chain
2	F	9	 11% 44% 11% 11% 33%
2	G	9	 11% 44% 22% 33%
2	H	9	 11% 11% 56% 33%

2 Entry composition i

There are 3 unique types of molecules in this entry. The entry contains 4054 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called streptavidin.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
			Total	C	N	O			
1	A	121	907	567	160	180	0	0	0
1	B	120	899	562	159	178	0	0	0
1	C	120	899	562	159	178	0	0	0
1	D	119	894	559	158	177	0	0	0

There are 16 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	13	MET	-	initiating methionine	UNP P22629
A	44	VAL	GLU	engineered mutation	UNP P22629
A	45	THR	SER	engineered mutation	UNP P22629
A	47	ARG	VAL	engineered mutation	UNP P22629
B	13	MET	-	initiating methionine	UNP P22629
B	44	VAL	GLU	engineered mutation	UNP P22629
B	45	THR	SER	engineered mutation	UNP P22629
B	47	ARG	VAL	engineered mutation	UNP P22629
C	13	MET	-	initiating methionine	UNP P22629
C	44	VAL	GLU	engineered mutation	UNP P22629
C	45	THR	SER	engineered mutation	UNP P22629
C	47	ARG	VAL	engineered mutation	UNP P22629
D	13	MET	-	initiating methionine	UNP P22629
D	44	VAL	GLU	engineered mutation	UNP P22629
D	45	THR	SER	engineered mutation	UNP P22629
D	47	ARG	VAL	engineered mutation	UNP P22629

- Molecule 2 is a protein called strep-tag II peptide.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
2	E	6	Total	C	N	O	0	0	0
			52	33	9	10			
2	F	6	Total	C	N	O	0	0	0
			55	36	10	9			
2	G	6	Total	C	N	O	0	0	0
			55	36	10	9			
2	H	6	Total	C	N	O	0	0	0
			55	36	10	9			


- Molecule 3 is water.


Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	73	Total	O	0	0
			73	73		
3	B	50	Total	O	0	0
			50	50		
3	C	54	Total	O	0	0
			54	54		
3	D	52	Total	O	0	0
			52	52		
3	E	2	Total	O	0	0
			2	2		
3	F	3	Total	O	0	0
			3	3		
3	H	4	Total	O	0	0
			4	4		

3 Residue-property plots [i](#)


These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density ($RSRZ > 2$). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

- Molecule 1: streptavidin

Chain A: 





- Molecule 1: streptavidin

Chain B: 





- Molecule 1: streptavidin

Chain C: 





- Molecule 1: streptavidin

Chain D: 



- Molecule 2: strep-tag II peptide


Chain E: 





- Molecule 2: strep-tag II peptide



- Molecule 2: strep-tag II peptide



- Molecule 2: strep-tag II peptide



4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	58.70Å 86.34Å 47.45Å 90.00° 98.92° 90.00°	Depositor
Resolution (Å)	36.00 – 1.70 35.98 – 1.69	Depositor EDS
% Data completeness (in resolution range)	94.8 (36.00-1.70) 94.5 (35.98-1.69)	Depositor EDS
R_{merge}	(Not available)	Depositor
R_{sym}	0.04	Depositor
$\langle I/\sigma(I) \rangle$ ¹	2.65 (at 1.70Å)	Xtrriage
Refinement program	CNS, REFMAC	Depositor
R, R_{free}	0.175 , 0.217 0.190 , 0.231	Depositor DCC
R_{free} test set	2440 reflections (4.98%)	wwPDB-VP
Wilson B-factor (Å ²)	25.4	Xtrriage
Anisotropy	0.061	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.37 , 57.0	EDS
L-test for twinning ²	$\langle L \rangle = 0.50$, $\langle L^2 \rangle = 0.33$	Xtrriage
Estimated twinning fraction	No twinning to report.	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	4054	wwPDB-VP
Average B, all atoms (Å ²)	28.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 6.75% of the height of the origin peak. No significant pseudotranslation is detected.*

¹Intensities estimated from amplitudes.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality i

5.1 Standard geometry i

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.93	0/930	1.65	9/1273 (0.7%)
1	B	0.87	0/921	1.49	8/1260 (0.6%)
1	C	0.88	0/921	1.52	12/1260 (1.0%)
1	D	0.85	0/916	1.40	5/1253 (0.4%)
2	E	0.80	0/54	1.14	0/72
2	F	1.08	1/57 (1.8%)	1.13	0/75
2	G	0.97	0/57	2.21	2/75 (2.7%)
2	H	0.72	0/57	1.21	0/75
All	All	0.88	1/3913 (0.0%)	1.52	36/5343 (0.7%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	F	4	HIS	N-CA	5.35	1.57	1.46

All (36) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	A	128	ASP	CB-CG-OD2	17.08	133.67	118.30
2	G	7	PHE	CB-CG-CD1	-12.02	112.39	120.80
2	G	7	PHE	CB-CG-CD2	11.20	128.64	120.80
1	A	128	ASP	CB-CG-OD1	-10.07	109.24	118.30
1	B	128	ASP	CB-CG-OD2	9.34	126.70	118.30
1	D	128	ASP	CB-CG-OD2	8.87	126.28	118.30
1	A	36	ASP	CB-CG-OD2	8.40	125.86	118.30
1	D	132	LYS	CD-CE-NZ	7.76	129.56	111.70
1	C	84	ARG	NE-CZ-NH2	-7.48	116.56	120.30
1	B	132	LYS	CD-CE-NZ	7.45	128.84	111.70
1	A	60	TYR	CB-CG-CD2	-6.91	116.86	121.00
1	C	67	ASP	CB-CG-OD2	6.74	124.36	118.30
1	B	128	ASP	OD1-CG-OD2	-6.43	111.08	123.30
1	A	53	ARG	NE-CZ-NH1	6.29	123.45	120.30
1	C	59	ARG	NE-CZ-NH2	-6.10	117.25	120.30

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
1	C	66	THR	CA-CB-CG2	-5.77	104.33	112.40
1	B	39	LEU	CB-CG-CD1	-5.75	101.22	111.00
1	B	84	ARG	NE-CZ-NH2	-5.72	117.44	120.30
1	A	132	LYS	CD-CE-NZ	5.68	124.77	111.70
1	D	122	SER	N-CA-CB	-5.67	102.00	110.50
1	C	36	ASP	CB-CG-OD1	5.63	123.37	118.30
1	C	29	PHE	N-CA-C	-5.57	95.97	111.00
1	D	131	THR	OG1-CB-CG2	-5.54	97.26	110.00
1	A	25	LEU	CA-CB-CG	-5.53	102.58	115.30
1	B	36	ASP	CB-CG-OD1	5.50	123.25	118.30
1	D	107	GLN	CA-CB-CG	-5.47	101.36	113.40
1	C	84	ARG	NE-CZ-NH1	5.44	123.02	120.30
1	A	75	TRP	O-C-N	5.40	131.34	122.70
1	C	29	PHE	N-CA-CB	-5.34	100.98	110.60
1	C	62	SER	CB-CA-C	-5.28	100.06	110.10
1	A	124	LEU	CB-CG-CD2	-5.28	102.02	111.00
1	C	108	TRP	CA-CB-CG	5.24	123.65	113.70
1	C	31	VAL	CA-CB-CG1	5.14	118.61	110.90
1	B	96	TYR	CG-CD1-CE1	-5.08	117.23	121.30
1	B	47	ARG	CG-CD-NE	5.02	122.34	111.80
1	C	130	PHE	CB-CG-CD1	-5.00	117.30	120.80

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	907	0	857	11	0
1	B	899	0	850	12	0
1	C	899	0	850	22	1
1	D	894	0	845	19	0
2	E	52	0	41	1	0
2	F	55	0	49	4	0
2	G	55	0	49	1	0
2	H	55	0	49	6	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
3	A	73	0	0	1	0
3	B	50	0	0	2	1
3	C	54	0	0	3	0
3	D	52	0	0	4	0
3	E	2	0	0	0	0
3	F	3	0	0	0	0
3	H	4	0	0	2	0
All	All	4054	0	3590	68	1

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 9.

All (68) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:24:GLN:NE2	3:C:163:HOH:O	2.01	0.94
1:C:15:ALA:O	3:C:166:HOH:O	1.86	0.93
1:D:107:GLN:OE1	3:D:171:HOH:O	1.89	0.89
2:F:9:LYS:H	2:F:9:LYS:CD	1.86	0.89
1:B:45:THR:OG1	1:B:52:SER:HB3	1.75	0.85
1:B:103:ARG:HD2	3:B:174:HOH:O	1.78	0.81
1:C:61:ASP:OD2	1:D:87:HIS:HD2	1.68	0.76
2:F:9:LYS:CD	2:F:9:LYS:N	2.52	0.72
1:C:24:GLN:OE1	3:C:179:HOH:O	2.08	0.71
2:F:9:LYS:H	2:F:9:LYS:HD3	1.55	0.71
1:C:44:VAL:HG22	1:C:53:ARG:HG2	1.72	0.71
2:F:9:LYS:N	2:F:9:LYS:HD2	2.07	0.70
1:A:101:GLU:O	1:A:101:GLU:HG3	1.92	0.69
1:C:103:ARG:HH11	1:C:103:ARG:HG3	1.58	0.66
1:C:44:VAL:CG2	1:C:53:ARG:HG2	2.25	0.66
1:C:15:ALA:HB3	1:C:18:THR:H	1.61	0.66
2:H:9:LYS:HE2	3:H:239:HOH:O	1.97	0.63
1:A:84:ARG:HD3	3:A:173:HOH:O	1.98	0.62
1:D:100:ALA:O	1:D:101:GLU:CB	2.46	0.62
1:D:100:ALA:O	1:D:101:GLU:HB2	2.00	0.61
2:H:9:LYS:CE	3:H:239:HOH:O	2.48	0.60
1:A:120:TRP:CH2	1:A:121:LYS:HD3	2.38	0.58
1:C:110:LEU:C	1:C:110:LEU:HD23	2.24	0.58
1:B:120:TRP:CE3	1:B:121:LYS:HB3	2.40	0.57
1:B:103:ARG:CD	3:B:174:HOH:O	2.47	0.57
1:D:46:ALA:HB3	2:H:8:GLU:O	2.05	0.56

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:73:LEU:HD12	1:C:73:LEU:C	2.26	0.55
1:B:47:ARG:HH11	1:B:47:ARG:HG2	1.72	0.55
1:A:25:LEU:HD21	1:D:120:TRP:CH2	2.42	0.54
1:C:46:ALA:HB3	2:G:8:GLU:O	2.08	0.54
1:B:73:LEU:C	1:B:73:LEU:HD12	2.28	0.53
1:C:49:ASN:OD1	1:C:51:GLU:OE1	2.27	0.53
1:B:120:TRP:CZ3	1:B:121:LYS:HB3	2.45	0.52
1:D:34:GLY:N	3:D:162:HOH:O	2.43	0.52
1:B:120:TRP:CZ2	1:B:121:LYS:HD3	2.45	0.52
1:B:47:ARG:HH11	1:B:47:ARG:CG	2.22	0.51
1:A:25:LEU:HD21	1:D:120:TRP:CZ3	2.45	0.51
1:B:45:THR:OG1	1:B:52:SER:CB	2.54	0.51
1:C:51:GLU:H	1:C:51:GLU:CD	2.15	0.49
1:D:100:ALA:C	1:D:101:GLU:HG2	2.32	0.49
1:D:51:GLU:HA	1:D:83:TYR:CD2	2.48	0.49
1:D:49:ASN:OD1	1:D:50:ALA:N	2.36	0.49
1:C:24:GLN:H	1:C:24:GLN:CD	2.16	0.48
2:H:4:HIS:CG	2:H:5:PRO:HD2	2.49	0.48
1:C:49:ASN:CG	1:C:51:GLU:OE1	2.53	0.47
1:A:47:ARG:HG2	1:A:49:ASN:ND2	2.30	0.47
1:C:103:ARG:HH11	1:C:103:ARG:CG	2.28	0.46
1:D:23:ASN:HB3	1:D:130:PHE:CE2	2.51	0.46
1:B:73:LEU:HD12	1:B:73:LEU:O	2.16	0.46
1:A:21:TRP:CZ2	1:A:132:LYS:HE3	2.51	0.45
1:D:103:ARG:CZ	1:D:105:ASN:HD21	2.29	0.45
1:C:110:LEU:C	1:C:110:LEU:CD2	2.87	0.44
1:D:110:LEU:C	1:D:110:LEU:HD23	2.39	0.43
1:C:44:VAL:HG22	1:C:53:ARG:HA	1.99	0.43
1:A:25:LEU:HD11	1:D:120:TRP:CE2	2.54	0.43
1:C:24:GLN:CD	1:C:24:GLN:N	2.72	0.43
1:D:131:THR:HB	3:D:150:HOH:O	2.20	0.42
1:A:45:THR:OG1	1:A:52:SER:HB2	2.20	0.42
1:D:107:GLN:CG	3:D:171:HOH:O	2.67	0.42
2:E:4:HIS:CG	2:E:5:PRO:HD2	2.55	0.42
1:A:120:TRP:CZ2	1:A:121:LYS:HD3	2.55	0.41
1:D:119:ALA:O	1:D:122:SER:N	2.52	0.41
2:H:4:HIS:HA	2:H:5:PRO:HD3	1.91	0.41
1:C:44:VAL:HG21	1:C:53:ARG:HG2	2.02	0.41
1:B:56:LEU:C	1:B:56:LEU:HD12	2.41	0.41
1:A:120:TRP:CE3	2:H:7:PHE:CD2	3.09	0.40
1:C:91:THR:HB	1:D:91:THR:HB	2.04	0.40

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:47:ARG:HB3	1:C:52:SER:OG	2.22	0.40

All (1) symmetry-related close contacts are listed below. The label for Atom-2 includes the symmetry operator and encoded unit-cell translations to be applied.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:132:LYS:NZ	3:B:178:HOH:O[1_455]	2.05	0.15

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	119/127 (94%)	118 (99%)	1 (1%)	0	100	100
1	B	118/127 (93%)	115 (98%)	3 (2%)	0	100	100
1	C	118/127 (93%)	116 (98%)	2 (2%)	0	100	100
1	D	117/127 (92%)	113 (97%)	3 (3%)	1 (1%)	17	5
2	E	4/9 (44%)	4 (100%)	0	0	100	100
2	F	4/9 (44%)	4 (100%)	0	0	100	100
2	G	4/9 (44%)	4 (100%)	0	0	100	100
2	H	4/9 (44%)	4 (100%)	0	0	100	100
All	All	488/544 (90%)	478 (98%)	9 (2%)	1 (0%)	47	30

All (1) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	D	101	GLU

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	90/93 (97%)	87 (97%)	3 (3%)	38	19
1	B	88/93 (95%)	85 (97%)	3 (3%)	37	18
1	C	88/93 (95%)	84 (96%)	4 (4%)	27	10
1	D	88/93 (95%)	85 (97%)	3 (3%)	37	18
2	E	6/9 (67%)	6 (100%)	0	100	100
2	F	6/9 (67%)	5 (83%)	1 (17%)	2	0
2	G	6/9 (67%)	6 (100%)	0	100	100
2	H	6/9 (67%)	6 (100%)	0	100	100
All	All	378/408 (93%)	364 (96%)	14 (4%)	34	15

All (14) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	101	GLU
1	A	107	GLN
1	A	121	LYS
1	B	24	GLN
1	B	47	ARG
1	B	128	ASP
1	C	24	GLN
1	C	49	ASN
1	C	118	ASN
1	C	128	ASP
1	D	73	LEU
1	D	103	ARG
1	D	121	LYS
2	F	9	LYS

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (9) such sidechains are listed below:

Mol	Chain	Res	Type
1	B	105	ASN
1	C	24	GLN
1	C	118	ASN
1	D	87	HIS
1	D	107	GLN
2	E	6	GLN
2	F	6	GLN
2	G	6	GLN
2	H	6	GLN

5.3.3 RNA [i](#)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no monosaccharides in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	121/127 (95%)	0.64	14 (11%) 4 5	16, 24, 43, 58	0
1	B	120/127 (94%)	0.62	10 (8%) 11 13	15, 24, 46, 53	0
1	C	120/127 (94%)	0.69	17 (14%) 2 3	16, 23, 47, 55	0
1	D	119/127 (93%)	0.76	14 (11%) 4 5	15, 25, 48, 57	0
2	E	6/9 (66%)	1.47	1 (16%) 1 1	25, 30, 40, 49	0
2	F	6/9 (66%)	0.54	0 100 100	23, 26, 38, 52	0
2	G	6/9 (66%)	0.73	1 (16%) 1 1	23, 27, 34, 49	0
2	H	6/9 (66%)	0.55	1 (16%) 1 1	24, 27, 39, 57	0
All	All	504/544 (92%)	0.69	58 (11%) 4 5	15, 24, 47, 58	0

All (58) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	D	120	TRP	9.3
1	C	15	ALA	7.3
2	E	3	SER	6.5
1	D	100	ALA	6.3
1	D	47	ARG	5.5
1	D	66	THR	5.0
1	B	66	THR	4.5
1	A	47	ARG	4.2
1	D	98	GLY	4.1
1	D	101	GLU	4.1
1	A	136	SER	4.0
1	B	15	ALA	4.0
1	B	25	LEU	3.7
1	C	49	ASN	3.5
1	D	99	GLY	3.4
1	A	109	LEU	3.4

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	B	109	LEU	3.4
1	D	117	ALA	3.3
1	A	35	ALA	3.3
1	C	46	ALA	3.3
1	B	47	ARG	3.2
2	G	9	LYS	3.0
1	A	135	PRO	3.0
1	A	49	ASN	2.9
1	B	35	ALA	2.8
1	C	29	PHE	2.7
1	A	120	TRP	2.7
1	A	125	VAL	2.7
1	C	24	GLN	2.7
1	C	134	LYS	2.6
1	D	125	VAL	2.6
1	D	68	GLY	2.6
1	C	47	ARG	2.6
1	A	133	VAL	2.6
1	C	51	GLU	2.5
1	C	116	GLU	2.5
1	B	108	TRP	2.5
1	A	66	THR	2.5
1	A	48	GLY	2.5
1	D	124	LEU	2.4
1	C	83	TYR	2.4
1	C	125	VAL	2.4
1	C	124	LEU	2.4
1	D	109	LEU	2.4
1	B	125	VAL	2.4
1	C	108	TRP	2.4
1	C	109	LEU	2.3
1	C	100	ALA	2.3
1	A	51	GLU	2.2
1	B	100	ALA	2.2
1	B	36	ASP	2.2
1	C	118	ASN	2.2
1	C	101	GLU	2.2
2	H	9	LYS	2.2
1	D	49	ASN	2.1
1	A	84	ARG	2.1
1	D	116	GLU	2.0
1	A	68	GLY	2.0

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no monosaccharides in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.