



Full wwPDB NMR Structure Validation Report ⓘ

May 28, 2020 – 10:32 pm BST

PDB ID : 2KLE
Title : ISIC Refined Solution Structure of the Calcium Binding Domain of the C-terminal Cytosolic Domain of Polycystin-2
Authors : Kalbitzer, H.R.
Deposited on : 2009-07-01

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : 2.11
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment was not calculated.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	158937	12864
Ramachandran outliers	154571	11451
Sidechain outliers	154315	11428

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	123	

2 Ensemble composition and analysis

This entry contains 10 models. Model 3 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:730-A:774 (45)	0.11	3

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 2 clusters and 2 single-model clusters were found.

Cluster number	Models
1	5, 6, 7, 8, 9, 10
2	2, 3
Single-model clusters	1; 4

3 Entry composition


There is only 1 type of molecule in this entry. The entry contains 1136 atoms, of which 543 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Polycystin-2.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	73	1136	356	543	105	131	1	0


There are 6 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	674	GLY	-	EXPRESSION TAG	UNP Q13563
A	675	SER	-	EXPRESSION TAG	UNP Q13563
A	676	THR	-	EXPRESSION TAG	UNP Q13563
A	677	ALA	-	EXPRESSION TAG	UNP Q13563
A	678	ILE	-	EXPRESSION TAG	UNP Q13563
A	679	GLY	-	EXPRESSION TAG	UNP Q13563




4.2.3 Score per residue for model 3 (medoid)

- Molecule 1: Polycystin-2




4.2.4 Score per residue for model 4

- Molecule 1: Polycystin-2



4.2.5 Score per residue for model 5

- Molecule 1: Polycystin-2




4.2.6 Score per residue for model 6

- Molecule 1: Polycystin-2

4.2.10 Score per residue for model 10

● Molecule 1: Polycystin-2



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *restrained molecular dynamics, simulated annealing*.

Of the 1000 calculated structures, 10 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CNS	refinement	1.1
Auremol-ISIC	refinement	

No chemical shift data was provided. No validations of the models with respect to experimental NMR restraints is performed at this time.

COVALENT-GEOMETRY INFOmissingINFO

5.1 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	361	332	332	19±3
All	All	3610	3320	3320	191

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 28.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:737:ASN:HA	1:A:769:GLU:N	0.70	2.02	10	10
1:A:737:ASN:HA	1:A:769:GLU:CA	0.66	2.21	2	10
1:A:737:ASN:HA	1:A:769:GLU:HA	0.63	1.68	1	10
1:A:767:ASP:HB3	1:A:770:LEU:HD13	0.62	1.70	5	5
1:A:748:LYS:N	1:A:748:LYS:HD3	0.59	2.12	2	3
1:A:748:LYS:HD3	1:A:748:LYS:N	0.59	2.13	10	5
1:A:759:PHE:HB2	1:A:763:ASP:HB2	0.57	1.77	2	10
1:A:741:LEU:HD22	1:A:768:GLN:CD	0.56	2.21	10	10
1:A:742:ARG:HA	1:A:745:LEU:HD12	0.56	1.76	5	10

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:737:ASN:CA	1:A:769:GLU:HA	0.55	2.31	2	10
1:A:768:GLN:O	1:A:769:GLU:HG3	0.55	2.01	9	1
1:A:748:LYS:CE	1:A:755:ILE:HD13	0.54	2.33	8	4
1:A:755:ILE:HA	1:A:758:ILE:HG22	0.54	1.79	4	10
1:A:742:ARG:O	1:A:746:LYS:N	0.51	2.44	1	10
1:A:748:LYS:HE2	1:A:755:ILE:HD13	0.50	1.83	8	7
1:A:735:LYS:HB2	1:A:771:THR:HA	0.50	1.83	2	8
1:A:741:LEU:HD21	1:A:745:LEU:HD21	0.50	1.84	8	8
1:A:731:GLN:OE1	1:A:736:LEU:HA	0.49	2.08	3	4
1:A:741:LEU:HD22	1:A:768:GLN:CG	0.48	2.38	8	9
1:A:731:GLN:CD	1:A:741:LEU:HA	0.47	2.29	10	7
1:A:740:GLU:O	1:A:743:GLN:HB2	0.47	2.09	8	7
1:A:737:ASN:HB2	1:A:740:GLU:CB	0.47	2.40	2	2
1:A:731:GLN:OE1	1:A:741:LEU:HA	0.45	2.10	3	2
1:A:748:LYS:HE3	1:A:755:ILE:HG21	0.45	1.88	8	1
1:A:737:ASN:CB	1:A:769:GLU:HA	0.45	2.41	2	3
1:A:756:GLU:HA	1:A:759:PHE:CE2	0.44	2.48	7	7
1:A:750:HIS:HB2	1:A:754:GLU:OE1	0.44	2.12	8	2
1:A:742:ARG:NH1	1:A:742:ARG:HG3	0.44	2.27	5	1
1:A:750:HIS:HB2	1:A:754:GLU:OE2	0.43	2.13	10	1
1:A:736:LEU:C	1:A:737:ASN:HD22	0.43	2.16	2	1
1:A:737:ASN:HB3	1:A:769:GLU:HA	0.42	1.91	10	1
1:A:735:LYS:CB	1:A:771:THR:HA	0.42	2.44	2	4
1:A:741:LEU:HD23	1:A:745:LEU:HD11	0.41	1.90	1	1
1:A:738:PHE:O	1:A:742:ARG:HG2	0.41	2.16	1	1
1:A:737:ASN:O	1:A:741:LEU:N	0.41	2.44	7	5
1:A:742:ARG:HD2	1:A:748:LYS:NZ	0.41	2.30	7	1

5.2 Torsion angles [i](#)

5.2.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	45/123 (37%)	35±1 (78±2%)	7±1 (15±3%)	3±1 (7±2%)	2	16
All	All	450/1230 (37%)	350 (78%)	68 (15%)	32 (7%)	2	16

All 4 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	769	GLU	10
1	A	768	GLN	10
1	A	734	GLY	6
1	A	748	LYS	6

5.2.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	37/108 (34%)	27±1 (74±4%)	10±1 (26±4%)	2 22
All	All	370/1080 (34%)	272 (74%)	98 (26%)	2 22

All 19 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	748	LYS	10
1	A	759	PHE	10
1	A	735	LYS	10
1	A	758	ILE	10
1	A	743	GLN	10
1	A	745	LEU	9
1	A	744	ASP	9
1	A	739	ASP	5
1	A	767	ASP	4
1	A	751	THR	4
1	A	736	LEU	3
1	A	769	GLU	3
1	A	742	ARG	3
1	A	771	THR	2
1	A	737	ASN	2
1	A	752	ASP	1
1	A	730	ARG	1
1	A	764	GLN	1
1	A	768	GLN	1

5.2.3 RNA [i](#)

There are no RNA molecules in this entry.

5.3 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.4 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.5 Ligand geometry [i](#)

There are no ligands in this entry.

5.6 Other polymers [i](#)

There are no such molecules in this entry.

5.7 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Chemical shift validation

No chemical shift data were provided