



Full wwPDB NMR Structure Validation Report ⓘ

Oct 30, 2024 – 06:38 AM EDT

PDB ID : 2KR1
BMRB ID : 16620
Title : Solution NMR structure of zinc binding N-terminal domain of ubiquitin-protein ligase E3A from Homo Sapiens. Northeast Structural Genomics Consortium (NESG) target HR3662
Authors : Lemak, A.; Yee, A.; Fares, C.; Semesi, A.; Xiao, R.; Montelione, G.; Dhe-Paganon, S.; Arrowsmith, C.; Northeast Structural Genomics Consortium (NESG); Structural Genomics Consortium (SGC)
Deposited on : 2009-11-27

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The types of validation reports are described at

<http://www.wwpdb.org/validation/2017/FAQs#types>.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)
wwPDB-RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
wwPDB-ShiftChecker : v1.2
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.39


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 88%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	210492	14027
Ramachandran outliers	207382	12486
Sidechain outliers	206894	12463

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	82	

2 Ensemble composition and analysis i

This entry contains 20 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:4-A:60 (57)	0.67	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 3 clusters and 1 single-model cluster was found.

Cluster number	Models
1	3, 4, 5, 6, 8, 9, 10, 13, 14, 15, 16, 17, 19, 20
2	1, 12, 18
3	7, 11
Single-model clusters	2

3 Entry composition

There are 2 unique types of molecules in this entry. The entry contains 992 atoms, of which 490 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Ubiquitin protein ligase E3A.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	64	991	312	490	90	92	7	0

There are 18 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	-17	MET	-	expression tag	UNP Q9H2G0
A	-16	HIS	-	expression tag	UNP Q9H2G0
A	-15	HIS	-	expression tag	UNP Q9H2G0
A	-14	HIS	-	expression tag	UNP Q9H2G0
A	-13	HIS	-	expression tag	UNP Q9H2G0
A	-12	HIS	-	expression tag	UNP Q9H2G0
A	-11	HIS	-	expression tag	UNP Q9H2G0
A	-10	SER	-	expression tag	UNP Q9H2G0
A	-9	SER	-	expression tag	UNP Q9H2G0
A	-8	GLY	-	expression tag	UNP Q9H2G0
A	-7	ARG	-	expression tag	UNP Q9H2G0
A	-6	GLU	-	expression tag	UNP Q9H2G0
A	-5	ASN	-	expression tag	UNP Q9H2G0
A	-4	LEU	-	expression tag	UNP Q9H2G0
A	-3	TYR	-	expression tag	UNP Q9H2G0
A	-2	PHE	-	expression tag	UNP Q9H2G0
A	-1	GLN	-	expression tag	UNP Q9H2G0
A	0	GLY	-	expression tag	UNP Q9H2G0

- Molecule 2 is ZINC ION (three-letter code: ZN) (formula: Zn).


Mol	Chain	Residues	Atoms	
			Total	Zn
2	A	1	1	1

4 Residue-property plots [i](#)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Ubiquitin protein ligase E3A




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Ubiquitin protein ligase E3A




4.2.2 Score per residue for model 2

- Molecule 1: Ubiquitin protein ligase E3A




4.2.3 Score per residue for model 3

- Molecule 1: Ubiquitin protein ligase E3A




4.2.4 Score per residue for model 4

- Molecule 1: Ubiquitin protein ligase E3A




4.2.5 Score per residue for model 5

- Molecule 1: Ubiquitin protein ligase E3A




4.2.6 Score per residue for model 6

- Molecule 1: Ubiquitin protein ligase E3A




4.2.7 Score per residue for model 7

- Molecule 1: Ubiquitin protein ligase E3A




4.2.8 Score per residue for model 8

- Molecule 1: Ubiquitin protein ligase E3A




4.2.9 Score per residue for model 9

- Molecule 1: Ubiquitin protein ligase E3A




4.2.10 Score per residue for model 10

- Molecule 1: Ubiquitin protein ligase E3A




4.2.11 Score per residue for model 11

- Molecule 1: Ubiquitin protein ligase E3A




4.2.12 Score per residue for model 12

- Molecule 1: Ubiquitin protein ligase E3A




4.2.13 Score per residue for model 13

- Molecule 1: Ubiquitin protein ligase E3A




4.2.14 Score per residue for model 14

- Molecule 1: Ubiquitin protein ligase E3A




4.2.15 Score per residue for model 15

- Molecule 1: Ubiquitin protein ligase E3A




4.2.16 Score per residue for model 16

- Molecule 1: Ubiquitin protein ligase E3A




4.2.17 Score per residue for model 17

- Molecule 1: Ubiquitin protein ligase E3A




4.2.18 Score per residue for model 18

- Molecule 1: Ubiquitin protein ligase E3A




4.2.19 Score per residue for model 19

- Molecule 1: Ubiquitin protein ligase E3A



4.2.20 Score per residue for model 20 (medoid)

- Molecule 1: Ubiquitin protein ligase E3A



5 Refinement protocol and experimental data overview

The models were refined using the following method: *restrained molecular dynamics*.

Of the 100 calculated structures, 20 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
CYANA	structure solution	
CNSSOLVE	refinement	

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	709
Number of shifts mapped to atoms	709
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	88%

6 Model quality i

6.1 Standard geometry i

Bond lengths and bond angles in the following residue types are not validated in this section:
ZN

There are no covalent bond-length or bond-angle outliers.

There are no bond-length outliers.

There are no bond-angle outliers.

There are no chirality outliers.

There are no planarity outliers.

6.2 Too-close contacts i

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	440	429	427	2±1
All	All	8820	8580	8540	35

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 2.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:26:CYS:HB3	1:A:34:CYS:SG	0.59	2.37	19	6
1:A:21:CYS:SG	1:A:23:ASN:OD1	0.55	2.64	1	1
1:A:40:MET:N	1:A:40:MET:SD	0.55	2.80	18	1
1:A:31:CYS:HB3	1:A:34:CYS:SG	0.50	2.47	17	6
1:A:40:MET:SD	1:A:45:ALA:HB2	0.48	2.48	20	2
1:A:8:HIS:O	1:A:12:ARG:HG3	0.47	2.10	5	6
1:A:48:LYS:O	1:A:52:LEU:HG	0.45	2.12	18	3
1:A:8:HIS:O	1:A:12:ARG:HG2	0.45	2.11	13	4
1:A:12:ARG:HB3	1:A:59:LEU:HD11	0.43	1.90	9	1
1:A:12:ARG:HB3	1:A:59:LEU:HD21	0.43	1.89	20	1
1:A:17:LEU:HD21	1:A:52:LEU:HD12	0.42	1.91	16	1
1:A:37:PHE:CZ	1:A:40:MET:SD	0.41	3.14	3	1

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:31:CYS:SG	1:A:32:ALA:O	0.41	2.79	18	1
1:A:21:CYS:SG	1:A:23:ASN:HB3	0.40	2.56	13	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles	
1	A	57/82 (70%)	52±1 (91±2%)	5±1 (9±2%)	0±0 (0±0%)	50	84
All	All	1140/1640 (70%)	1039 (91%)	100 (9%)	1 (0%)	50	84

All 1 unique Ramachandran outliers are listed below.

Mol	Chain	Res	Type	Models (Total)
1	A	58	LYS	1

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	45/68 (66%)	43±1 (95±3%)	2±1 (5±3%)	22	74
All	All	900/1360 (66%)	853 (95%)	47 (5%)	22	74

All 11 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	9	LEU	13
1	A	29	GLU	7
1	A	40	MET	7

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	Models (Total)
1	A	56	ASN	4
1	A	54	LYS	4
1	A	24	GLU	3
1	A	39	ARG	3
1	A	7	LYS	2
1	A	42	ASN	2
1	A	50	LEU	1
1	A	11	GLU	1

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no oligosaccharides in this entry.

6.6 Ligand geometry [i](#)

Of 1 ligands modelled in this entry, 1 is monoatomic - leaving 0 for Mogul analysis.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 88% for the well-defined parts and 83% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	709
Number of shifts mapped to atoms	709
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

The following table shows the suggested chemical shift referencing corrections.

Nucleus	# values	Correction \pm precision, ppm	Suggested action
$^{13}\text{C}_\alpha$	62	-0.29 ± 0.15	None needed (< 0.5 ppm)
$^{13}\text{C}_\beta$	59	0.31 ± 0.14	None needed (< 0.5 ppm)
$^{13}\text{C}'$	54	-0.03 ± 0.09	None needed (< 0.5 ppm)
^{15}N	55	-0.89 ± 0.42	Should be applied

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 88%, i.e. 661 atoms were assigned a chemical shift out of a possible 753. 0 out of 6 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	^1H	^{13}C	^{15}N
Backbone	272/285 (95%)	111/115 (97%)	109/114 (96%)	52/56 (93%)
Sidechain	359/405 (89%)	246/264 (93%)	108/125 (86%)	5/16 (31%)

Continued on next page...

Continued from previous page...

	Total	¹ H	¹³ C	¹⁵ N
Aromatic	30/63 (48%)	20/30 (67%)	10/29 (34%)	0/4 (0%)
Overall	661/753 (88%)	377/409 (92%)	227/268 (85%)	57/76 (75%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 83%, i.e. 709 atoms were assigned a chemical shift out of a possible 858. 0 out of 6 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	290/316 (92%)	119/127 (94%)	116/128 (91%)	55/61 (90%)
Sidechain	389/471 (83%)	266/306 (87%)	118/145 (81%)	5/20 (25%)
Aromatic	30/71 (42%)	20/34 (59%)	10/31 (32%)	0/6 (0%)
Overall	709/858 (83%)	405/467 (87%)	244/304 (80%)	60/87 (69%)

7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition. If well-defined core and ill-defined regions are not identified then it is shown as gray bars.

Random coil index (RCI) for chain A:

