

Full wwPDB X-ray Structure Validation Report (i)

Jun 19, 2024 – 01:08 AM EDT

PDB ID	:	4KR0
Title	:	Complex structure of MERS-CoV spike RBD bound to CD26
Authors	:	Lu, G.; Hu, Y.; Wang, Q.; Qi, J.; Gao, F.; Li, Y.; Zhang, Y.; Zhang, W.;
		Yuan, Y.; Zhang, B.; Shi, Y.; Yan, J.; Gao, G.F.
Deposited on	:	2013-05-15
Resolution	:	2.70 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.37.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $X\text{-}RAY \, DIFFRACTION$

The reported resolution of this entry is 2.70 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	$\begin{array}{c} {\rm Whole \ archive} \\ (\#{\rm Entries}) \end{array}$	${f Similar\ resolution}\ (\#{ m Entries,\ resolution\ range}({ m \AA}))$
R _{free}	130704	2808 (2.70-2.70)
Clashscore	141614	3122 (2.70-2.70)
Ramachandran outliers	138981	3069 (2.70-2.70)
Sidechain outliers	138945	3069 (2.70-2.70)
RSRZ outliers	127900	2737 (2.70-2.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain				
1	А	739	^{2%} 82%			15%	••
2	В	251	^{2%} 63% 1	18%	•	17%	_
3	С	2	100%				
3	D	2	100%				
3	F	2	100%				



Mol	Chain	Length	Quality of chain	
3	G	2	100%	
4	Е	3	67%	33%



2 Entry composition (i)

There are 6 unique types of molecules in this entry. The entry contains 7944 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Dipeptidyl peptidase 4.

Mol	Chain	Residues		A	toms			ZeroOcc	AltConf	Trace
1	А	728	Total 5963	C 3827	N 982	O 1128	S 26	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
А	34	ALA	-	expression tag	UNP P27487
А	35	ASP	-	expression tag	UNP P27487
А	36	GLY	-	expression tag	UNP P27487
А	37	ILE	-	expression tag	UNP P27487
А	38	GLN	-	expression tag	UNP P27487
А	767	HIS	-	expression tag	UNP P27487
А	768	HIS	-	expression tag	UNP P27487
А	769	HIS	-	expression tag	UNP P27487
А	770	HIS	-	expression tag	UNP P27487
А	771	HIS	-	expression tag	UNP P27487
А	772	HIS	-	expression tag	UNP P27487

• Molecule 2 is a protein called S protein.

Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace	
2	В	208	Total 1608	C 1026	N 256	0 315	S 11	0	0	0

There are 11 discrepancies between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
В	362	ALA	-	expression tag	UNP K0BRG7
В	363	ASP	-	expression tag	UNP K0BRG7
В	364	GLY	-	expression tag	UNP K0BRG7
В	365	ILE	-	expression tag	UNP K0BRG7
В	366	GLN	-	expression tag	UNP K0BRG7



00100000	ica ji cint pi c	ere ae page			
Chain	Residue	Modelled	Actual	Comment	Reference
В	607	HIS	-	expression tag	UNP K0BRG7
В	608	HIS	-	expression tag	UNP K0BRG7
В	609	HIS	-	expression tag	UNP K0BRG7
В	610	HIS	-	expression tag	UNP K0BRG7
В	611	HIS	-	expression tag	UNP K0BRG7
В	612	HIS	-	expression tag	UNP K0BRG7

• Molecule 3 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
3	С	2	Total C N O 28 16 2 10	0	0	0
3	D	2	Total C N O 28 16 2 10	0	0	0
3	F	2	Total C N O 28 16 2 10	0	0	0
3	G	2	Total C N O 28 16 2 10	0	0	0

• Molecule 4 is an oligosaccharide called beta-D-mannopyranose-(1-4)-2-acetamido-2-deoxy-b eta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose.



Mol	Chain	Residues	Atoms				ZeroOcc	AltConf	Trace
4	Е	3	Total 39	C 22	N 2	O 15	0	0	0

• Molecule 5 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$).





Mol	Chain	Residues	Atoms				ZeroOcc	AltConf
5	А	1	Total 14	C 8	N 1	O 5	0	0
5	А	1	Total 14	C 8	N 1	O 5	0	0

• Molecule 6 is water.

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
6	А	177	Total O 177 177	0	0
6	В	17	Total O 17 17	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.



• Molecule 1: Dipeptidyl peptidase 4

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C:

100%



NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain	D:
-------	----

100%

100%

NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain F:

NAG1 NAG2

• Molecule 3: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain G:	100%	
NAG2 NAG2		
• Molecule 4: beta-D-m	nannopyranose-(1-4)-2-acetamido-2-dec	oxy-beta-D-glu

α · α		
Chain E:	67%	33%

NAG1 NAG2 BMA3



4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 61 2 2	Depositor
Cell constants	110.22Å 110.22Å 527.60Å	Depositor
a, b, c, α , β , γ	90.00° 90.00° 120.00°	Depositor
Bosolution(A)	48.85 - 2.70	Depositor
Resolution (A)	48.85 - 2.70	EDS
% Data completeness	99.9 (48.85-2.70)	Depositor
(in resolution range)	99.9 (48.85 - 2.70)	EDS
R_{merge}	0.12	Depositor
R_{sym}	0.12	Depositor
$< I/\sigma(I) > 1$	4.31 (at 2.69 Å)	Xtriage
Refinement program	PHENIX (phenix.refine: 1.8_1069)	Depositor
P. P.	0.198 , 0.232	Depositor
n, n_{free}	0.200 , 0.234	DCC
R_{free} test set	2712 reflections (5.07%)	wwPDB-VP
Wilson B-factor $(Å^2)$	48.3	Xtriage
Anisotropy	0.472	Xtriage
Bulk solvent $k_{sol}(e/Å^3), B_{sol}(Å^2)$	0.33 , 47.0	EDS
L-test for $twinning^2$	$< L >=0.50, < L^2>=0.33$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.93	EDS
Total number of atoms	7944	wwPDB-VP
Average B, all atoms $(Å^2)$	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 2.46% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: NAG, BMA

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond	angles
	Unam	RMSZ	# Z > 5	RMSZ	# Z > 5
1	А	0.34	0/6135	0.50	0/8344
2	В	0.33	0/1647	0.50	0/2249
All	All	0.34	0/7782	0.50	0/10593

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	А	5963	0	5678	60	0
2	В	1608	0	1574	23	0
3	С	28	0	25	0	0
3	D	28	0	25	0	0
3	F	28	0	25	0	0
3	G	28	0	25	0	0
4	Е	39	0	34	1	0
5	А	28	0	26	2	0
6	А	177	0	0	12	0
6	В	17	0	0	0	0
All	All	7944	0	7412	84	0



The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 6.

All (84) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom_1	Atom_2	Interatomic	Clash
Atom-1	Atom-2	distance (Å)	overlap (Å)
1:A:211:TYR:OH	6:A:1016:HOH:O	1.94	0.82
1:A:588:ASP:OD2	6:A:1023:HOH:O	2.09	0.69
2:B:408:ASN:HB3	2:B:587:LYS:HB3	1.74	0.68
1:A:400:GLY:O	6:A:923:HOH:O	2.11	0.68
1:A:696:LYS:HG3	1:A:728:VAL:HG22	1.76	0.67
1:A:640:LEU:HD11	1:A:650:GLY:HA3	1.76	0.67
5:A:813:NAG:O6	6:A:953:HOH:O	2.13	0.66
2:B:493:LYS:NZ	2:B:565:GLU:O	2.28	0.66
1:A:77:LEU:HD23	1:A:88:VAL:HA	1.79	0.65
2:B:501:ASN:ND2	2:B:559:SER:OG	2.31	0.63
1:A:242:SER:OG	1:A:243:ASP:N	2.32	0.62
1:A:435:GLN:NE2	6:A:936:HOH:O	2.29	0.60
1:A:70:TYR:HB3	1:A:79:PHE:CE2	2.38	0.59
1:A:329:ASP:OD1	1:A:343:ARG:NH1	2.36	0.58
2:B:457:SER:HB3	2:B:460:SER:HB3	1.85	0.58
1:A:173:TYR:CE2	1:A:184:ARG:HG3	2.39	0.58
1:A:658:ARG:NH1	6:A:1003:HOH:O	2.37	0.57
2:B:399:PHE:O	2:B:523:TYR:OH	2.15	0.54
1:A:184:ARG:HD3	1:A:186:THR:O	2.07	0.54
1:A:184:ARG:HD2	1:A:187:TRP:CE2	2.42	0.54
1:A:358:ARG:NH1	6:A:1071:HOH:O	2.40	0.54
1:A:135:TYR:HD1	1:A:142:LEU:HD12	1.72	0.54
1:A:648:LYS:NZ	1:A:762:CYS:O	2.31	0.53
1:A:630:SER:OG	1:A:740:HIS:NE2	2.36	0.53
1:A:70:TYR:HB3	1:A:79:PHE:HE2	1.72	0.53
1:A:201:TRP:CZ2	1:A:205:GLU:HG3	2.44	0.52
2:B:467:PHE:O	2:B:524:SER:HB2	2.10	0.52
1:A:105:TYR:OH	6:A:987:HOH:O	2.18	0.52
1:A:614:SER:HA	1:A:619:VAL:HB	1.91	0.52
2:B:480:ILE:HB	2:B:571:PHE:HB2	1.91	0.51
2:B:383:CYS:N	2:B:408:ASN:O	2.44	0.51
1:A:729:ASP:OD2	6:A:941:HOH:O	2.19	0.50
1:A:49:LEU:HD22	1:A:749:GLN:HA	1.94	0.49
1:A:734:TRP:O	6:A:959:HOH:O	2.20	0.49
1:A:89:PHE:CE1	1:A:107:ILE:HD13	2.48	0.49
1:A:115:LEU:HD11	1:A:132:TYR:HB3	1.96	0.48
2:B:581:THR:O	2:B:583:SER:N	2.47	0.47



	io as page	Interatomic	Clash	
Atom-1	Atom-2	distance (Å)	overlap (Å)	
2:B:519:ASN:HB2	2:B:522:GLN:OE1	2.13	0.47	
2:B:485:PRO:O	2:B:566:GLN:HG2	2.15	0.46	
1:A:123:GLN:HB3	1:A:127:SER:OG	2.15	0.46	
1:A:446:SER:HB2	1:A:457:TYR:CE1	2.50	0.46	
1:A:600:THR:OG1	1:A:601:PHE:N	2.48	0.46	
1:A:745:SER:O	1:A:749:GLN:HG3	2.15	0.46	
2:B:408:ASN:HA	2:B:585:CYS:O	2.15	0.46	
1:A:201:TRP:CZ2	1:A:710:ASN:HA	2.51	0.45	
1:A:388:GLN:HB2	1:A:391:LYS:HB2	1.98	0.45	
1:A:302:ASP:OD1	6:A:926:HOH:O	2.21	0.45	
2:B:498:SER:HB3	2:B:534:VAL:HG23	1.99	0.45	
1:A:146:GLU:O	1:A:175:LYS:NZ	2.49	0.45	
1:A:435:GLN:HB3	1:A:441:LYS:HB2	1.97	0.45	
1:A:103:ASN:OD1	1:A:120:TYR:HB3	2.17	0.44	
1:A:422:TYR:CE2	1:A:423:LYS:HD2	2.51	0.44	
2:B:436:ASN:ND2	2:B:577:TYR:OH	2.38	0.44	
2:B:394:PRO:HG3	2:B:400:LYS:HG3	1.99	0.44	
1:A:75:ASN:OD1	1:A:92:ASN:N	2.50	0.44	
1:A:535:ASP:OD1	1:A:537:SER:OG	2.29	0.44	
1:A:630:SER:HB3	1:A:631:TYR:H	1.54	0.43	
2:B:456:LEU:HD12	2:B:456:LEU:HA	1.88	0.43	
1:A:302:ASP:HB3	1:A:314:GLN:HB2	2.00	0.43	
1:A:53:TYR:HB3	1:A:500:LEU:HD11	2.01	0.43	
2:B:587:LYS:HE3	2:B:587:LYS:HB2	1.74	0.43	
1:A:461:PHE:CD2	1:A:468:TYR:HB3	2.54	0.43	
1:A:158:SER:HB3	1:A:163:LYS:HB2	2.00	0.43	
1:A:561:LEU:HD12	1:A:561:LEU:HA	1.82	0.42	
1:A:482:LEU:HD13	1:A:491:LEU:HD12	2.01	0.42	
1:A:544:LEU:HD12	1:A:544:LEU:HA	1.86	0.42	
1:A:294:LEU:O	2:B:542:ARG:HD3	2.19	0.41	
1:A:489:LYS:HB3	1:A:489:LYS:HE3	1.79	0.41	
2:B:484:VAL:HA	2:B:485:PRO:HD3	1.72	0.41	
2:B:479:LEU:HD12	2:B:479:LEU:HA	1.91	0.41	
1:A:397:ILE:HD12	1:A:434:ILE:HD13	2.02	0.41	
1:A:177:GLU:HA	1:A:178:PRO:HD3	1.82	0.41	
1:A:739:ASP:HB2	6:A:939:HOH:O	2.21	0.41	
2:B:504:SER:HB3	2:B:515:PRO:HA	2.03	0.41	
1:A:402:TRP:CD2	1:A:421:GLU:HB2	2.55	0.41	
1:A:472:CYS:O	1:A:478:PRO:HA	2.21	0.41	
1:A:517:ILE:HD12	1:A:612:GLN:HG3	2.03	0.41	
2:B:487:ASN:OD1	2:B:487:ASN:N	2.54	0.41	



Atom-1	Atom-2	Interatomic	Clash
	1100111 2	distance $(Å)$	overlap (Å)
1:A:306:ALA:HB3	1:A:310:ARG:HG2	2.03	0.40
1:A:136:ASP:HB3	1:A:139:LYS:HB2	2.04	0.40
2:B:484:VAL:O	2:B:566:GLN:HB3	2.21	0.40
1:A:74:ASN:ND2	5:A:813:NAG:O7	2.49	0.40
1:A:405:ILE:HG12	1:A:419:SER:HA	2.03	0.40
4:E:2:NAG:C4	4:E:3:BMA:O2	2.67	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	А	726/739~(98%)	690~(95%)	34~(5%)	2~(0%)	41 66
2	В	206/251~(82%)	188 (91%)	15 (7%)	3~(2%)	10 26
All	All	932/990~(94%)	878 (94%)	49 (5%)	5(0%)	29 54

All (5) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	А	520	ASN
2	В	485	PRO
2	В	582	ASN
2	В	382	GLU
1	А	82	GLU

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent side chain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	А	653/662~(99%)	620~(95%)	33~(5%)	24 50		
2	В	190/224~(85%)	172 (90%)	18 (10%)	8 20		
All	All	843/886~(95%)	792 (94%)	51 (6%)	19 42		

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

All (51) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	А	41	LYS
1	А	78	VAL
1	А	86	SER
1	А	120	TYR
1	А	142	LEU
1	А	184	ARG
1	А	202	VAL
1	А	265	THR
1	А	277	SER
1	А	313	LEU
1	А	326	ASP
1	А	351	THR
1	А	358	ARG
1	А	385	CYS
1	А	412	SER
1	А	415	LEU
1	А	436	LEU
1	А	442	VAL
1	А	448	GLU
1	А	482	LEU
1	А	489	LYS
1	А	507	VAL
1	А	543	LEU
1	А	546	VAL
1	A	561	LEU
1	А	566	TYR
1	A	630	SER
1	A	660	GLU
1	А	673	LEU
1	A	684	ARG
1	А	685	ASN
1	А	726	VAL
1	А	736	THR



Mol	Chain	Res	Type
2	В	411	LEU
2	В	423	PHE
2	В	450	LEU
2	В	458	VAL
2	В	465	SER
2	В	473	PHE
2	В	479	LEU
2	В	481	LEU
2	В	484	VAL
2	В	487	ASN
2	В	488	LEU
2	В	490	THR
2	В	510	ASP
2	В	535	TRP
2	В	555	VAL
2	В	565	GLU
2	В	573	ILE
2	В	588	LEU

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (1) such sidechains are listed below:

Mol	Chain	Res	Type
1	А	748	HIS

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

11 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond



Mal	Trune	Chain	Dec	T in le	Bo	ond leng	ths	В	ond ang	les
IVIOI	туре	Chain	nes		Counts	RMSZ	# Z >2	Counts	RMSZ	# Z > 2
3	NAG	С	1	1,3	14,14,15	0.55	0	17,19,21	0.96	1 (5%)
3	NAG	С	2	3	14,14,15	0.54	0	17,19,21	0.97	1 (5%)
3	NAG	D	1	1,3	14,14,15	0.55	0	17,19,21	1.08	1 (5%)
3	NAG	D	2	3	14,14,15	0.53	0	17,19,21	0.95	1 (5%)
4	NAG	Е	1	4,1	14,14,15	0.62	0	17,19,21	1.00	2 (11%)
4	NAG	Е	2	4	14,14,15	0.63	0	17,19,21	0.99	1 (5%)
4	BMA	Е	3	4	11,11,12	0.25	0	$15,\!15,\!17$	0.53	0
3	NAG	F	1	1,3	14,14,15	0.57	0	17,19,21	1.03	1 (5%)
3	NAG	F	2	3	14,14,15	0.49	0	17,19,21	1.22	2 (11%)
3	NAG	G	1	1,3	14,14,15	0.45	0	17,19,21	1.09	1 (5%)
3	NAG	G	2	3	14,14,15	0.53	0	17,19,21	1.14	2 (11%)

length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
3	NAG	С	1	1,3	-	1/6/23/26	0/1/1/1
3	NAG	С	2	3	-	4/6/23/26	0/1/1/1
3	NAG	D	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	D	2	3	-	0/6/23/26	0/1/1/1
4	NAG	Е	1	4,1	-	0/6/23/26	0/1/1/1
4	NAG	Е	2	4	-	2/6/23/26	0/1/1/1
4	BMA	Е	3	4	-	2/2/19/22	0/1/1/1
3	NAG	F	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	F	2	3	-	4/6/23/26	0/1/1/1
3	NAG	G	1	1,3	-	0/6/23/26	0/1/1/1
3	NAG	G	2	3	_	3/6/23/26	0/1/1/1

There are no bond length outliers.

All (13) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms		$Observed(^{o})$	$Ideal(^{o})$
3	F	2	NAG	C4-C3-C2	-2.95	106.69	111.02



Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
3	D	1	NAG	C4-C3-C2	2.68	114.94	111.02
3	G	2	NAG	C1-O5-C5	2.61	115.68	112.19
3	F	1	NAG	C1-O5-C5	2.47	115.50	112.19
3	D	2	NAG	C1-O5-C5	2.42	115.43	112.19
3	С	1	NAG	C3-C4-C5	2.25	114.31	110.23
4	Ε	2	NAG	C2-N2-C7	-2.13	120.05	122.90
3	G	2	NAG	C2-N2-C7	2.11	125.72	122.90
3	С	2	NAG	C1-O5-C5	-2.07	109.41	112.19
3	G	1	NAG	C1-O5-C5	2.04	114.92	112.19
4	Ε	1	NAG	O4-C4-C3	-2.03	105.59	110.38
3	F	2	NAG	C1-C2-N2	2.02	113.61	110.43
4	Е	1	NAG	C2-N2-C7	-2.02	120.20	122.90

There are no chirality outliers.

Mol	Chain	Res	Type	Atoms
3	С	2	NAG	C8-C7-N2-C2
3	С	2	NAG	O7-C7-N2-C2
3	G	2	NAG	C1-C2-N2-C7
4	Е	2	NAG	C8-C7-N2-C2
3	F	2	NAG	O5-C5-C6-O6
3	G	2	NAG	C8-C7-N2-C2
3	G	2	NAG	O7-C7-N2-C2
4	Е	2	NAG	O7-C7-N2-C2
3	С	2	NAG	C4-C5-C6-O6
4	Е	3	BMA	C4-C5-C6-O6
3	С	2	NAG	O5-C5-C6-O6
4	Е	3	BMA	O5-C5-C6-O6
3	С	1	NAG	O5-C5-C6-O6
3	F	2	NAG	C4-C5-C6-O6
3	F	2	NAG	C1-C2-N2-C7
3	F	2	NAG	C3-C2-N2-C7

All (16) torsion outliers are listed below:

There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
4	Е	3	BMA	1	0
4	Е	2	NAG	1	0





The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.

















5.6 Ligand geometry (i)

2 ligands are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol T	Turne	rpo Chain		Tinle	Bo	Bond lengths			Bond angles		
INIOI	туре	Chain	nes	LINK	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2	
5	NAG	А	803	1	14,14,15	0.50	0	17,19,21	0.82	1 (5%)	
5	NAG	А	813	1	14,14,15	0.49	0	17,19,21	0.97	1 (5%)	

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.



Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
5	NAG	А	803	1	-	0/6/23/26	0/1/1/1
5	NAG	А	813	1	-	1/6/23/26	0/1/1/1

There are no bond length outliers.

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	$Observed(^{o})$	$Ideal(^{o})$
5	А	813	NAG	C1-O5-C5	3.13	116.38	112.19
5	А	803	NAG	C1-O5-C5	2.13	115.03	112.19

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
5	А	813	NAG	O5-C5-C6-O6

There are no ring outliers.

1 monomer is involved in 2 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	А	813	NAG	2	0

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	#RSRZ>2	$OWAB(Å^2)$	Q<0.9
1	А	728/739~(98%)	-0.24	12 (1%) 72 74	26, 44, 82, 119	0
2	В	208/251~(82%)	0.06	6 (2%) 51 52	38, 67, 107, 130	0
All	All	936/990~(94%)	-0.17	18 (1%) 66 69	26, 48, 92, 130	0

All (18) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	А	39	SER	5.1
2	В	588	LEU	4.5
2	В	580	ASP	3.6
1	А	128	TYR	3.5
1	А	766	PRO	3.3
1	А	127	SER	3.0
2	В	486	HIS	3.0
1	А	506	ASN	2.9
2	В	587	LYS	2.8
1	А	96	ASP	2.5
1	А	123	GLN	2.5
1	А	99	GLY	2.5
2	В	391	GLY	2.2
1	А	124	TRP	2.2
1	А	95	PHE	2.2
1	А	81	ALA	2.1
2	В	381	VAL	2.1
1	А	83	TYR	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.



6.3 Carbohydrates (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
4	BMA	Е	3	11/12	0.56	0.20	85,103,108,113	0
3	NAG	С	1	14/15	0.78	0.17	89,106,113,117	0
3	NAG	F	2	14/15	0.84	0.21	40,63,72,82	0
3	NAG	С	2	14/15	0.84	0.33	108,117,120,122	0
3	NAG	G	2	14/15	0.85	0.25	93,106,110,111	0
3	NAG	G	1	14/15	0.85	0.19	74,79,90,98	0
4	NAG	Е	2	14/15	0.88	0.17	47,60,68,69	0
3	NAG	D	2	14/15	0.92	0.18	70,76,83,88	0
3	NAG	D	1	14/15	0.93	0.17	52,65,76,83	0
3	NAG	F	1	14/15	0.94	0.12	42,50,57,62	0
4	NAG	E	1	14/15	0.98	0.10	30,42,48,50	0

The following is a graphical depiction of the model fit to experimental electron density for oligosaccharide. Each fit is shown from different orientation to approximate a three-dimensional view.

















6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$B-factors(Å^2)$	Q<0.9
5	NAG	A	813	14/15	0.66	0.29	89,96,101,102	0
5	NAG	А	803	14/15	0.76	0.24	70,85,91,94	0

6.5 Other polymers (i)

There are no such residues in this entry.

