



Full wwPDB X-ray Structure Validation Report ⓘ

May 12, 2020 – 11:33 pm BST

PDB ID : 4KRE
Title : Structure of Human Argonaute-1 bound to endogenous Sf9 RNA
Authors : Faehnle, C.R.; Elkayam, E.; Joshua-Tor, L.
Deposited on : 2013-05-16
Resolution : 1.75 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/XrayValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

MolProbity : 4.02b-467
Xtriage (Phenix) : 1.13
EDS : 2.11
Percentile statistics : 20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac : 5.8.0158
CCP4 : 7.0.044 (Gargrove)
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : 2.11


1 Overall quality at a glance i

The following experimental techniques were used to determine the structure:

X-RAY DIFFRACTION

The reported resolution of this entry is 1.75 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	2340 (1.76-1.76)
Clashscore	141614	2466 (1.76-1.76)
Ramachandran outliers	138981	2437 (1.76-1.76)
Sidechain outliers	138945	2437 (1.76-1.76)
RSRZ outliers	127900	2298 (1.76-1.76)
RNA backbone	3102	1079 (2.40-1.12)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments on the lower bar indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$. The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	858	
2	R	20	

2 Entry composition [i](#)

There are 3 unique types of molecules in this entry. The entry contains 7179 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called Protein argonaute-1.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	S			
1	A	799	6475	4154	1144	1138	39	0	9	0

There is a discrepancy between the modelled and reference sequences:

Chain	Residue	Modelled	Actual	Comment	Reference
A	0	ALA	-	EXPRESSION TAG	UNP Q9UL18

- Molecule 2 is a RNA chain called RNA (5'-R(P*AP*AP*UP*AP*UP*UP*AP*AP*A*AP*AP*AP*AP*AP*AP*AP*AP*AP*AP*A)-3').

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	P			
2	R	11	234	107	46	71	10	0	0	0

- Molecule 3 is water.

Mol	Chain	Residues	Atoms		ZeroOcc	AltConf
3	A	458	Total	O	0	0
			458	458		
3	R	12	Total	O	0	0
			12	12		

4 Data and refinement statistics

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants a, b, c, α , β , γ	70.47Å 98.63Å 73.21Å 90.00° 110.96° 90.00°	Depositor
Resolution (Å)	32.93 – 1.75 49.31 – 1.75	Depositor EDS
% Data completeness (in resolution range)	98.1 (32.93-1.75) 98.1 (49.31-1.75)	Depositor EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$\langle I/\sigma(I) \rangle$ ¹	1.84 (at 1.75Å)	Xtrriage
Refinement program	PHENIX 1.8.1_1168	Depositor
R, R_{free}	0.171 , 0.202 0.171 , 0.203	Depositor DCC
R_{free} test set	4581 reflections (5.00%)	wwPDB-VP
Wilson B-factor (Å ²)	23.2	Xtrriage
Anisotropy	0.283	Xtrriage
Bulk solvent k_{sol} (e/Å ³), B_{sol} (Å ²)	0.39 , 54.4	EDS
L-test for twinning ²	$\langle L \rangle = 0.49$, $\langle L^2 \rangle = 0.32$	Xtrriage
Estimated twinning fraction	0.022 for l,-k,h	Xtrriage
F_o, F_c correlation	0.96	EDS
Total number of atoms	7179	wwPDB-VP
Average B, all atoms (Å ²)	30.0	wwPDB-VP

Xtrriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 4.55% of the height of the origin peak. No significant pseudotranslation is detected.*

¹ Intensities estimated from amplitudes.

² Theoretical values of $\langle |L| \rangle$, $\langle L^2 \rangle$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality [i](#)

5.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	# Z >5	RMSZ	# Z >5
1	A	0.43	0/6651	0.58	0/9005
2	R	0.82	1/262 (0.4%)	1.19	2/402 (0.5%)
All	All	0.45	1/6913 (0.0%)	0.62	2/9407 (0.0%)

All (1) bond length outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(Å)	Ideal(Å)
2	R	1	A	OP3-P	-8.84	1.50	1.61

All (2) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)
2	R	2	A	N1-C6-N6	6.61	122.57	118.60
2	R	2	A	C2-N3-C4	-5.56	107.82	110.60

There are no chirality outliers.

There are no planarity outliers.

5.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	6475	0	6584	15	0
2	R	234	0	121	1	0
3	A	458	0	0	0	0
3	R	12	0	0	0	0
All	All	7179	0	6705	15	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 1.

All (15) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:759:ARG:NH2	2:R:8:A:OP2	2.20	0.73
1:A:250:ASP:OD1	1:A:253:ARG:NH2	2.36	0.58
1:A:115:VAL:HG22	1:A:125:ILE:HG12	1.89	0.55
1:A:519:ILE:HD12	1:A:550:VAL:HG21	1.90	0.52
1:A:254:VAL:HG12	1:A:258:LYS:HE2	1.93	0.51
1:A:467:ASN:O	1:A:471:GLN:HG2	2.10	0.50
1:A:392:ILE:HG23	1:A:397:ILE:HB	1.93	0.50
1:A:705[B]:VAL:HG12	1:A:763:TYR:CD2	2.48	0.49
1:A:489:PHE:CD1	1:A:506:LEU:HD21	2.49	0.47
1:A:208[A]:MET:HG3	1:A:209:TRP:CD1	2.52	0.44
1:A:190:LEU:O	1:A:358:GLN:HG2	2.19	0.43
1:A:63:LYS:HE3	1:A:63:LYS:HB2	1.84	0.41
1:A:451:ILE:HD11	1:A:469:THR:HG21	2.03	0.41
1:A:522:GLY:O	1:A:524:THR:HG22	2.20	0.40
1:A:349:ARG:HB2	1:A:349:ARG:HE	1.71	0.40

There are no symmetry-related clashes.

5.3 Torsion angles [i](#)

5.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	790/858 (92%)	777 (98%)	13 (2%)	0	100 100

There are no Ramachandran outliers to report.

5.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	714/746 (96%)	710 (99%)	4 (1%)	86 79

All (4) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	629[A]	VAL
1	A	629[B]	VAL
1	A	764	TYR
1	A	802	TYR

Some sidechains can be flipped to improve hydrogen bonding and reduce clashes. There are no such sidechains identified.

5.3.3 RNA [i](#)

Mol	Chain	Analysed	Backbone Outliers	Pucker Outliers
2	R	9/20 (45%)	0	0

There are no RNA backbone outliers to report.

There are no RNA pucker outliers to report.

5.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

5.6 Ligand geometry [i](#)

There are no ligands in this entry.

5.7 Other polymers [i](#)

There are no such residues in this entry.

5.8 Polymer linkage issues [i](#)

There are no chain breaks in this entry.

6 Fit of model and data

6.1 Protein, DNA and RNA chains

In the following table, the column labelled ‘#RSRZ> 2’ contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled ‘Q< 0.9’ lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2	OWAB(Å ²)	Q<0.9
1	A	799/858 (93%)	0.11	35 (4%) 34 40	11, 27, 57, 75	0
2	R	11/20 (55%)	0.58	1 (9%) 9 11	23, 30, 79, 90	0
All	All	810/878 (92%)	0.11	36 (4%) 34 40	11, 27, 58, 90	0

All (36) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	239	ARG	5.3
1	A	251	SER	4.9
1	A	484	GLN	4.5
1	A	254	VAL	4.3
1	A	113	PHE	3.9
1	A	357	ASN	3.7
1	A	237	ASP	3.6
1	A	244	GLN	3.5
1	A	238	ILE	3.5
1	A	835	ARG	3.4
1	A	248	LEU	3.3
1	A	353	LYS	3.2
1	A	250	ASP	3.2
1	A	364	LYS	3.2
1	A	605	LYS	3.0
1	A	186	TYR	3.0
1	A	356	ASP	2.9
1	A	458	LYS	2.9
1	A	351	ILE	2.9
2	R	9	A	2.8
1	A	187	TYR	2.5
1	A	633	ARG	2.5
1	A	111	VAL	2.5
1	A	631	ARG	2.5

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type	RSRZ
1	A	236	LEU	2.5
1	A	680[A]	TYR	2.4
1	A	256	PHE	2.4
1	A	354	LEU	2.4
1	A	352	LYS	2.4
1	A	114	GLU	2.3
1	A	84	ILE	2.2
1	A	284	ARG	2.2
1	A	671	GLU	2.2
1	A	255	ARG	2.1
1	A	447	LYS	2.1
1	A	329	GLY	2.1

6.2 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.4 Ligands [i](#)

There are no ligands in this entry.

6.5 Other polymers [i](#)

There are no such residues in this entry.