

wwPDB X-ray Structure Validation Summary Report (i)

Dec 11, 2024 – 04:09 PM JST

PDB ID : 9KU6

Title : Crystal structure of the complex of lactoperoxidase with nitric oxide at 1.72

A resolution

Authors: Maurya, A.; Sharma, P.; Sharma, S.; Singh, T.P.

Deposited on : 2024-12-03

Resolution : 1.72 Å(reported)

This is a wwPDB X-ray Structure Validation Summary Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.orgA user guide is available at

https://www.wwpdb.org/validation/2017/XrayValidationReportHelp with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity: 4.02b-467

Mogul : 1.8.5 (274361), CSD as541be (2020)

Xtriage (Phenix) : 1.21

EDS : 3.0

buster-report : 1.1.7 (2018)

Percentile statistics : 20231227.v01 (using entries in the PDB archive December 27th 2023)

CCP4 : 9.0.004 (Gargrove)

Density-Fitness : 1.0.11

Ideal geometry (proteins) : Engh & Huber (2001) Ideal geometry (DNA, RNA) : Parkinson et al. (1996)


Validation Pipeline (wwPDB-VP) : 2.40

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: X-RAY DIFFRACTION

The reported resolution of this entry is 1.72 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive $(\# \mathrm{Entries})$	$\begin{array}{c} {\rm Similar\ resolution} \\ (\#{\rm Entries},{\rm resolution\ range}({\rm \AA})) \end{array}$
R_{free}	164625	7106 (1.74-1.70)
Clashscore	180529	7746 (1.74-1.70)
Ramachandran outliers	177936	7654 (1.74-1.70)
Sidechain outliers	177891	7654 (1.74-1.70)
RSRZ outliers	164620	7104 (1.74-1.70)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain			
1	A	595	85%	14%	•	
2	С	2	100%			

The following table lists non-polymeric compounds, carbohydrate monomers and non-standard residues in protein, DNA, RNA chains that are outliers for geometric or electron-density-fit criteria:



Mol	Type	Chain	Res	Chirality	Geometry	Clashes	Electron density
4	IOD	A	608	-	-	X	-
4	IOD	A	641	-	-	X	-
4	IOD	A	642	-	-	X	-
4	IOD	A	644	-	-	X	-
4	IOD	A	648	-	-	X	-
4	IOD	A	650	_	_	X	-
4	IOD	A	651	-	-	X	-
5	SCN	A	622	-	-	X	-
5	SCN	A	628	-	-	X	-
5	SCN	A	629	-	-	X	-
7	NO2	A	634	_	_	X	-
7	NO2	A	637	-	-	X	-
8	NO	A	635	-	-	X	-



2 Entry composition (i)

There are 11 unique types of molecules in this entry. The entry contains 5522 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

• Molecule 1 is a protein called Lactoperoxidase.


Mol	Chain	Residues		At	oms			ZeroOcc	AltConf	Trace
1	A	595	Total 4792	C 3050	N 849	O 866	S 27	0	3	0

• Molecule 2 is an oligosaccharide called 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-a cetamido-2-deoxy-beta-D-glucopyranose.



\mathbf{Mol}	Chain	Residues	Atoms	ZeroOcc	AltConf	Trace
2	С	2	Total C N O 28 16 2 10	0	0	0

• Molecule 3 is 2-acetamido-2-deoxy-beta-D-glucopyranose (three-letter code: NAG) (formula: $C_8H_{15}NO_6$) (labeled as "Ligand of Interest" by depositor).






Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
3	A	1	Total C N O 14 8 1 5	0	0
3	A	1	Total C N O 14 8 1 5	0	0
3	A	1	Total C N O 14 8 1 5	0	0

• Molecule 4 is IODIDE ION (three-letter code: IOD) (formula: I) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
4	A	24	Total I 24 24	0	0

• Molecule 5 is THIOCYANATE ION (three-letter code: SCN) (formula: CNS) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0


Continued on next page...



 $Continued\ from\ previous\ page...$

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0
5	A	1	Total C N S 3 1 1 1	0	0


• Molecule 6 is TRIETHYLENE GLYCOL (three-letter code: PGE) (formula: $C_6H_{14}O_4$) (labeled as "Ligand of Interest" by depositor).





I	Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
	6	A	1	Total C C 10 6 4	0	0


• Molecule 7 is NITRITE ION (three-letter code: NO2) (formula: NO₂) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
7	A	1	Total N O 3 1 2	0	0
7	A	1	Total N O 3 1 2	0	0


• Molecule 8 is NITRIC OXIDE (three-letter code: NO) (formula: NO) (labeled as "Ligand of Interest" by depositor).





Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
8	A	1	Total N O 2 1 1	0	0
8	A	1	Total N O 2 1 1	0	0

• Molecule 9 is PROTOPORPHYRIN IX CONTAINING FE (three-letter code: HEM) (formula: $C_{34}H_{32}FeN_4O_4$) (labeled as "Ligand of Interest" by depositor).



Mol	Chain	Residues	Atoms			ZeroOcc	AltConf		
9	A	1	Total 43	C 34	Fe 1	N 4	O 4	0	0



• Molecule 10 is CALCIUM ION (three-letter code: CA) (formula: Ca) (labeled as "Ligand of Interest" by depositor).

Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
10	A	1	Total Ca 1 1	0	0

• Molecule 11 is water.


Mol	Chain	Residues	Atoms	ZeroOcc	AltConf
11	A	521	Total O 521 521	0	0



3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: Lactoperoxidase



• Molecule 2: 2-acetamido-2-deoxy-beta-D-glucopyranose-(1-4)-2-acetamido-2-deoxy-beta-D-glucopyranose

Chain C: 100%





4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 1 21 1	Depositor
Cell constants	54.02Å 80.41Å 75.84Å	Donositor
a, b, c, α , β , γ	90.00° 103.12° 90.00°	Depositor
Resolution (Å)	73.86 - 1.72	Depositor
Resolution (A)	73.86 - 1.72	EDS
% Data completeness	99.0 (73.86-1.72)	Depositor
(in resolution range)	99.0 (73.86-1.72)	EDS
R_{merge}	0.06	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) > 1$	1.27 (at 1.72Å)	Xtriage
Refinement program	REFMAC 5.8.0425	Depositor
D D.	0.186 , 0.235	Depositor
R, R_{free}	0.196 , 0.244	DCC
R_{free} test set	1330 reflections (1.99%)	wwPDB-VP
Wilson B-factor (Å ²)	36.0	Xtriage
Anisotropy	0.234	Xtriage
Bulk solvent $k_{sol}(e/Å^3)$, $B_{sol}(Å^2)$	0.35, 57.4	EDS
L-test for twinning ²	$ < L >=0.50, < L^2>=0.34$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o, F_c correlation	0.97	EDS
Total number of atoms	5522	wwPDB-VP
Average B, all atoms (Å ²)	53.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: The largest off-origin peak in the Patterson function is 5.69% of the height of the origin peak. No significant pseudotranslation is detected.

²Theoretical values of <|L|>, $<L^2>$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.



¹Intensities estimated from amplitudes.

5 Model quality (i)

5.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: SCN, IOD, NO2, NO, NAG, HEM, CA, PGE, CSD

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mal	Chain	Bond	lengths	Bond angles		
IVIOI	Chain	RMSZ	# Z > 5	RMSZ	# Z > 5	
1	A	0.40	0/4913	0.84	4/6667 (0.1%)	

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a maintain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	A	0	2

There are no bond length outliers.

All (4) bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
1	A	167	CYS	CB-CA-C	6.84	124.07	110.40
1	A	418	ARG	CB-CA-C	6.52	123.45	110.40
1	A	17	GLU	CB-CA-C	5.04	120.48	110.40
1	A	11	PRO	N-CA-CB	-5.01	97.09	102.60

There are no chirality outliers.

All (2) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	A	551	ARG	Sidechain
1	A	67	ARG	Sidechain



5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	4792	0	4703	69	0
2	С	28	0	25	1	0
3	A	42	0	39	0	0
4	A	24	0	0	27	0
5	A	51	0	0	11	0
6	A	10	0	14	3	0
7	A	6	0	0	5	0
8	A	4	0	0	7	0
9	A	43	0	30	16	0
10	A	1	0	0	0	0
11	A	521	0	0	14	0
All	All	5522	0	4811	96	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 10.

The worst 5 of 96 close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	$\begin{array}{c} {\rm Interatomic} \\ {\rm distance} \ ({\rm \AA}) \end{array}$	$\begin{array}{c} \text{Clash} \\ \text{overlap } (\text{\AA}) \end{array}$
1:A:108:ASP:OD2	9:A:639:HEM:CMD	1.67	1.40
4:A:608:IOD:I	5:A:628:SCN:N	0.83	1.38
1:A:258:GLU:OE2	9:A:639:HEM:CMB	1.72	1.37
5:A:621:SCN:N	11:A:701:HOH:O	1.66	1.28
4:A:608:IOD:I	5:A:628:SCN:C	1.44	1.21

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.



The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	594/595 (100%)	567 (96%)	20 (3%)	7 (1%)	11 2

5 of 7 Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	167	CYS
1	A	11	PRO
1	A	56	ALA
1	A	169	THR
1	A	10	VAL

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles		
1	A	519/517 (100%)	506 (98%)	13 (2%)	42 24		

5 of 13 residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type		
1	A	347	PHE		
1	A	485	LYS		
1	A	574	HIS		
1	A	513	CYS		
1	A	543	SER		

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	A	135	GLN
1	A	521	GLN



5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

2 non-standard protein/DNA/RNA residues are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			В	ond ang	gles
					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
1	CSD	A	441[A]	1	3,7,8	0.68	0	1,8,10	1.19	0
1	CSD	A	441[B]	1	3,7,8	0.56	0	1,8,10	0.96	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
1	CSD	A	441[A]	1	-	0/2/6/8	-
1	CSD	A	441[B]	1	-	1/2/6/8	-

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms	
1	A	441[B]	CSD	N-CA-CB-SG	

There are no ring outliers.

1 monomer is involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
1	A	441[A]	CSD	1	0



5.5 Carbohydrates (i)

2 monosaccharides are modelled in this entry.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
MIOI					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
2	NAG	С	1	2,1	14,14,15	0.50	0	17,19,21	0.88	0
2	NAG	С	2	2	14,14,15	0.54	0	17,19,21	0.90	0

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

\mathbf{Mol}	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	NAG	С	1	2,1	-	2/6/23/26	0/1/1/1
2	NAG	С	2	2	-	2/6/23/26	0/1/1/1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (4) torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
2	С	2	NAG	C8-C7-N2-C2
2	С	2	NAG	O7-C7-N2-C2
2	С	1	NAG	O5-C5-C6-O6
2	С	1	NAG	C4-C5-C6-O6


There are no ring outliers.

2 monomers are involved in 1 short contact:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
2	С	2	NAG	1	0
2	С	1	NAG	1	0



The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for oligosaccharide.



5.6 Ligand geometry (i)

Of 51 ligands modelled in this entry, 25 are monoatomic - leaving 26 for Mogul analysis.

In the following table, the Counts columns list the number of bonds (or angles) for which Mogul statistics could be retrieved, the number of bonds (or angles) that are observed in the model and the number of bonds (or angles) that are defined in the Chemical Component Dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 2 is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Type	Chain	Res	Link	Bond lengths			Bond angles		
IVIOI					Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SCN	A	621	-	1,2,2	0.20	0	0,1,1	-	-



Mal	Т	Clasia	Dan	T !1.	Во	ond leng	ths	В	ond ang	gles
Mol	Type	Chain	Res	Link	Counts	RMSZ	# Z > 2	Counts	RMSZ	# Z > 2
5	SCN	A	626	-	1,2,2	0.06	0	0,1,1	-	-
3	NAG	A	638	1	14,14,15	0.51	0	17,19,21	1.52	2 (11%)
5	SCN	A	624	-	1,2,2	0.38	0	0,1,1	-	-
5	SCN	A	627	-	1,2,2	0.28	0	0,1,1	-	-
5	SCN	A	619	-	1,2,2	1.52	0	0,1,1	-	-
5	SCN	A	618	-	1,2,2	1.29	0	0,1,1	-	-
6	PGE	A	632	-	9,9,9	0.19	0	8,8,8	0.24	0
8	NO	A	635	-	0,1,1	-	-	-		
5	SCN	A	616	-	1,2,2	0.12	0	0,1,1	-	-
5	SCN	A	625	ı	1,2,2	0.03	0	0,1,1	ı	-
5	SCN	A	629	-	1,2,2	0.54	0	0,1,1	ı	-
5	SCN	A	622	-	1,2,2	0.55	0	0,1,1	-	-
5	SCN	A	633	-	1,2,2	0.43	0	0,1,1	-	-
7	NO2	A	634	-	1,2,2	0.68	0	0,1,1	-	-
8	NO	A	636	-	0,1,1	-	-	-		
5	SCN	A	623	-	1,2,2	0.66	0	0,1,1	_	-
5	SCN	A	617	-	1,2,2	1.09	0	0,1,1	_	-
3	NAG	A	615	1	14,14,15	0.41	0	17,19,21	1.03	1 (5%)
7	NO2	A	637	4	1,2,2	1.06	0	0,1,1	-	-
9	HEM	A	639	1	41,50,50	1.34	5 (12%)	45,82,82	2.04	14 (31%)
5	SCN	A	630	4	1,2,2	0.67	0	0,1,1	-	-
5	SCN	A	628	4	1,2,2	0.82	0	0,1,1	-	-
5	SCN	A	631	-	1,2,2	1.00	0	0,1,1	-	-
5	SCN	A	620	-	1,2,2	0.08	0	0,1,1	-	-
3	NAG	A	601	1	14,14,15	0.45	0	17,19,21	0.90	1 (5%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the Chemical Component Dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
9	HEM	A	639	1	-	4/12/54/54	-
3	NAG	A	638	1	-	2/6/23/26	0/1/1/1
6	PGE	A	632	-	-	4/7/7/7	-
3	NAG	A	601	1	-	2/6/23/26	0/1/1/1
3	NAG	A	615	1	-	2/6/23/26	0/1/1/1

All (5) bond length outliers are listed below:



Mol	Chain	Res	Type	Atoms	Z	$\operatorname{Observed}(\text{\AA})$	$Ideal(\AA)$
9	A	639	HEM	C1B-NB	-3.65	1.34	1.40
9	A	639	HEM	C4D-ND	-3.12	1.34	1.40
9	A	639	HEM	FE-NB	2.90	2.11	1.96
9	A	639	HEM	CHB-C1B	2.77	1.42	1.35
9	A	639	HEM	CHA-C4D	2.16	1.40	1.35

The worst 5 of 18 bond angle outliers are listed below:

Mol	Chain	Res	Type	Atoms	\mathbf{Z}	$Observed(^o)$	$Ideal(^{o})$
9	A	639	HEM	C1B-NB-C4B	5.11	110.35	105.07
3	A	638	NAG	C1-O5-C5	5.09	119.09	112.19
9	A	639	HEM	CMD-C2D-C1D	4.21	131.45	125.04
9	A	639	HEM	CBA-CAA-C2A	-4.09	105.64	112.62
9	A	639	HEM	C4B-CHC-C1C	3.81	127.59	122.56

There are no chirality outliers.

5 of 14 torsion outliers are listed below:

Mol	Chain	Res	Type	Atoms
3	A	638	NAG	C8-C7-N2-C2
6	A	632	PGE	C6-C5-O3-C4
3	A	601	NAG	O5-C5-C6-O6
6	A	632	PGE	O1-C1-C2-O2
3	A	601	NAG	C4-C5-C6-O6

There are no ring outliers.


11 monomers are involved in 37 short contacts:

Mol	Chain	Res	Type	Clashes	Symm-Clashes
5	A	621	SCN	1	0
6	A	632	PGE	3	0
8	A	635	NO	6	0
5	A	629	SCN	2	0
5	A	622	SCN	5	0
7	A	634	NO2	3	0
8	A	636	NO	1	0
7	A	637	NO2	2	0
9	A	639	HEM	16	0
5	A	628	SCN	2	0
5	Ā	620	SCN	1	0


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths,




bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier. The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.













































5.7 Other polymers (i)

There are no such residues in this entry.



5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ>2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95^{th} percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ $>$	# RSRZ > 2	$OWAB(Å^2)$	Q < 0.9
1	A	594/595 (99%)	0.34	24 (4%) 43 47	26, 45, 114, 204	2 (0%)

The worst 5 of 24 RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	A	12	LEU	5.1
1	A	10	VAL	5.1
1	A	173	GLN	5.1
1	A	119	LEU	3.8
1	A	8	ALA	3.7

6.2 Non-standard residues in protein, DNA, RNA chains (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum, median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
1	CSD	A	441[A]	8/9	0.95	0.12	35,41,58,59	8
1	CSD	A	441[B]	8/9	0.95	0.12	28,29,32,33	8

6.3 Carbohydrates (i)

SUGAR-RSR INFOmissingINFO

6.4 Ligands (i)

In the following table, the Atoms column lists the number of modelled atoms in the group and the number defined in the chemical component dictionary. The B-factors column lists the minimum,



median, 95^{th} percentile and maximum values of B factors of atoms in the group. The column labelled 'Q< 0.9' lists the number of atoms with occupancy less than 0.9.

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\operatorname{B-factors}(\mathring{\mathbf{A}}^2)$	Q < 0.9
3	NAG	A	638	14/15	0.71	0.13	76,89,94,96	0
3	NAG	A	601	14/15	0.84	0.10	66,76,88,90	0
5	SCN	A	630	3/3	0.88	0.11	63,63,75,76	3
3	NAG	A	615	14/15	0.90	0.10	52,68,73,80	0
4	IOD	A	650	1/1	0.91	0.10	73,73,73,73	1
4	IOD	A	645	1/1	0.91	0.11	73,73,73,73	1
5	SCN	A	633	3/3	0.91	0.10	65,65,65,69	0
7	NO2	A	637	3/3	0.91	0.09	43,43,44,45	0
5	SCN	A	629	3/3	0.92	0.10	23,23,24,29	3
6	PGE	A	632	10/10	0.92	0.10	60,61,67,72	0
5	SCN	A	626	3/3	0.92	0.09	36,36,48,60	0
8	NO	A	635	2/2	0.92	0.12	37,37,37,45	2
5	SCN	A	619	3/3	0.94	0.08	19,19,23,40	3
4	IOD	A	609	1/1	0.94	0.09	70,70,70,70	1
4	IOD	A	643	1/1	0.94	0.09	62,62,62,62	1
4	IOD	A	651	1/1	0.94	0.08	61,61,61,61	1
5	SCN	A	617	3/3	0.95	0.07	31,31,32,32	3
5	SCN	A	627	3/3	0.95	0.08	50,50,53,66	0
5	SCN	A	628	3/3	0.95	0.10	30,30,46,55	3
5	SCN	A	622	3/3	0.96	0.05	20,20,23,30	3
5	SCN	A	623	3/3	0.96	0.10	41,41,47,57	0
5	SCN	A	625	3/3	0.96	0.06	25,25,27,40	3
4	IOD	A	644	1/1	0.96	0.07	51,51,51,51	1
4	IOD	A	610	1/1	0.96	0.12	59,59,59,59	1
4	IOD	A	607	1/1	0.96	0.06	65,65,65,65	1
8	NO	A	636	2/2	0.96	0.07	40,40,40,48	0
5	SCN	A	616	3/3	0.97	0.06	32,32,38,43	3
4	IOD	A	648	1/1	0.97	0.06	60,60,60,60	1
5	SCN	A	618	3/3	0.97	0.07	43,43,45,50	0
4	IOD	A	614	1/1	0.97	0.07	51,51,51,51	1
5	SCN	A	620	3/3	0.97	0.06	27,27,27,30	3
4	IOD	A	606	1/1	0.97	0.08	91,91,91,91	1
5	SCN	A	624	3/3	0.98	0.07	39,39,43,52	0
5	SCN	A	631	3/3	0.98	0.05	20,20,20,25	3
4	IOD	A	642	1/1	0.98	0.08	66,66,66,66	1
4	IOD	A	647	1/1	0.98	0.07	58,58,58,58	1
7	NO2	A	634	3/3	0.98	0.04	30,30,31,31	3
5	SCN	A	621	3/3	0.98	0.10	30,30,32,32	3
4	IOD	A	608	1/1	0.98	0.05	59,59,59,59	1
4	IOD	A	603	1/1	0.98	0.04	53,53,53,53	1
9	HEM	A	639	43/43	0.98	0.05	26,31,35,38	0


Continued on next page...



Continued from previous page...

Mol	Type	Chain	Res	Atoms	RSCC	RSR	$\mathbf{B} ext{-}\mathbf{factors}(\mathbf{\mathring{A}}^2)$	Q<0.9
4	IOD	A	612	1/1	0.99	0.03	41,41,41,41	1
4	IOD	A	605	1/1	0.99	0.04	39,39,39,39	1
4	IOD	A	646	1/1	0.99	0.06	49,49,49,49	1
4	IOD	A	604	1/1	0.99	0.04	45,45,45,45	1
4	IOD	A	611	1/1	0.99	0.03	47,47,47,47	1
4	IOD	A	641	1/1	1.00	0.01	27,27,27,27	1
4	IOD	A	613	1/1	1.00	0.04	48,48,48,48	1
4	IOD	A	649	1/1	1.00	0.05	41,41,41,41	1
4	IOD	A	602	1/1	1.00	0.05	32,32,32,32	1
10	CA	A	640	1/1	1.00	0.02	34,34,34,34	0


The following is a graphical depiction of the model fit to experimental electron density of all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the geometry validation Tables will also be included. Each fit is shown from different orientation to approximate a three-dimensional view.







Electron density around NAG A 601:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SCN A 630:


 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)

















Electron density around IOD A 650:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around SCN A 633:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NO2 A 637:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around SCN A 629:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around PGE A 632:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around SCN A 626:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around NO A 635:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)













Electron density around SCN A 619: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray

and green (positive)















Electron density around IOD A 609:





















Electron density around SCN A 617:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SCN A 627:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around SCN A 628:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)











Electron density around SCN A 622: 2mF_o-DF_c (at 0.7 rmsd) in gray mF_o-DF_c (at 3 rmsd) in purple (negative) and green (positive)




Electron density around SCN A 623:

 $2 {
m mF}_o {
m -DF}_c$ (at 0.7 rmsd) in gray ${
m mF}_o {
m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around SCN A 625:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around IOD A 644: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)



Electron density around IOD A 610:












Electron density around IOD A 607:












Electron density around NO A 636:

 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SCN A 616:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around IOD A 648: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray ${ m mF}_o{ m -DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)














Electron density around IOD A 606:













Electron density around SCN A 624:
















Electron density around IOD A 642:


 $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)









Electron density around IOD A 647:












Electron density around NO2 A 634:

 $2 {\rm mF}_o\text{-}{\rm DF}_c$ (at 0.7 rmsd) in gray ${\rm mF}_o\text{-}{\rm DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)










Electron density around SCN A 621:

 $2 \mathrm{mF}_o\text{-DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around IOD A 608: $2 \text{mF}_o\text{-DF}_c \text{ (at 0.7 rmsd) in gray} \\ \text{mF}_o\text{-DF}_c \text{ (at 3 rmsd) in purple (negative)}$

and green (positive)













Electron density around IOD A 603:



















Electron density around IOD A 605: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_{o}\text{-}\mathrm{DF}_{c}$ (at 3 rmsd) in purple (negative) and green (positive)












Electron density around IOD A 611:












Electron density around IOD A 641:

 $2mF_o$ -DF_c (at 0.7 rmsd) in gray mF_o -DF_c (at 3 rmsd) in purple (negative) and green (positive)












Electron density around IOD A 613:












Electron density around IOD A 649:










Electron density around IOD A 602: $2 \mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 0.7 rmsd) in gray $\mathrm{mF}_o\text{-}\mathrm{DF}_c$ (at 3 rmsd) in purple (negative) and green (positive)





6.5 Other polymers (i)

There are no such residues in this entry.

