

Full wwPDB NMR Structure Validation Report (i)

Sep 21, 2021 – 10:16 AM JST

PDB ID	:	6KXZ
Title	:	HUMAN PARALLEL STRANDED 7-MER G-QUADRUPLEX COM-
		PLEXED WITH 2 EPIRUBICIN (EPI) MOLECULES
Authors	:	Barthwal, R.; Raje, S.; Pandav, K.
Deposited on	:	2019-09-15

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org* A user guide is available at https://www.wwpdb.org/validation/2017/NMRValidationReportHelp with specific help available everywhere you see the (i) symbol.

The following versions of software and data (see references (1)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	1.8.5 (274361), CSD as541be (2020)
buster-report	:	1.1.7(2018)
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
RCI	:	v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV	:	Wang et al. (2010)
ShiftChecker	:	2.23.1
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP)	:	2.23.1

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure: $SOLUTION\ NMR$

The overall completeness of chemical shifts assignment is 10%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.

Metric	Percen	tile Ranks	Value
Clashscore			0
	Worse		Better
	Percentile relative to all structur	es	
	Percentile relative to all NMR str	ructures	
	TT71 1 1 •		1
Motric	Whole archive	NMR archive	

Metric	$egin{array}{c} { m Whole \ archive} \ (\#{ m Entries}) \end{array}$	${f NMR} { m archive} \ (\#{ m Entries})$
Clashscore	158937	12864

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5%

Mol	Chain	Length		Quality of chain
1	А	7	43%	57%
1	В	7	43%	57%
1	С	7	29%	71%
1	D	7	14%	86%



2 Ensemble composition and analysis (i)

This entry contains 15 models. This entry does not contain polypeptide chains, therefore identification of well-defined residues and clustering analysis are not possible. All residues are included in the validation scores.




3 Entry composition (i)

There are 2 unique types of molecules in this entry. The entry contains 1036 atoms, of which 382 are hydrogens and 0 are deuteriums.

• Molecule 1 is a DNA chain called [TTAGGGT]4-EPI-Complex.

Mol	Chain	Residues		A	Aton	ns			Trace
1	Λ	7	Total	С	Η	Ν	Ο	Р	0
	A	1	225	70	81	26	42	6	0
1	В	7	Total	С	Η	Ν	Ο	Р	0
	D	1	225	70	81	26	42	6	0
1	С	7	Total	С	Η	Ν	Ο	Р	0
	U	1	225	70	81	26	42	6	0
1	Л	7	Total	С	Η	Ν	Ο	Р	0
	D	1	225	70	81	26	42	6	0

• Molecule 2 is 4'-EPIDOXORUBICIN (three-letter code: DM6) (formula: $C_{27}H_{30}NO_{11}$).



Mol	Chain	Residues		At	oms		
0	۸	1	Total	С	Η	Ν	0
	A	1	68	27	29	1	11
0	٨	1	Total	С	Η	Ν	0
	A	1	68	27	29	1	11



4 Residue-property plots (i)

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

• Molecule 1: [TTAGGGT]4-EPI-Complex

Chain A:	43%	57%	
T1 T2 A3 T7			
• Molecule	e 1: [TTAGGGT]4-EPI-Comp	lex	
Chain B:	43%	57%	
T8 T9 T14 T14			
• Molecule	e 1: [TTAGGGT]4-EPI-Comp	blex	
_			
Chain C:	29%	71%	
T15 T16 A17 G20 T21			
• Molecule	e 1: [TTAGGGT]4-EPI-Comp	olex	
Chain D.			
Chain D:	14%	86%	
T22 T23 A24 G25 G26 G27 T28			

4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.



4.2.1 Score per residue for model 1

Chain A:	43%	43%	14%
11 12 A3 17			
• Molecule 1:	[TTAGGGT]4-EPI-Cor	nplex	
Chain B:	43%	57%	
T8 T9 T14			
• Molecule 1:	[TTAGGGT]4-EPI-Cor	nplex	
Chain C:	43%	57%	
115 116 117 121			
• Molecule 1:	[TTAGGGT]4-EPI-Cor	nplex	
Chain D: 14	4%	86%	
122 123 424 625 626 128 128			
4.2.2 Score	e per residue for mod	el 2	
• Molecule 1:	[TTAGGGT]4-EPI-Cor	nplex	
Chain A:	43%	57%	
11 12 43 17			
• Molecule 1:	[TTAGGGT]4-EPI-Cor	nplex	
Chain B:	29%	71%	
T8 T9 611 T14			
• Molecule 1:	[TTAGGGT]4-EPI-Cor	nplex	
Chain C:	29%	57%	14%



T15 T16 A17 G20 T21


• Molecule 1: [TTAGGGT]4-EPI-Complex

Chain D:	29%	57%	14%
122 123 123 124 128 128			

- 4.2.3 Score per residue for model 3
- Molecule 1: [TTAGGGT]4-EPI-Complex

Chain A:	43%	43%	14%
T1 T2 A3 T7			
• Molecule	e 1: [TTAGGGT]4-EPI-Cor	nplex	
Chain B:	43%	57%	
T8 T9 A10 T14			
• Molecule	e 1: [TTAGGGT]4-EPI-Cor	nplex	
Chain C:	29%	71%	
T15 T16 A17 G20 T21			
• Molecule	e 1: [TTAGGGT]4-EPI-Cor	nplex	
Chain D:	29%	71%	
T22 T23 A24 G25 G26 G27 T28			

4.2.4 Score per residue for model 4



• Molecule 1:	[TTAGGGT]4-EP		
Chain B:	29%	71%	
T8 T19 G11 T14			
• Molecule 1:	[TTAGGGT]4-EP	I-Complex	
Chain C:	29%	57%	14%
115 116 117 118 118 121			
• Molecule 1:	[TTAGGGT]4-EP	PI-Complex	
Chain D:	29%	71%	
T22 T23 A24 G25 G26 T28 T28			
	e per residue for [TTAGGGT]4-EP		
Chain A:	43%	57%	
Chain A:	43%	57%	
T1 T2 A3 T7	43% [TTAGGGT]4-EP		
T1 T2 A3 T7			
<mark>₽₿₽</mark> ► • Molecule 1:	[TTAGGGT]4-EP	'I-Complex	
<mark>문 원 및</mark> • Molecule 1: Chain B: 말 윤 원 등 문	[TTAGGGT]4-EP	PI-Complex 71%	
<mark>문 원 및</mark> • Molecule 1: Chain B: 말 윤 원 등 문	[TTAGGGT]4-EP 29%	PI-Complex 71%	
타 유 및 타 • Molecule 1: Chain B: 윤 유 및 타 • Molecule 1:	[TTAGGGT]4-EP 29% [TTAGGGT]4-EP	'I-Complex 71% 'I-Complex	
E E P P E • Molecule 1: Chain B: E E P E E E • Molecule 1: Chain C: E E E E E E E E E	[TTAGGGT]4-EP 29% [TTAGGGT]4-EP	PI-Complex 71% PI-Complex 71%	
E E P P E • Molecule 1: Chain B: E E P E E E • Molecule 1: Chain C: E E E E E E E E E	[TTAGGGT]4-EP 29% [TTAGGGT]4-EP 29%	PI-Complex 71% PI-Complex 71%	29%



4.2.6 Score per residue for model 6

Chain A:	43%	57%	
11 12 13 17			
• Molecule 1:	[TTAGGGT]4-EPI-Comp	lex	
Chain B:	43%	57%	
T9 A10 T14			
• Molecule 1:	[TTAGGGT]4-EPI-Comp	lex	
Chain C:	29%	57%	14%
115 116 A17 G18 T21			
• Molecule 1:	[TTAGGGT]4-EPI-Comp	lex	
Chain D: 1	4%	71%	14%
122 123 424 625 625 728 128			
4.2.7 Score	e per residue for model	7	
• Molecule 1:	[TTAGGGT]4-EPI-Comp	lex	
Chain A:	29%	71%	
11 12 43 17			
• Molecule 1:	[TTAGGGT]4-EPI-Comp	lex	
Chain B:	43%	57%	
T8 T9 T10 T14			
• Molecule 1:	[TTAGGGT]4-EPI-Comp	lex	
Chain C:	29%	43%	29%



T15 T16 A17 G18 G19 G20 T21

• Molecule 1: [TTAGGGT]4-EPI-Complex

Chain D:	43%	57%
122 123 124 128 128		

4.2.8 Score per residue for model 8

• Molecule 1: [TTAGGGT]4-EPI-Complex

Chain A:	43%	57%	
T1 T2 A3 T7			
• Molecule	e 1: [TTAGGGT]4-EPI	-Complex	
Chain B:	29%	57%	14%
T8 T9 A10 G11 G12 G12 G13 T14			
• Molecule	e 1: [TTAGGGT]4-EPI	-Complex	
Chain C:	29%	71%	
T15 T16 A17 G20 T21			
• Molecule	e 1: [TTAGGGT]4-EPI	-Complex	
Chain D:	29%	57%	14%
T22 T23 A24 G25 T28			

4.2.9 Score per residue for model 9





• Molecule 1: [TTAGGGT]4-EPI-Complex

Chain B:	43%		57%	
T9 19 114				
• Molecule 1: ['	TTAGGGT]4-EF	PI-Complex		
Chain C: 14%	6	71%		14%
115 116 A17 G18 G19 G20 T21				
• Molecule 1: ['	TTAGGGT]4-EF	PI-Complex		
Chain D:	29%		71%	
T22 T23 A24 G25 T28				
4.2.10 Score	e per residue fo	r model 10		
• Molecule 1: ['	TTAGGGT]4-EF	PI-Complex		
Chain A:	43%		43%	14%
11 12 13 17				
• Molecule 1: ['	TTAGGGT]4-EF	PI-Complex		
Chain B:	43%		57%	
T8 19 14 14				
• Molecule 1: [TTAGGGT]4-EF	PI-Complex		
Chain C:	43%		43%	14%
115 116 117 121				

Chain D:	14%	71%	14%
T22 T23 A24 G25 G26 T28			

4.2.11 Score per residue for model 11

Chain A: 43%	57%	
11 12 17		
• Molecule 1: [TTAGGGT]4-2	EPI-Complex	
Chain B: 43%	57%	
18 10 14		
• Molecule 1: [TTAGGGT]4-2	EPI-Complex	
Chain C: 29%	71%	
115 116 117 120 121		
• Molecule 1: [TTAGGGT]4-2	EPI-Complex	
Chain D: 14%	71% 14	4%
122 123 025 026 027 128		
4.2.12 Score per residue	for model 12	
• Molecule 1: [TTAGGGT]4-	EPI-Complex	
Chain A: 29%	71%	
11 8 8 17 17		
• Molecule 1: [TTAGGGT]4-	EPI-Complex	
Chain B: 43%	57%	
18 10 14		
• Molecule 1: [TTAGGGT]4-	EPI-Complex	
Chain C: 43%	57%	



T15 T16 A17 T21

• Molecule 1: [TTAGGGT]4-EPI-Complex

Chain D: 29% 71%

- 4.2.13 Score per residue for model 13
- \bullet Molecule 1: [TTAGGGT]4-EPI-Complex

Chain A: 43% 57% • Molecule 1: [TTAGGGT]4-EPI-Complex Chain B: 43% 57% • Molecule 1: [TTAGGGT]4-EPI-Complex Chain C: 29% 43% 29% • Molecule 1: [TTAGGGT]4-EPI-Complex Chain D: 86% 14% T22 T23 A24 G25 G26 G26 G26 G27 T28

4.2.14 Score per residue for model 14



• Molecule 1: [[TTAGGGT]4-EPI-Comp	blex	
Chain B:	43%	57%	
T8 T9 A10 T14			
• Molecule 1: [[TTAGGGT]4-EPI-Comp	blex	
Chain C:	43%	57%	
T15 T16 A17 T21			
• Molecule 1: [[TTAGGGT]4-EPI-Comp	blex	
Chain D:	29%	71%	
T22 T23 A24 G25 T28			
4.2.15 Score	e per residue for mode	el 15	
• Molecule 1: [[TTAGGGT]4-EPI-Comp	blex	
Chain A:	29%	57%	14%
11 83 68 8 98			
• Molecule 1: [[TTAGGGT]4-EPI-Comp	blex	
Chain B:	29%	71%	
T8 A10 G11 T14			
• Molecule 1: [[TTAGGGT]4-EPI-Comp	blex	
Chain C:	43%	57%	
T15 T16 A17 T21			
• Molecule 1: [[TTAGGGT]4-EPI-Comp	blex	
Chain D: 14	%	71%	14%
T22 T23 A24 G25 G26 G27 T28			



5 Refinement protocol and experimental data overview (i)

The models were refined using the following method: *simulated annealing*.

Of the 35 calculated structures, 15 were deposited, based on the following criterion: *structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
Amber	refinement	
TopSpin	structure solution	3.6

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	working_cs.cif
Number of chemical shift lists	1
Total number of shifts	65
Number of shifts mapped to atoms	65
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	10%



6 Model quality (i)

6.1 Standard geometry (i)

Bond lengths and bond angles in the following residue types are not validated in this section: $\mathrm{DM6}$

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with |Z| > 5 is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol Chain RMS		B	ond lengths	Bond angles	
		RMSZ	$\#Z{>}5$	RMSZ	#Z > 5
1	А	$1.69 {\pm} 0.01$	$0{\pm}0/161~(~0.0{\pm}~0.0\%)$	2.27 ± 0.02	$10{\pm}1/248~(~4.0{\pm}~0.4\%)$
1	В	$1.67 {\pm} 0.01$	$0{\pm}0/161~(~0.0{\pm}~0.0\%)$	$2.24{\pm}0.04$	$9{\pm}1/248~(~3.8{\pm}~0.4\%)$
1	С	$1.66 {\pm} 0.02$	$0{\pm}0/161~(~0.0{\pm}~0.0\%)$	$2.33 {\pm} 0.04$	$10{\pm}1/248~(~4.0{\pm}~0.5\%)$
1	D	$1.65 {\pm} 0.01$	$0{\pm}0/161~(~0.0{\pm}~0.0\%)$	$2.39{\pm}0.04$	$12{\pm}1/248~(~4.8{\pm}~0.6\%)$
All	All	1.67	0/9660~(~0.0%)	2.31	614/14880~(~4.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	А	$0.0{\pm}0.0$	$0.6{\pm}0.8$
1	С	$0.0{\pm}0.0$	$0.6{\pm}0.7$
1	D	$0.0{\pm}0.0$	$0.6{\pm}0.6$
1	В	$0.0{\pm}0.0$	$0.1{\pm}0.3$
All	All	0	29

There are no bond-length outliers.

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mal	Chain	hain Res Type Atoms Z Observed		Observed ⁽⁰⁾	$Ideal(^{o})$	Models			
Mol	Chain	nes	туре	Atoms		$\mathbf{Observed}(^{o})$	Ideal(')	Worst	Total
1	С	17	DA	N1-C6-N6	-10.60	112.24	118.60	9	15
1	А	3	DA	N1-C6-N6	-10.52	112.29	118.60	15	15
1	D	24	DA	N1-C6-N6	-10.28	112.43	118.60	1	15
1	В	10	DA	N1-C6-N6	-9.63	112.82	118.60	4	15
1	D	24	DA	C5-C6-N1	8.19	121.79	117.70	4	15
1	А	3	DA	C5-C6-N1	8.07	121.73	117.70	1	15

Continued on next page...



Mol	Chain	Res	Type	Atoms	Z	Observed(°)	$Ideal(^{o})$	Mo	dels
WIOI	Ullalli	nes	туре	Atoms		Observed()	Ideal()	Worst	Total
1	С	17	DA	C5-C6-N1	8.07	121.74	117.70	12	15
1	В	10	DA	C5-C6-N1	7.85	121.62	117.70	1	15
1	В	11	DG	O4'-C1'-N9	7.49	113.25	108.00	15	3
1	D	26	DG	O4'-C1'-N9	7.43	113.20	108.00	4	8
1	С	17	DA	C4-C5-C6	-7.24	113.38	117.00	12	15
1	D	24	DA	C4-C5-C6	-7.21	113.39	117.00	15	15
1	А	3	DA	C4-C5-C6	-7.19	113.41	117.00	13	15
1	С	21	DT	C6-C5-C7	-7.17	118.60	122.90	13	15
1	В	10	DA	C4-C5-C6	-7.16	113.42	117.00	14	15
1	С	16	DT	C6-C5-C7	-6.98	118.71	122.90	15	15
1	D	23	DT	C6-C5-C7	-6.95	118.73	122.90	13	15
1	D	24	DA	O4'-C1'-N9	6.93	112.85	108.00	11	12
1	В	9	DT	C6-C5-C7	-6.93	118.74	122.90	1	15
1	А	1	DT	C6-C5-C7	-6.83	118.80	122.90	7	15
1	D	27	DG	O4'-C1'-N9	6.77	112.74	108.00	13	2
1	А	7	DT	C6-C5-C7	-6.66	118.91	122.90	12	15
1	В	14	DT	C6-C5-C7	-6.65	118.91	122.90	3	15
1	С	18	DG	O4'-C1'-N9	6.50	112.55	108.00	13	2
1	D	28	DT	C6-C5-C7	-6.41	119.05	122.90	3	15
1	В	8	DT	C6-C5-C7	-6.30	119.12	122.90	13	15
1	В	9	DT	O4'-C1'-N1	6.26	112.38	108.00	15	3
1	А	2	DT	C6-C5-C7	-6.24	119.16	122.90	15	15
1	С	15	DT	C6-C5-C7	-6.22	119.17	122.90	6	15
1	В	8	DT	O4'-C1'-N1	6.21	112.35	108.00	11	4
1	С	20	DG	O4'-C1'-N9	6.19	112.33	108.00	8	4
1	D	25	DG	O4'-C1'-N9	6.12	112.28	108.00	5	3
1	С	19	DG	O4'-C1'-N9	6.08	112.25	108.00	7	2
1	А	2	DT	N3-C2-O2	-6.02	118.69	122.30	8	14
1	D	22	DT	N3-C2-O2	-5.98	118.71	122.30	3	15
1	С	15	DT	O4'-C1'-N1	5.97	112.18	108.00	15	3
1	D	22	DT	C6-C5-C7	-5.97	119.32	122.90	15	15
1	А	1	DT	N3-C2-O2	-5.86	118.79	122.30	4	15
1	С	15	DT	N3-C2-O2	-5.81	118.81	122.30	11	15
1	A	1	DT	O4'-C1'-N1	5.79	112.05	108.00	4	4
1	D	23	DT	O4'-C1'-N1	5.68	111.98	108.00	10	4
1	А	3	DA	O4'-C1'-N9	5.64	111.95	108.00	13	4
1	С	21	DT	O4'-C1'-N1	5.63	111.94	108.00	11	3
1	С	21	DT	N3-C2-O2	-5.63	118.92	122.30	8	14
1	D	25	DG	N1-C6-O6	-5.61	116.53	119.90	9	10
1	В	8	DT	N3-C2-O2	-5.61	118.94	122.30	15	12
1	B	9	DT	N3-C2-O2	-5.60	118.94	122.30	4	15

Continued from previous page...

Continued on next page...



Mol	Chain	Dec	Turne	Atoma	Z	Observed ⁽⁰⁾	$Ideal(^{o})$	Mod	lels
IVIOI	Chain	Res	Type	Atoms	L	$\mathbf{Observed}(^{o})$	Ideal(*)	Worst	Total
1	D	28	DT	N3-C2-O2	-5.59	118.94	122.30	10	14
1	В	14	DT	O4'-C1'-N1	5.59	111.92	108.00	8	1
1	А	6	DG	O4'-C4'-C3'	5.59	109.35	106.00	14	2
1	D	23	DT	N3-C2-O2	-5.58	118.95	122.30	7	15
1	С	16	DT	O4'-C1'-N1	5.58	111.90	108.00	7	2
1	А	7	DT	N3-C2-O2	-5.54	118.97	122.30	2	15
1	В	14	DT	N3-C2-O2	-5.54	118.98	122.30	6	12
1	D	25	DG	O4'-C4'-C3'	5.51	109.31	106.00	14	1
1	С	20	DG	N3-C2-N2	-5.40	116.12	119.90	9	4
1	С	16	DT	N3-C2-O2	-5.39	119.06	122.30	5	9
1	D	27	DG	N3-C2-N2	-5.36	116.15	119.90	11	1
1	А	2	DT	O4'-C1'-N1	5.30	111.71	108.00	4	1
1	А	6	DG	O4'-C1'-N9	5.28	111.70	108.00	7	1
1	D	26	DG	C1'-O4'-C4'	-5.18	104.92	110.10	5	1
1	С	18	DG	N3-C2-N2	-5.17	116.28	119.90	6	1
1	D	24	DA	C1'-O4'-C4'	-5.14	104.96	110.10	11	1
1	В	12	DG	O4'-C1'-N9	5.12	111.58	108.00	8	1
1	А	5	DG	O4'-C1'-N9	5.01	111.51	108.00	15	1

Continued from previous page...

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	С	17	DA	Sidechain	5
1	D	24	DA	Sidechain	4
1	А	3	DA	Sidechain	4
1	D	25	DG	Sidechain	4
1	А	2	DT	Sidechain	2
1	С	20	DG	Sidechain	2
1	А	6	DG	Sidechain	1
1	А	7	DT	Sidechain	1
1	В	11	DG	Sidechain	1
1	D	26	DG	Sidechain	1
1	С	16	DT	Sidechain	1
1	В	14	DT	Sidechain	1
1	А	1	DT	Sidechain	1
1	С	18	DG	Sidechain	1



6.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
All	All	9810	5730	5820	-

The all-atom clash score is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clash score for this structure is -.

There are no clashes.

6.3 Torsion angles (i)

6.3.1 Protein backbone (i)

There are no protein molecules in this entry.

6.3.2 Protein sidechains (i)

There are no protein molecules in this entry.

6.3.3 RNA (i)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates (i)

There are no monosaccharides in this entry.

6.6 Ligand geometry (i)

2 ligands are modelled in this entry.



In the following table, the Counts columns list the number of bonds for which Mogul statistics could be retrieved, the number of bonds that are observed in the model and the number of bonds that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond length is the number of standard deviations the observed value is removed from the expected value. A bond length with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond lengths.

Mal	Trune	Chain	Dec	Tinle		Bond leng	gths
	туре	Chain	nes		Counts	RMSZ	#Z>2
2	DM6	А	101	-	41,43,43	$1.11 {\pm} 0.02$	2±1 (3±1%)
2	DM6	А	102	-	41,43,43	1.11 ± 0.02	1±1 (3±1%)

In the following table, the Counts columns list the number of angles for which Mogul statistics could be retrieved, the number of angles that are observed in the model and the number of angles that are defined in the chemical component dictionary. The Link column lists molecule types, if any, to which the group is linked. The Z score for a bond angle is the number of standard deviations the observed value is removed from the expected value. A bond angle with |Z| > 2 is considered an outlier worth inspection. RMSZ is the average root-mean-square of all Z scores of the bond angles.

Mal	Trune	Chain	Dec	Tinle		Bond ang	gles
IVIOI	туре	Chain	nes	LIIIK	Counts	RMSZ	$\#Z{>}2$
2	DM6	А	101	-	55,67,67	$1.00{\pm}0.06$	3 ± 1 (6±1%)
2	DM6	А	102	-	55,67,67	$1.01 {\pm} 0.07$	3 ± 1 (6±1%)

In the following table, the Chirals column lists the number of chiral outliers, the number of chiral centers analysed, the number of these observed in the model and the number defined in the chemical component dictionary. Similar counts are reported in the Torsion and Rings columns. '-' means no outliers of that kind were identified.

Mol	Type	Chain	Res	Link	Chirals	Torsions	Rings
2	DM6	А	102	-	-	$0\pm0,13,60,60$	$0\pm 0,5,5,5$
2	DM6	А	101	-	-	$0\pm0,13,60,60$	$0\pm 0,5,5,5$

All unique bond outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Dec	Turne	Atoma	Z	Observed(Å)	Ideal(Å)	Mod	dels
	Unain	nes	туре	Atoms		Observed(A)	Ideal(A)	Worst	Total
2	А	101	DM6	C8-C7	2.25	1.56	1.52	15	4
2	А	102	DM6	C20-C7	2.20	1.55	1.51	11	9
2	А	102	DM6	C8-C7	2.18	1.56	1.52	12	3
2	А	102	DM6	C17-C5	2.16	1.53	1.47	8	5

Continued on next page...



Mal	Chain	Dag	T 0	A 4 a a a a	7	Observed(Å)		Models	
IVI01	Chain	Res	Type	Atoms	L	Observed(A)	Ideal(A)	Worst	Total
2	А	101	DM6	C10-C19	2.15	1.53	1.51	3	8
2	А	101	DM6	C20-C7	2.13	1.55	1.51	9	8
2	А	102	DM6	C10-C19	2.12	1.53	1.51	9	3
2	А	101	DM6	C17-C5	2.08	1.52	1.47	1	3

Continued from previous page...

All unique angle outliers are listed below. They are sorted according to the Z-score of the worst occurrence in the ensemble.

Mol	Chain	Res	Tuno	Atoms	Z	Observed(°)	Ideal(°)	Mod	dels
MOI	Ullalli	nes	Type	Atoms	2	Observed()	Iueai()	Worst	Total
2	А	102	DM6	O4-C4-C16	4.81	122.55	115.85	14	15
2	А	102	DM6	C21-O4-C4	4.59	124.46	117.53	8	9
2	А	101	DM6	C21-O4-C4	4.52	124.34	117.53	13	9
2	А	101	DM6	O4-C4-C16	4.49	122.10	115.85	12	15
2	А	102	DM6	O4-C4-C3	4.42	116.80	124.37	14	15
2	А	101	DM6	O4-C4-C3	4.07	117.39	124.37	12	15
2	А	101	DM6	C6-C20-C7	2.31	122.21	118.59	7	4
2	А	102	DM6	C8-C7-C20	2.18	115.84	110.68	11	11
2	А	101	DM6	C8-C7-C20	2.18	115.83	110.68	15	6
2	А	102	DM6	C6-C20-C7	2.13	121.92	118.59	14	1
2	А	101	DM6	C19-C20-C7	2.05	117.62	120.96	7	1

There are no chirality outliers.

All unique torsion outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.


Mol	Chain	Res	Type	Atoms	Models (Total)
2	А	102	DM6	C9-C13-C14-O14	1
2	А	102	DM6	O13-C13-C14-O14	1

There are no ring outliers.


The following is a two-dimensional graphical depiction of Mogul quality analysis of bond lengths, bond angles, torsion angles, and ring geometry for all instances of the Ligand of Interest. In addition, ligands with molecular weight > 250 and outliers as shown on the validation Tables will also be included. For torsion angles, if less then 5% of the Mogul distribution of torsion angles is within 10 degrees of the torsion angle in question, then that torsion angle is considered an outlier. Any bond that is central to one or more torsion angles identified as an outlier by Mogul will be highlighted in the graph. For rings, the root-mean-square deviation (RMSD) between the ring in question and similar rings identified by Mogul is calculated over all ring torsion angles. If the average RMSD is greater than 60 degrees and the minimal RMSD between the ring in question and any Mogul-identified rings is also greater than 60 degrees, then that ring is considered an outlier.



The outliers are highlighted in purple. The color gray indicates Mogul did not find sufficient equivalents in the CSD to analyse the geometry.







6.7 Other polymers (i)

There are no such molecules in this entry.

6.8 Polymer linkage issues (i)

There are no chain breaks in this entry.



7 Chemical shift validation (i)

The completeness of assignment taking into account all chemical shift lists is 10% for the well-defined parts and 10% for the entire structure.

7.1 Chemical shift list 1

File name: working_cs.cif

Chemical shift list name: DM6-HTel7Mer-NMR.str

7.1.1 Bookkeeping (i)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	65
Number of shifts mapped to atoms	65
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing (i)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments (i)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 10%, i.e. 56 atoms were assigned a chemical shift out of a possible 548. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	0/0~(%)	0/0~(-%)	0/0 (%)	$0/0 \ (\%)$
Sidechain	0/0~(-%)	0/0~(-%)	0/0 (%)	$0/0 \ (\%)$
Aromatic	0/0~(-%)	0/0~(-%)	0/0 (%)	0/0 (%)
Overall	56/548~(10%)	56/324~(17%)	0/184~(0%)	0/40~(0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 10%, i.e. 56 atoms were assigned a chemical shift out of a possible 548. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.



	Total	$^{1}\mathbf{H}$	$^{13}\mathrm{C}$	$^{15}\mathbf{N}$
Backbone	0/0 (%)	0/0~(-%)	0/0 (%)	$0/0 \ (\%)$
Sidechain	0/0 (%)	0/0~(-%)	0/0 (%)	0/0 (%)
Aromatic	0/0~(-%)	0/0~(-%)	0/0 (%)	$0/0 \ (\%)$
Overall	56/548~(10%)	56/324~(17%)	0/184~(0%)	0/40~(0%)

7.1.4 Statistically unusual chemical shifts (i)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots (i)

No *random coil index* (RCI) plot could be generated from the current chemical shift list (DM6-HTel7Mer-NMR.str). RCI is only applicable to proteins.

