Full wwPDB X-ray Structure Validation Report (i)

Jun 19, 2024 – 06:30 AM EDT

PDB ID	:	4KY3
Title	:	Three-dimensional Structure of the orthorhombic crystal of computationally designed insertion domain , Northeast Structural Genomics Consortium (NESG) Target OR327
Authors	:	Kuzin, A.; Su, M.; Seetharaman, J.; Maglaqui, M.; Xiao, R.; Lee, D.; Gleixner, J.; Baker, D.; Everett, J.K.; Acton, T.B.; Montelione, G.T.; Tong, L.; Hunt, J.F.; Northeast Structural Genomics Consortium (NESG)
Deposited on	:	2013-05-28
Resolution	:	2.96 Å(reported)

This is a Full wwPDB X-ray Structure Validation Report for a publicly released PDB entry.

We welcome your comments at *validation@mail.wwpdb.org*
A user guide is available at
https://www.wwpdb.org/validation/2017/XrayValidationReportHelp
with specific help available everywhere you see the (i) symbol.

The types of validation reports are described at
http://www.wwpdb.org/validation/2017/FAQs#types.

The following versions of software and data (see references (i)) were used in the production of this report:

MolProbity	:	4.02b-467
Mogul	:	2022.3.0, CSD as543be (2022)
Xtriage (Phenix)	:	1.20.1
EDS	:	2.37.1
Percentile statistics	:	20191225.v01 (using entries in the PDB archive December 25th 2019)
Refmac	:	5.8.0158
CCP4	:	7.0.044 (Gargrove)
Ideal geometry (proteins)	:	Engh & Huber (2001)
Ideal geometry (DNA, RNA)	:	Parkinson et al. (1996)

1 Overall quality at a glance (i)

The following experimental techniques were used to determine the structure:
X-RAY DIFFRACTION

The reported resolution of this entry is 2.96 Å.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	Similar resolution (#Entries, resolution range(Å))
R_{free}	130704	3104 (3.00-2.92)
Clashscore	141614	3462 (3.00-2.92)
Ramachandran outliers	138981	3340 (3.00-2.92)
Sidechain outliers	138945	3343 (3.00-2.92)
RSRZ outliers	127900	2986 (3.00-2.92)

The table below summarises the geometric issues observed across the polymeric chains and their fit to the electron density. The red, orange, yellow and green segments of the lower bar indicate the fraction of residues that contain outliers for >=3, 2, 1 and 0 types of geometric quality criteria respectively. A grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions <=5% The upper red bar (where present) indicates the fraction of residues that have poor fit to the electron density. The numeric value is given above the bar.

Mol	Chain	Length	Quality of chain
1	A	172	48% 34% 10% 7%
1	B	172	51% 34% 8% • 7%
1	C	172	3% 48% 37% 8% 7%

Validation Pipeline (wwPDB-VP) : 2.37.1

2 Entry composition (i)

There is only 1 type of molecule in this entry. The entry contains 3839 atoms, of which 0 are hydrogens and 0 are deuteriums.

In the tables below, the ZeroOcc column contains the number of atoms modelled with zero occupancy, the AltConf column contains the number of residues with at least one atom in alternate conformation and the Trace column contains the number of residues modelled with at most 2 atoms.

- Molecule 1 is a protein called designed protein OR327.

Mol	Chain	Residues	Atoms					ZeroOcc	AltConf	Trace
			Total	C	N	O	Se			
1	A	160	1281	809	219	250	3	0	0	0
			Total	C	N	O	Se			
1	B	160	1277	807	219	248	3	0	0	0
			Total	C	N	O	Se			
1	C	160	1281	809	219	250	3	0	0	0

3 Residue-property plots (i)

These plots are drawn for all protein, RNA, DNA and oligosaccharide chains in the entry. The first graphic for a chain summarises the proportions of the various outlier classes displayed in the second graphic. The second graphic shows the sequence view annotated by issues in geometry and electron density. Residues are color-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. A red dot above a residue indicates a poor fit to the electron density (RSRZ > 2). Stretches of 2 or more consecutive residues without any outlier are shown as a green connector. Residues present in the sample, but not in the model, are shown in grey.

• Molecule 1: designed protein OR327

Chain A: 48% 34% 10% 7%



• Molecule 1: designed protein OR327

Chain B: 51% 34% 8% • 7%

• Molecule 1: designed protein OR327

Chain C: 3% 48% 37% 8% 7%




I152
T153
V158
Q161
L162
GLY
SER
LEU
GLU
HIS
HIS
HIS
HIS
HIS
HIS

4 Data and refinement statistics (i)

Property	Value	Source
Space group	P 21 21 21	Depositor
Cell constants a, b, c, α, β, γ	61.21Å 71.85Å 118.01Å 90.00° 90.00° 90.00°	Depositor
Resolution (Å)	28.16 – 2.96 28.16 – 2.96	Depositor EDS
% Data completeness (in resolution range)	98.1 (28.16-2.96) 98.1 (28.16-2.96)	Depositor EDS
R_{merge}	0.10	Depositor
R_{sym}	(Not available)	Depositor
$< I/\sigma(I) >$ [1]	1.93 (at 2.95Å)	Xtriage
Refinement program	PHENIX dev_1269	Depositor
R, R_{free}	0.187 , 0.271 0.189 , 0.267	Depositor DCC
R_{free} test set	554 reflections (4.99%)	wwPDB-VP
Wilson B-factor (Å^2)	75.8	Xtriage
Anisotropy	0.489	Xtriage
Bulk solvent k_{sol}(e/Å^3), B_{sol}(Å^2)	0.24 , 58.4	EDS
L-test for twinning[2]	$< \|L\| > = 0.45$, $< L^2 > = 0.27$	Xtriage
Estimated twinning fraction	No twinning to report.	Xtriage
F_o,F_c correlation	0.95	EDS
Total number of atoms	3839	wwPDB-VP
Average B, all atoms (Å^2)	94.0	wwPDB-VP

Xtriage's analysis on translational NCS is as follows: *The largest off-origin peak in the Patterson function is 7.21% of the height of the origin peak. No significant pseudotranslation is detected.*

[1]Intensities estimated from amplitudes.

[2]Theoretical values of $< |L| >$, $< L^2 >$ for acentric reflections are 0.5, 0.333 respectively for untwinned datasets, and 0.375, 0.2 for perfectly twinned datasets.

5 Model quality (i)

5.1 Standard geometry (i)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	$\#\|Z\| >5$	RMSZ	$\#\|Z\| >5$
1	A	0.50	0/1290	0.72	0/1731
1	B	0.50	0/1286	0.75	0/1725
1	C	0.41	0/1290	0.62	0/1731
All	All	0.47	0/3866	0.70	0/5187

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand.A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	#Chirality outliers	#Planarity outliers
1	B	0	1

There are no bond length outliers.

There are no bond angle outliers.

There are no chirality outliers.

All (1) planarity outliers are listed below:

Mol	Chain	Res	Type	Group
1	B	152	ILE	Peptide

5.2 Too-close contacts (i)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in the chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes within the asymmetric unit, whereas Symm-Clashes lists symmetry-related clashes.

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	A	1281	0	1317	50	0
1	B	1277	0	1316	40	0

Continued on next page...

Continued from previous page...

Mol	Chain	Non-H	H(model)	H(added)	Clashes	Symm-Clashes
1	C	1281	0	1317	44	0
All	All	3839	0	3950	127	0

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 16.

All (127) close contacts within the same asymmetric unit are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:99:LEU:HB2	1:A:122:LEU:HD21	1.56	0.86
1:A:140:GLU:OE2	1:A:143:ARG:NH1	2.09	0.86
1:A:18:ASP:HB2	1:C:18:ASP:HB2	1.61	0.81
1:B:106:ALA:HA	1:B:137:LEU:HD21	1.69	0.75
1:B:25:THR:OG1	1:B:26:GLU:N	2.19	0.73
1:B:15:LYS:HB3	1:C:115:TYR:CZ	2.26	0.70
1:A:47:ARG:HD3	1:A:161:GLN:HG2	1.72	0.70
1:B:144:LEU:HD23	1:B:148:PHE:HE2	1.56	0.70
1:C:95:LEU:HD11	1:C:120:ASN:HA	1.74	0.70
1:A:89:GLY:O	1:A:120:ASN:HB2	1.92	0.69
1:A:49:ARG:HG3	1:A:159:GLU:HB3	1.75	0.69
1:B:99:LEU:HB2	1:B:122:LEU:HD21	1.75	0.67
1:C:76:TYR:HA	1:C:162:LEU:HD12	1.77	0.67
1:B:102:MSE:HE1	1:B:141:ALA:HB2	1.78	0.66
1:C:102:MSE:HA	1:C:102:MSE:HE2	1.76	0.65
1:B:67:LEU:HD13	1:B:158:VAL:HG12	1.79	0.65
1:A:80:ASN:HB3	1:A:159:GLU:HG2	1.77	0.65
1:C:90:ASP:OD2	1:C:120:ASN:ND2	2.31	0.63
1:A:130:PRO:HG2	1:A:133:VAL:HG23	1.81	0.61
1:C:5:GLN:HE21	1:C:20:THR:HG22	1.65	0.61
1:A:102:MSE:HA	1:A:102:MSE:HE2	1.80	0.61
1:C:30:GLN:O	1:C:34:ASN:ND2	2.26	0.60
1:C:140:GLU:OE2	1:C:143:ARG:NH1	2.36	0.59
1:C:12:ASP:OD1	1:C:46:LYS:N	2.34	0.58
1:C:143:ARG:HA	1:C:146:LYS:HB3	1.84	0.58
1:C:72:ALA:O	1:C:75:GLY:N	2.28	0.57
1:B:142:GLU:HA	1:B:152:ILE:HD12	1.87	0.57
1:A:92:LEU:HA	1:A:95:LEU:HD12	1.87	0.57
1:B:99:LEU:HA	1:B:102:MSE:HB3	1.88	0.56
1:C:99:LEU:O	1:C:103:ILE:HG12	2.07	0.55
1:C:117:LEU:HB2	1:C:122:LEU:HD13	1.89	0.55
1:C:86:ARG:HB2	1:C:153:THR:HB	1.89	0.55

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:C:131:ARG:NH1	1:C:135:GLU:OE2	2.40	0.54
1:C:90:ASP:HA	1:C:120:ASN:HB3	1.89	0.54
1:B:109:PHE:O	1:B:111:GLY:N	2.32	0.54
1:A:82:LYS:HB3	1:A:157:THR:HB	1.88	0.54
1:B:55:ARG:HH21	1:B:153:THR:HG22	1.72	0.54
1:A:38:ASP:O	1:A:42:LYS:HG3	2.08	0.54
1:B:49:ARG:HG3	1:B:159:GLU:HB2	1.90	0.54
1:B:6:VAL:HG22	1:B:52:ILE:HG12	1.90	0.53
1:C:143:ARG:O	1:C:147:GLU:N	2.41	0.53
1:B:144:LEU:HD23	1:B:148:PHE:CE2	2.39	0.53
1:B:117:LEU:HD12	1:B:121:ASP:O	2.10	0.52
1:C:62:LYS:O	1:C:66:ILE:HG13	2.09	0.52
1:C:7:GLN:HG3	1:C:20:THR:HG23	1.92	0.52
1:A:26:GLU:OE2	1:A:62:LYS:NZ	2.25	0.52
1:A:54:ALA:HB1	1:A:59:GLU:HB3	1.92	0.52
1:B:98:ALA:HA	1:B:144:LEU:HD22	1.91	0.51
1:C:83:MSE:HE1	1:C:137:LEU:HB2	1.92	0.51
1:B:46:LYS:O	1:B:161:GLN:HG2	2.10	0.51
1:A:119:GLY:O	1:A:121:ASP:N	2.43	0.51
1:A:127:THR:HG22	1:A:128:GLY:N	2.26	0.51
1:B:81:ARG:HG2	1:B:158:VAL:HG23	1.91	0.51
1:A:117:LEU:HB2	1:A:122:LEU:HD13	1.93	0.51
1:A:131:ARG:O	1:A:135:GLU:HG3	2.11	0.51
1:A:65:ALA:HB2	1:A:131:ARG:HH21	1.74	0.51
1:C:130:PRO:HG2	1:C:133:VAL:HG23	1.92	0.51
1:B:39:TYR:CD2	1:B:43:GLN:HG3	2.46	0.50
1:C:88:ARG:HB2	1:C:151:THR:HB	1.93	0.50
1:C:108:LYS:HB3	1:C:109:PHE:CE2	2.47	0.50
1:C:69:LYS:NZ	1:C:73:GLU:OE2	2.45	0.49
1:A:3:ASP:HB3	1:A:4:ILE:HD12	1.92	0.49
1:A:25:THR:OG1	1:A:28:GLU:HG3	2.13	0.49
1:B:85:VAL:HG13	1:B:154:ILE:HG12	1.95	0.48
1:A:102:MSE:HG2	1:A:144:LEU:HD23	1.95	0.48
1:B:82:LYS:HG3	1:B:127:THR:HG23	1.95	0.48
1:C:28:GLU:O	1:C:32:VAL:HG13	2.13	0.48
1:A:127:THR:HG22	1:A:128:GLY:H	1.78	0.48
1:B:84:THR:HA	1:B:124:ILE:O	2.15	0.47
1:B:140:GLU:O	1:B:144:LEU:HD12	2.14	0.47
1:B:95:LEU:HD23	1:B:95:LEU:HA	1.63	0.47
1:A:40:ILE:HG23	1:A:45:ALA:H	1.80	0.47
1:A:126:ILE:HD13	1:A:137:LEU:HD13	1.97	0.47

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:18:ASP:HB2	1:C:18:ASP:CB	2.38	0.46
1:B:47:ARG:HE	1:B:47:ARG:HB3	1.57	0.46
1:C:78:ASP:HB3	1:C:161:GLN:O	2.16	0.46
1:A:102:MSE:HG2	1:A:144:LEU:CD2	2.46	0.45
1:C:5:GLN:HB3	1:C:53:THR:HB	1.98	0.45
1:A:74:LEU:HB3	1:A:76:TYR:CD1	2.51	0.45
1:A:103:ILE:HD12	1:A:107:ARG:HH12	1.83	0.44
1:C:95:LEU:HD22	1:C:122:LEU:HB2	1.99	0.44
1:A:14:GLY:HA2	1:C:22:THR:HB	2.00	0.44
1:A:162:LEU:HD23	1:A:162:LEU:HA	1.85	0.44
1:B:88:ARG:O	1:B:150:ILE:HG22	2.18	0.44
1:C:25:THR:C	1:C:27:SER:H	2.22	0.44
1:A:66:ILE:O	1:A:70:VAL:HG23	2.18	0.43
1:B:57:SER:OG	1:B:139:LYS:HD2	2.18	0.43
1:A:92:LEU:HD11	1:B:107:ARG:NE	2.33	0.43
1:C:92:LEU:HA	1:C:95:LEU:HD12	2.01	0.43
1:C:102:MSE:HE1	1:C:140:GLU:HB3	2.01	0.43
1:B:113:VAL:HA	1:B:125:THR:O	2.18	0.43
1:A:6:VAL:HB	1:A:21:TYR:HB2	2.01	0.43
1:A:36:LEU:O	1:A:39:TYR:HB3	2.18	0.43
1:A:107:ARG:NH2	1:B:96:GLU:OE1	2.51	0.43
1:B:142:GLU:O	1:B:146:LYS:HG2	2.18	0.43
1:A:49:ARG:HA	1:A:159:GLU:HA	2.01	0.42
1:A:96:GLU:O	1:A:99:LEU:N	2.52	0.42
1:C:72:ALA:O	1:C:74:LEU:N	2.52	0.42
1:A:67:LEU:HD23	1:A:67:LEU:HA	1.84	0.42
1:A:34:ASN:HA	1:A:37:MSE:HE3	2.02	0.42
1:B:27:SER:O	1:B:31:LYS:HB3	2.19	0.42
1:B:63:PHE:O	1:B:67:LEU:HG	2.19	0.42
1:C:91:ASP:OD1	1:C:93:GLU:HB2	2.19	0.42
1:A:143:ARG:NH2	1:A:147:GLU:OE1	2.45	0.42
1:C:26:GLU:O	1:C:29:LEU:HB3	2.20	0.42
1:C:87:PHE:HA	1:C:151:THR:O	2.19	0.42
1:A:118:ASP:HA	1:B:114:THR:HG22	2.02	0.42
1:C:40:ILE:HG23	1:C:45:ALA:N	2.35	0.41
1:A:62:LYS:HG2	1:A:66:ILE:HD11	2.03	0.41
1:A:62:LYS:O	1:A:65:ALA:HB3	2.20	0.41
1:A:64:LEU:HA	1:A:67:LEU:HB2	2.02	0.41
1:B:12:ASP:O	1:B:13:ASN:HB2	2.20	0.41
1:B:66:ILE:O	1:B:70:VAL:HG23	2.21	0.41
1:A:23:VAL:HG21	1:A:29:LEU:HB2	2.03	0.41

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Interatomic distance (Å)	Clash overlap (Å)
1:A:56:SER:HB2	1:A:59:GLU:HB2	2.03	0.41
1:B:83:MSE:HE2	1:B:83:MSE:HB3	1.88	0.41
1:A:15:LYS:HB2	1:A:15:LYS:HE2	1.81	0.41
1:B:74:LEU:HD12	1:B:74:LEU:HA	1.90	0.41
1:C:11:ASP:OD1	1:C:16:ASN:HA	2.21	0.41
1:C:34:ASN:HA	1:C:37:MSE:HG3	2.03	0.41
1:A:23:VAL:HB	1:A:28:GLU:HB2	2.03	0.41
1:B:134:LEU:HD12	1:B:134:LEU:HA	1.90	0.41
1:C:106:ALA:O	1:C:111:GLY:HA3	2.20	0.41
1:C:109:PHE:C	1:C:111:GLY:H	2.24	0.40
1:C:13:ASN:N	1:C:13:ASN:OD1	2.54	0.40
1:B:71:PHE:CD2	1:B:79:ILE:HD11	2.56	0.40
1:A:28:GLU:O	1:A:31:LYS:N	2.53	0.40

There are no symmetry-related clashes.

5.3 Torsion angles (i)

5.3.1 Protein backbone (i)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the backbone conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	158/172 (92%)	137 (87%)	14 (9%)	7 (4%)	2 12
1	B	158/172 (92%)	139 (88%)	15 (10%)	4 (2%)	5 25
1	C	158/172 (92%)	131 (83%)	22 (14%)	5 (3%)	4 19
All	All	474/516 (92%)	407 (86%)	51 (11%)	16 (3%)	3 17

All (16) Ramachandran outliers are listed below:

Mol	Chain	Res	Type
1	A	120	ASN
1	C	81	ARG
1	C	120	ASN
1	A	119	GLY

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	42	LYS
1	A	158	VAL
1	B	110	ALA
1	B	130	PRO
1	C	80	ASN
1	C	111	GLY
1	A	13	ASN
1	A	82	LYS
1	B	13	ASN
1	B	98	ALA
1	C	73	GLU
1	A	44	GLY

5.3.2 Protein sidechains (i)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all X-ray entries followed by that with respect to entries of similar resolution.

The Analysed column shows the number of residues for which the sidechain conformation was analysed, and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles
1	A	139/145 (96%)	117 (84%)	22 (16%)	2 10
1	B	138/145 (95%)	117 (85%)	21 (15%)	3 12
1	C	139/145 (96%)	118 (85%)	21 (15%)	3 12
All	All	416/435 (96%)	352 (85%)	64 (15%)	2 11

All (64) residues with a non-rotameric sidechain are listed below:

Mol	Chain	Res	Type
1	A	3	ASP
1	A	7	GLN
1	A	13	ASN
1	A	15	LYS
1	A	18	ASP
1	A	24	THR
1	A	26	GLU
1	A	47	ARG
1	A	58	LYS
1	A	64	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	A	74	LEU
1	A	83	MSE
1	A	92	LEU
1	A	102	MSE
1	A	125	THR
1	A	129	VAL
1	A	134	LEU
1	A	143	ARG
1	A	144	LEU
1	A	153	THR
1	A	158	VAL
1	A	159	GLU
1	B	9	ASN
1	B	16	ASN
1	B	18	ASP
1	B	19	TYR
1	B	25	THR
1	B	28	GLU
1	B	31	LYS
1	B	35	GLU
1	B	47	ARG
1	B	55	ARG
1	B	74	LEU
1	B	79	ILE
1	B	121	ASP
1	B	127	THR
1	B	129	VAL
1	B	134	LEU
1	B	137	LEU
1	B	143	ARG
1	B	150	ILE
1	B	152	ILE
1	B	156	VAL
1	C	3	ASP
1	C	13	ASN
1	C	16	ASN
1	C	26	GLU
1	C	31	LYS
1	C	32	VAL
1	C	53	THR
1	C	90	ASP
1	C	92	LEU

Continued on next page...

Continued from previous page...

Mol	Chain	Res	Type
1	C	96	GLU
1	C	113	VAL
1	C	114	THR
1	C	117	LEU
1	C	123	GLU
1	C	127	THR
1	C	129	VAL
1	C	134	LEU
1	C	143	ARG
1	C	149	ASN
1	C	151	THR
1	C	158	VAL

Sometimes sidechains can be flipped to improve hydrogen bonding and reduce clashes. All (2) such sidechains are listed below:

Mol	Chain	Res	Type
1	C	5	GLN
1	C	120	ASN

5.3.3 RNA (i)

There are no RNA molecules in this entry.

5.4 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

5.5 Carbohydrates (i)

There are no monosaccharides in this entry.

5.6 Ligand geometry (i)

There are no ligands in this entry.

5.7 Other polymers (i)

There are no such residues in this entry.

5.8 Polymer linkage issues (i)

There are no chain breaks in this entry.

6 Fit of model and data (i)

6.1 Protein, DNA and RNA chains (i)

In the following table, the column labelled '#RSRZ> 2' contains the number (and percentage) of RSRZ outliers, followed by percent RSRZ outliers for the chain as percentile scores relative to all X-ray entries and entries of similar resolution. The OWAB column contains the minimum, median, 95th percentile and maximum values of the occupancy-weighted average B-factor per residue. The column labelled 'Q< 0.9' lists the number of (and percentage) of residues with an average occupancy less than 0.9.

Mol	Chain	Analysed	<RSRZ>	#RSRZ>2			OWAB(Å^2)	Q<0.9
1	A	157/172 (91%)	-0.42	0	100	100	44, 83, 118, 180	0
1	B	157/172 (91%)	-0.49	0	100	100	46, 79, 118, 194	0
1	C	157/172 (91%)	-0.05	5 (3%)	47	31	61, 107, 172, 222	0
All	All	471/516 (91%)	-0.32	5 (1%)	80	65	44, 88, 145, 222	0

All (5) RSRZ outliers are listed below:

Mol	Chain	Res	Type	RSRZ
1	C	89	GLY	4.8
1	C	77	ASN	3.3
1	C	162	LEU	3.1
1	C	55	ARG	2.4
1	C	51	SER	2.0

6.2 Non-standard residues in protein, DNA, RNA chains (i)

There are no non-standard protein/DNA/RNA residues in this entry.

6.3 Carbohydrates (i)

There are no monosaccharides in this entry.

6.4 Ligands (i)

There are no ligands in this entry.

6.5 Other polymers (i)

There are no such residues in this entry.