



Full wwPDB NMR Structure Validation Report ⓘ

Feb 19, 2018 – 07:49 pm GMT

PDB ID : 1KZ0
Title : Solution structure of the third helix of Antennapedia homeodomain
Authors : Czajlik, A.; Mesko, E.; Penke, B.; Perczel, A.
Deposited on : 2002-02-06

This is a Full wwPDB NMR Structure Validation Report for a publicly released PDB entry.

We welcome your comments at validation@mail.wwpdb.org

A user guide is available at

<https://www.wwpdb.org/validation/2017/NMRValidationReportHelp>

with specific help available everywhere you see the ⓘ symbol.

The following versions of software and data (see [references ⓘ](#)) were used in the production of this report:

Cyrange : Kirchner and Güntert (2011)
NmrClust : Kelley et al. (1996)
MolProbity : 4.02b-467
Percentile statistics : 20171227.v01 (using entries in the PDB archive December 27th 2017)
RCI : v_1n_11_5_13_A (Berjanski et al., 2005)
PANAV : Wang et al. (2010)
ShiftChecker : trunk30686
Ideal geometry (proteins) : Engh & Huber (2001)
Ideal geometry (DNA, RNA) : Parkinson et al. (1996)
Validation Pipeline (wwPDB-VP) : trunk30686


1 Overall quality at a glance

The following experimental techniques were used to determine the structure:

SOLUTION NMR

The overall completeness of chemical shifts assignment is 49%.

Percentile scores (ranging between 0-100) for global validation metrics of the entry are shown in the following graphic. The table shows the number of entries on which the scores are based.



Metric	Whole archive (#Entries)	NMR archive (#Entries)
Clashscore	136279	12091
Ramachandran outliers	132675	10835
Sidechain outliers	132484	10811

The table below summarises the geometric issues observed across the polymeric chains and their fit to the experimental data. The red, orange, yellow and green segments indicate the fraction of residues that contain outliers for ≥ 3 , 2, 1 and 0 types of geometric quality criteria. A cyan segment indicates the fraction of residues that are not part of the well-defined cores, and a grey segment represents the fraction of residues that are not modelled. The numeric value for each fraction is indicated below the corresponding segment, with a dot representing fractions $\leq 5\%$

Mol	Chain	Length	Quality of chain
1	A	16	

2 Ensemble composition and analysis

This entry contains 26 models. Model 20 is the overall representative, medoid model (most similar to other models). The authors have identified model 1 as representative, based on the following criterion: *fewest violations, lowest energy*.

The following residues are included in the computation of the global validation metrics.

Well-defined (core) protein residues			
Well-defined core	Residue range (total)	Backbone RMSD (Å)	Medoid model
1	A:1-A:16 (16)	0.06	20

Ill-defined regions of proteins are excluded from the global statistics.

Ligands and non-protein polymers are included in the analysis.

The models can be grouped into 4 clusters and 4 single-model clusters were found.

Cluster number	Models
1	2, 5, 7, 8, 9, 10, 16, 17, 18, 20, 21, 22
2	4, 13, 14, 15, 19, 26
3	24, 25
4	1, 12
Single-model clusters	3; 6; 11; 23

3 Entry composition

There is only 1 type of molecule in this entry. The entry contains 334 atoms, of which 175 are hydrogens and 0 are deuteriums.

- Molecule 1 is a protein called Antennapedia protein.

Mol	Chain	Residues	Atoms						Trace
			Total	C	H	N	O	S	
1	A	16	334	104	175	34	20	1	0

4 Residue-property plots

4.1 Average score per residue in the NMR ensemble

These plots are provided for all protein, RNA and DNA chains in the entry. The first graphic is the same as shown in the summary in section 1 of this report. The second graphic shows the sequence where residues are colour-coded according to the number of geometric quality criteria for which they contain at least one outlier: green = 0, yellow = 1, orange = 2 and red = 3 or more. Stretches of 2 or more consecutive residues without any outliers are shown as green connectors. Residues which are classified as ill-defined in the NMR ensemble, are shown in cyan with an underline colour-coded according to the previous scheme. Residues which were present in the experimental sample, but not modelled in the final structure are shown in grey.

- Molecule 1: Antennapedia protein




4.2 Scores per residue for each member of the ensemble

Colouring as in section 4.1 above.


4.2.1 Score per residue for model 1

- Molecule 1: Antennapedia protein



4.2.2 Score per residue for model 2

- Molecule 1: Antennapedia protein



4.2.3 Score per residue for model 3

- Molecule 1: Antennapedia protein



4.2.4 Score per residue for model 4

- Molecule 1: Antennapedia protein




4.2.5 Score per residue for model 5

- Molecule 1: Antennapedia protein




4.2.6 Score per residue for model 6

- Molecule 1: Antennapedia protein




4.2.7 Score per residue for model 7

- Molecule 1: Antennapedia protein




4.2.8 Score per residue for model 8

- Molecule 1: Antennapedia protein



4.2.9 Score per residue for model 9

- Molecule 1: Antennapedia protein




4.2.10 Score per residue for model 10

- Molecule 1: Antennapedia protein




4.2.11 Score per residue for model 11

- Molecule 1: Antennapedia protein



4.2.12 Score per residue for model 12

- Molecule 1: Antennapedia protein




4.2.13 Score per residue for model 13

- Molecule 1: Antennapedia protein



4.2.14 Score per residue for model 14

- Molecule 1: Antennapedia protein



4.2.15 Score per residue for model 15

- Molecule 1: Antennapedia protein




4.2.16 Score per residue for model 16

- Molecule 1: Antennapedia protein



4.2.17 Score per residue for model 17

- Molecule 1: Antennapedia protein



4.2.18 Score per residue for model 18

- Molecule 1: Antennapedia protein



4.2.19 Score per residue for model 19

- Molecule 1: Antennapedia protein



4.2.20 Score per residue for model 20 (medoid)

- Molecule 1: Antennapedia protein



4.2.21 Score per residue for model 21

- Molecule 1: Antennapedia protein




4.2.22 Score per residue for model 22

- Molecule 1: Antennapedia protein




4.2.23 Score per residue for model 23

- Molecule 1: Antennapedia protein



4.2.24 Score per residue for model 24

- Molecule 1: Antennapedia protein



4.2.25 Score per residue for model 25

- Molecule 1: Antennapedia protein



4.2.26 Score per residue for model 26

- Molecule 1: Antennapedia protein



5 Refinement protocol and experimental data overview

The models were refined using the following method: *simulated annealing and molecular dynamics*.

Of the 50 calculated structures, 26 were deposited, based on the following criterion: *structures with the least restraint violations, structures with the lowest energy*.

The following table shows the software used for structure solution, optimisation and refinement.

Software name	Classification	Version
X-PLOR	refinement	3.851
X-PLOR	structure solution	3.851

The following table shows chemical shift validation statistics as aggregates over all chemical shift files. Detailed validation can be found in section 7 of this report.

Chemical shift file(s)	BMRB entry 5289
Number of chemical shift lists	2
Total number of shifts	294
Number of shifts mapped to atoms	294
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Assignment completeness (well-defined parts)	49%

No validations of the models with respect to experimental NMR restraints is performed at this time.

6 Model quality [i](#)

6.1 Standard geometry [i](#)

The Z score for a bond length (or angle) is the number of standard deviations the observed value is removed from the expected value. A bond length (or angle) with $|Z| > 5$ is considered an outlier worth inspection. RMSZ is the (average) root-mean-square of all Z scores of the bond lengths (or angles).

Mol	Chain	Bond lengths		Bond angles	
		RMSZ	#Z>5	RMSZ	#Z>5
1	A	0.98±0.00	0±0/163 (0.0±0.0%)	1.61±0.01	0±0/212 (0.1±0.2%)
All	All	0.98	0/4238 (0.0%)	1.61	7/5512 (0.1%)

Chiral center outliers are detected by calculating the chiral volume of a chiral center and verifying if the center is modelled as a planar moiety or with the opposite hand. A planarity outlier is detected by checking planarity of atoms in a peptide group, atoms in a mainchain group or atoms of a sidechain that are expected to be planar.

Mol	Chain	Chirality	Planarity
1	A	0.0±0.0	3.0±0.0
All	All	0	78

There are no bond-length outliers.

All unique angle outliers are listed below.

Mol	Chain	Res	Type	Atoms	Z	Observed(°)	Ideal(°)	Models	
								Worst	Total
1	A	14	TRP	NE1-CE2-CZ2	5.09	136.00	130.40	1	7

There are no chirality outliers.

All unique planar outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Group	Models (Total)
1	A	1	ARG	Sidechain	26
1	A	11	ARG	Sidechain	26
1	A	10	ARG	Sidechain	26

6.2 Too-close contacts [i](#)

In the following table, the Non-H and H(model) columns list the number of non-hydrogen atoms and hydrogen atoms in each chain respectively. The H(added) column lists the number of hydrogen atoms added and optimized by MolProbity. The Clashes column lists the number of clashes averaged over the ensemble.

Mol	Chain	Non-H	H(model)	H(added)	Clashes
1	A	159	175	175	52±2
All	All	4134	4550	4550	1362

The all-atom clashscore is defined as the number of clashes found per 1000 atoms (including hydrogen atoms). The all-atom clashscore for this structure is 157.

All unique clashes are listed below, sorted by their clash magnitude.

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:3:ILE:HG22	1:A:3:ILE:O	0.83	1.73	6	15
1:A:3:ILE:O	1:A:3:ILE:HG22	0.82	1.73	7	11
1:A:3:ILE:HA	1:A:6:TRP:CD1	0.77	2.14	11	26
1:A:5:ILE:C	1:A:5:ILE:HD12	0.77	2.01	16	14
1:A:5:ILE:HD12	1:A:5:ILE:C	0.77	2.00	10	12
1:A:7:PHE:O	1:A:11:ARG:HB2	0.72	1.85	11	26
1:A:5:ILE:HD12	1:A:6:TRP:N	0.72	2.00	16	26
1:A:3:ILE:HA	1:A:6:TRP:CG	0.71	2.19	11	26
1:A:3:ILE:CG2	1:A:3:ILE:O	0.70	2.40	11	15
1:A:3:ILE:O	1:A:3:ILE:CG2	0.70	2.39	14	11
1:A:6:TRP:CE2	1:A:7:PHE:CD1	0.69	2.80	26	26
1:A:15:LYS:CD	1:A:15:LYS:N	0.68	2.56	26	10
1:A:9:ASN:O	1:A:12:MET:HG2	0.67	1.90	7	25
1:A:15:LYS:N	1:A:15:LYS:CD	0.66	2.58	13	16
1:A:3:ILE:HA	1:A:6:TRP:CD2	0.66	2.26	11	26
1:A:6:TRP:CG	1:A:7:PHE:N	0.66	2.63	11	26
1:A:3:ILE:HA	1:A:6:TRP:CE2	0.65	2.26	11	26
1:A:6:TRP:CD1	1:A:6:TRP:N	0.63	2.67	11	15
1:A:7:PHE:O	1:A:10:ARG:HG2	0.62	1.94	10	26
1:A:6:TRP:N	1:A:6:TRP:CD1	0.62	2.68	15	11
1:A:7:PHE:O	1:A:11:ARG:N	0.62	2.32	26	26
1:A:3:ILE:CA	1:A:6:TRP:CD1	0.62	2.82	11	26
1:A:7:PHE:CD1	1:A:8:GLN:N	0.61	2.68	23	26
1:A:6:TRP:NE1	1:A:7:PHE:CE1	0.61	2.68	26	26
1:A:11:ARG:O	1:A:14:TRP:HB2	0.60	1.96	6	26
1:A:3:ILE:C	1:A:6:TRP:CD1	0.60	2.75	11	26
1:A:7:PHE:HD1	1:A:8:GLN:N	0.58	1.96	11	26

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:3:ILE:HA	1:A:6:TRP:NE1	0.57	2.13	3	26
1:A:12:MET:CG	1:A:13:LYS:HE3	0.57	2.29	6	1
1:A:6:TRP:NE1	1:A:7:PHE:CD1	0.57	2.72	11	26
1:A:9:ASN:O	1:A:13:LYS:HG2	0.57	1.99	25	26
1:A:12:MET:HG3	1:A:13:LYS:HE3	0.56	1.76	23	1
1:A:14:TRP:CE3	1:A:16:LYS:CD	0.55	2.90	19	24
1:A:5:ILE:CD1	1:A:5:ILE:C	0.55	2.75	10	13
1:A:5:ILE:C	1:A:5:ILE:CD1	0.54	2.75	25	13
1:A:7:PHE:C	1:A:7:PHE:CD1	0.54	2.80	10	13
1:A:7:PHE:CD1	1:A:7:PHE:C	0.54	2.80	6	13
1:A:8:GLN:HA	1:A:11:ARG:HB2	0.54	1.79	1	26
1:A:9:ASN:O	1:A:12:MET:CG	0.54	2.56	23	4
1:A:14:TRP:CZ3	1:A:16:LYS:HD2	0.52	2.40	2	16
1:A:15:LYS:HD2	1:A:15:LYS:N	0.51	2.21	26	9
1:A:8:GLN:O	1:A:8:GLN:NE2	0.50	2.44	23	12
1:A:14:TRP:CE3	1:A:16:LYS:HD3	0.50	2.41	24	18
1:A:3:ILE:HG23	1:A:6:TRP:CZ2	0.50	2.41	16	26
1:A:3:ILE:CD1	1:A:3:ILE:N	0.50	2.75	25	19
1:A:6:TRP:CZ2	1:A:7:PHE:CD2	0.50	3.00	11	26
1:A:3:ILE:N	1:A:3:ILE:CD1	0.50	2.75	2	7
1:A:2:GLN:O	1:A:3:ILE:HB	0.50	2.06	11	26
1:A:7:PHE:CZ	1:A:11:ARG:NH2	0.49	2.81	17	6
1:A:8:GLN:NE2	1:A:8:GLN:O	0.48	2.46	9	14
1:A:4:LYS:N	1:A:6:TRP:CD1	0.48	2.82	17	26
1:A:9:ASN:HA	1:A:12:MET:HG2	0.48	1.85	23	1
1:A:3:ILE:C	1:A:6:TRP:NE1	0.48	2.67	16	26
1:A:15:LYS:N	1:A:15:LYS:HD2	0.47	2.23	1	17
1:A:14:TRP:CZ3	1:A:16:LYS:CD	0.47	2.97	4	7
1:A:3:ILE:HG13	1:A:6:TRP:CZ2	0.47	2.44	26	26
1:A:6:TRP:CE2	1:A:7:PHE:CG	0.47	3.02	11	26
1:A:7:PHE:CZ	1:A:11:ARG:CZ	0.47	2.98	26	19
1:A:9:ASN:O	1:A:12:MET:HE2	0.47	2.10	7	1
1:A:7:PHE:CZ	1:A:11:ARG:NH1	0.47	2.83	18	20
1:A:6:TRP:CZ2	1:A:7:PHE:CE2	0.46	3.03	11	16
1:A:3:ILE:CA	1:A:6:TRP:NE1	0.46	2.78	16	26
1:A:12:MET:HG3	1:A:13:LYS:N	0.46	2.24	22	3
1:A:9:ASN:C	1:A:12:MET:HG2	0.46	2.31	9	4
1:A:6:TRP:CE2	1:A:7:PHE:CE1	0.46	3.03	26	26
1:A:12:MET:HG3	1:A:13:LYS:CE	0.45	2.42	6	1
1:A:9:ASN:CA	1:A:12:MET:HG2	0.45	2.41	23	1
1:A:7:PHE:CE1	1:A:11:ARG:CZ	0.44	3.00	3	16

Continued on next page...

Continued from previous page...

Atom-1	Atom-2	Clash(Å)	Distance(Å)	Models	
				Worst	Total
1:A:2:GLN:C	1:A:4:LYS:HD3	0.44	2.33	7	25
1:A:6:TRP:O	1:A:9:ASN:N	0.43	2.51	23	26
1:A:2:GLN:O	1:A:4:LYS:CD	0.42	2.67	4	25
1:A:5:ILE:O	1:A:9:ASN:N	0.42	2.49	22	26
1:A:6:TRP:C	1:A:8:GLN:N	0.42	2.73	26	26
1:A:11:ARG:O	1:A:12:MET:C	0.42	2.58	23	1
1:A:5:ILE:O	1:A:9:ASN:ND2	0.42	2.47	23	1
1:A:12:MET:HG3	1:A:13:LYS:CD	0.41	2.46	18	8
1:A:12:MET:CE	1:A:13:LYS:HD3	0.41	2.46	8	1
1:A:12:MET:CG	1:A:13:LYS:N	0.41	2.83	26	6
1:A:8:GLN:CA	1:A:11:ARG:HB2	0.41	2.44	1	16
1:A:9:ASN:HA	1:A:12:MET:HE3	0.41	1.92	25	1
1:A:14:TRP:C	1:A:15:LYS:HD2	0.41	2.36	26	1
1:A:4:LYS:N	1:A:4:LYS:CD	0.40	2.85	11	1

6.3 Torsion angles [i](#)

6.3.1 Protein backbone [i](#)

In the following table, the Percentiles column shows the percent Ramachandran outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR entries. The Analysed column shows the number of residues for which the backbone conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Favoured	Allowed	Outliers	Percentiles
1	A	14/16 (88%)	8±0 (57±0%)	4±0 (28±2%)	2±0 (15±2%)	1 5
All	All	364/416 (88%)	208 (57%)	103 (28%)	53 (15%)	1 5

All 3 unique Ramachandran outliers are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	6	TRP	26
1	A	3	ILE	25
1	A	13	LYS	2

6.3.2 Protein sidechains [i](#)

In the following table, the Percentiles column shows the percent sidechain outliers of the chain as a percentile score with respect to all PDB entries followed by that with respect to all NMR

entries. The Analysed column shows the number of residues for which the sidechain conformation was analysed and the total number of residues.

Mol	Chain	Analysed	Rotameric	Outliers	Percentiles	
1	A	16/16 (100%)	10±1 (60±4%)	6±1 (40±4%)	0	5
All	All	416/416 (100%)	251 (60%)	165 (40%)	0	5

All 7 unique residues with a non-rotameric sidechain are listed below. They are sorted by the frequency of occurrence in the ensemble.

Mol	Chain	Res	Type	Models (Total)
1	A	4	LYS	26
1	A	14	TRP	26
1	A	15	LYS	26
1	A	7	PHE	26
1	A	13	LYS	25
1	A	16	LYS	19
1	A	10	ARG	17

6.3.3 RNA [i](#)

There are no RNA molecules in this entry.

6.4 Non-standard residues in protein, DNA, RNA chains [i](#)

There are no non-standard protein/DNA/RNA residues in this entry.

6.5 Carbohydrates [i](#)

There are no carbohydrates in this entry.

6.6 Ligand geometry [i](#)

There are no ligands in this entry.

6.7 Other polymers [i](#)

There are no such molecules in this entry.

6.8 Polymer linkage issues

There are no chain breaks in this entry.

7 Chemical shift validation [i](#)

The completeness of assignment taking into account all chemical shift lists is 49% for the well-defined parts and 49% for the entire structure.

7.1 Chemical shift list 1

File name: BMRB entry 5289

Chemical shift list name: *assigned_chem_shift_list_1*

7.1.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	154
Number of shifts mapped to atoms	154
Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.1.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.1.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 49%, i.e. 138 atoms were assigned a chemical shift out of a possible 281. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	31/80 (39%)	31/32 (97%)	0/32 (0%)	0/16 (0%)
Sidechain	90/168 (54%)	90/102 (88%)	0/50 (0%)	0/16 (0%)
Aromatic	17/33 (52%)	17/17 (100%)	0/14 (0%)	0/2 (0%)
Overall	138/281 (49%)	138/151 (91%)	0/96 (0%)	0/34 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 49%, i.e. 138 atoms were assigned a chemical shift out of a possible 281. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	31/80 (39%)	31/32 (97%)	0/32 (0%)	0/16 (0%)
Sidechain	90/168 (54%)	90/102 (88%)	0/50 (0%)	0/16 (0%)
Aromatic	17/33 (52%)	17/17 (100%)	0/14 (0%)	0/2 (0%)
Overall	138/281 (49%)	138/151 (91%)	0/96 (0%)	0/34 (0%)


7.1.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.1.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:



7.2 Chemical shift list 2

File name: BMRB entry 5289

Chemical shift list name: *assigned_chem_shift_list_2*

7.2.1 Bookkeeping [i](#)

The following table shows the results of parsing the chemical shift list and reports the number of nuclei with statistically unusual chemical shifts.

Total number of shifts	140
Number of shifts mapped to atoms	140

Number of unparsed shifts	0
Number of shifts with mapping errors	0
Number of shifts with mapping warnings	0
Number of shift outliers (ShiftChecker)	0

7.2.2 Chemical shift referencing [i](#)

No chemical shift referencing corrections were calculated (not enough data).

7.2.3 Completeness of resonance assignments [i](#)

The following table shows the completeness of the chemical shift assignments for the well-defined regions of the structure. The overall completeness is 44%, i.e. 123 atoms were assigned a chemical shift out of a possible 281. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	32/80 (40%)	32/32 (100%)	0/32 (0%)	0/16 (0%)
Sidechain	85/168 (51%)	85/102 (83%)	0/50 (0%)	0/16 (0%)
Aromatic	6/33 (18%)	6/17 (35%)	0/14 (0%)	0/2 (0%)
Overall	123/281 (44%)	123/151 (81%)	0/96 (0%)	0/34 (0%)

The following table shows the completeness of the chemical shift assignments for the full structure. The overall completeness is 44%, i.e. 123 atoms were assigned a chemical shift out of a possible 281. 0 out of 0 assigned methyl groups (LEU and VAL) were assigned stereospecifically.

	Total	¹ H	¹³ C	¹⁵ N
Backbone	32/80 (40%)	32/32 (100%)	0/32 (0%)	0/16 (0%)
Sidechain	85/168 (51%)	85/102 (83%)	0/50 (0%)	0/16 (0%)
Aromatic	6/33 (18%)	6/17 (35%)	0/14 (0%)	0/2 (0%)
Overall	123/281 (44%)	123/151 (81%)	0/96 (0%)	0/34 (0%)

7.2.4 Statistically unusual chemical shifts [i](#)

There are no statistically unusual chemical shifts.

7.2.5 Random Coil Index (RCI) plots [i](#)

The image below reports *random coil index* values for the protein chains in the structure. The height of each bar gives a probability of a given residue to be disordered, as predicted from the available chemical shifts and the amino acid sequence. A value above 0.2 is an indication

of significant predicted disorder. The colour of the bar shows whether the residue is in the well-defined core (black) or in the ill-defined residue ranges (cyan), as described in section 2 on ensemble composition.

Random coil index (RCI) for chain A:

